

THE KNESER-POULSEN CONJECTURE FOR SPHERICAL POLYTOPES

KÁROLY BEZDEK AND ROBERT CONNELLY

Eötvös University and Cornell University

January 14, 2003

ABSTRACT. If a finite set of balls of radius $\pi/2$ (hemispheres) in the unit sphere \mathbb{S}^n is rearranged so that the distance between each pair of centers does not decrease, then the (spherical) volume of the intersection does not increase, and the (spherical) volume of the union does not decrease. This result is a spherical analog to a conjecture by Kneser (1954) and Poulsen (1955) in the case when the radii are all equal to $\pi/2$.

1. Introduction.

Let \mathbb{S}^n be the unit sphere in Euclidean $(n + 1)$ -dimensional space, and let $X(\mathbf{p})$ be a finite intersection of balls of radius $\pi/2$ (closed hemispheres) in \mathbb{S}^n whose configuration of centers is $\mathbf{p} = (\mathbf{p}_1, \dots, \mathbf{p}_N)$. We say that another configuration $\mathbf{q} = (\mathbf{q}_1, \dots, \mathbf{q}_N)$ is a *contraction* of \mathbf{p} if, for all $1 \leq i < j \leq N$, the spherical distance between \mathbf{p}_i and \mathbf{p}_j is not less than the spherical distance between \mathbf{q}_i and \mathbf{q}_j . We denote n -dimensional spherical volume by $\text{Vol}_n[\]$. Our main result is the following.

Theorem 1. *If \mathbf{q} is a configuration in \mathbb{S}^n that is a contraction of the configuration \mathbf{p} , then*

$$\text{Vol}_n[X(\mathbf{p})] \leq \text{Vol}_n[X(\mathbf{q})].$$

This result is to be compared with our previous result [2], where the ambient space is Euclidean space \mathbb{E}^n , and the set X is the intersection of balls of arbitrary radius, but for $n \geq 3$ we need to assume that there is a piecewise-analytic monotone motion of the configuration in \mathbb{E}^{n+2} . This uses the results [4] and [5] of Csikós, where the ambient space \mathbb{E}^n , and the balls which make up the space X are of arbitrary radius, but there must be a continuous contraction of the configuration \mathbf{p} to the configuration \mathbf{q} in \mathbb{E}^n . For a history of earlier related results and the conjectures of Kneser [9] and Poulsen [10] as well as related conjectures, see our paper [2].

1991 *Mathematics Subject Classification.* 52A10, 52A40 (52A38, 51M25, 51M16).

The first author was partially supported by the Hung. Nat. Sci. Found. (OTKA), grant no. T029786. The second author was partially supported as an Erdős Visiting Professor at Eötvös University, Budapest, Hungary, and the National Science Foundation Grant number DMS-0209595.

The method we use here follows the same general outline as in our paper [2]. We use a leapfrog lemma to move one configuration to the other in an analytic and monotone way, but only in higher dimensions. Then the higher-dimensional balls have their combined volume (their intersections or unions) change monotonically, a fact that we prove using Schöffli differential formula. Then we apply an integral formula to relate the volume of the higher dimensional object (or the volume of the boundary of the higher dimension set in our previous case) to the volume of the lower-dimensional object, obtaining the volume inequality for the more general discrete motions.

The following are some corollaries following from Theorem 1 and its proof. The first one is an immediate set theoretic consequence.

Corollary 1. *If a finite set of balls of radius $\pi/2$ in the unit sphere \mathbb{S}^n is rearranged so that the distance between each pair of centers does not increase, then the (spherical) volume of the union does not increase.*

Corollary 2. *Let $\mathbf{p} = (\mathbf{p}_1, \dots, \mathbf{p}_N)$ be N points on a hemisphere of \mathbb{S}^2 (resp., points in \mathbb{E}^2), and let $\mathbf{q} = (\mathbf{q}_1, \dots, \mathbf{q}_N)$ be a contraction of \mathbf{p} in \mathbb{S}^2 (resp., in \mathbb{E}^2). Then perimeter of the convex hull of \mathbf{q} is less than or equal to the perimeter of the convex hull of \mathbf{p} .*

Proof. This theorem in \mathbb{S}^2 follows from taking the spherical polar of the convex hull of the configurations \mathbf{p} and \mathbf{q} , which are the intersections of the hemispheres with centers at the vertices of the corresponding configurations. In this case the area of the polar and the perimeter of the convex hull sum up to 2π , and the result follows. Finally, the claim in \mathbb{E}^2 follows easily from the spherical version via a standard limiting procedure.

The Euclidean part of Corollary 2 has been proved independently by R. Alexander [1], V. Capovleas and J. Pach [3], and V. N. Sudakov[11].

Recall that the *extreme points* of a compact convex set X are the points that do not lie on the relative interior of a segment joining two other points of X . The following is a result of a compactness argument applied to a sequence of points that are dense in X .

Corollary 3. *If there is a contraction of the extreme points of a convex set in an open hemisphere of \mathbb{S}^n , then the polar of X has volume no larger than the volume of the polar of the convex hull of the image.*

2. Leapfrog Lemmas.

We repeat here one of our previous Lemmas from [2], the Leapfrog Lemma.

Lemma 1. *Suppose that $\mathbf{p} = (\mathbf{p}_1, \dots, \mathbf{p}_N)$ and $\mathbf{q} = (\mathbf{q}_1, \dots, \mathbf{q}_N)$ are two configurations in \mathbb{E}^n . Then the following is a continuous motion $\mathbf{p}(t) = (\mathbf{p}_1(t), \dots, \mathbf{p}_N(t))$ in \mathbb{E}^{2n} , that is analytic in t , such that $\mathbf{p}(0) = \mathbf{p}$, $\mathbf{p}(1) = \mathbf{q}$ and for $0 \leq t \leq 1$, $|\mathbf{p}_i(t) - \mathbf{p}_j(t)|$ is monotone:*

$$(1) \quad \mathbf{p}_i(t) = \left(\frac{\mathbf{p}_i + \mathbf{q}_i}{2} + (\cos \pi t) \frac{\mathbf{p}_i - \mathbf{q}_i}{2}, (\sin \pi t) \frac{\mathbf{p}_i - \mathbf{q}_i}{2} \right), \quad 1 \leq i < j \leq N.$$

We need to apply this to a sphere, rather than Euclidean space. Here we consider the unit spheres $\mathbb{S}^n \subset \mathbb{S}^{n+1} \subset \mathbb{S}^{n+2} \dots$ in such a way that each \mathbb{S}^n is the set of points that are a unit distance from the origin in \mathbb{E}^{n+1} . So we need the following.

Corollary 4. *Suppose that $\mathbf{p} = (\mathbf{p}_1, \dots, \mathbf{p}_N)$ and $\mathbf{q} = (\mathbf{q}_1, \dots, \mathbf{q}_N)$ are two configurations in \mathbb{S}^n . Then there is a monotone analytic motion from $\mathbf{p} = (\mathbf{p}_1, \dots, \mathbf{p}_N)$ to $\mathbf{q} = (\mathbf{q}_1, \dots, \mathbf{q}_N)$ in \mathbb{S}^{2n+1} .*

Proof. Apply Lemma 1 to each configuration $\mathbf{p} = (\mathbf{p}_1, \dots, \mathbf{p}_N)$ and $\mathbf{q} = (\mathbf{q}_1, \dots, \mathbf{q}_N)$ with the origin as an additional configuration point for each. So for each t , the configuration $\mathbf{p}(t) = (\mathbf{p}_1(t), \dots, \mathbf{p}_N(t))$ lies at a unit distance from the origin in \mathbb{E}^{2n+2} , which is just \mathbb{S}^{2n+1} .

3. The differentiable case.

We look at the case when there is a smooth motion of the configuration $\mathbf{p}(t)$ in \mathbb{S}^n . More precisely we consider the family $X(t) = X(\mathbf{p}(t))$ of convex spherical n -polytopes in \mathbb{S}^n having the same combinatorial face structure with facet hyperplanes being differentiable in the parameter t . The following classical theorem of Schläfli (see for example [8]) describes how the volume of $X(t)$ changes as a function of its dihedral angles and the volume of its $(n - 2)$ -dimensional faces.

Theorem 2 (Schläfli). *For each $(n - 2)$ -face $F_{ij}(t)$ of the convex spherical n -polytope $X(t)$ in \mathbb{S}^n let $\alpha_{ij}(t)$ represent the dihedral angle between the two facets $F_i(t)$ and $F_j(t)$ meeting at $F_{ij}(t)$. Then the following holds:*

$$(2) \quad \frac{d}{dt} \text{Vol}_n[X(t)] = \frac{1}{n-1} \sum_{F_{ij}} \frac{d\alpha_{ij}(t)}{dt} \text{Vol}_{n-2}[F_{ij}(t)],$$

to be summed over all $(n - 2)$ -faces.

Corollary 5. *If the configuration $\mathbf{p}(t)$ is a differentiable contraction in t , then with the notation of Theorem 2,*

$$\frac{d}{dt} \text{Vol}_n[X(t)] \geq 0.$$

Proof. As the distance between $\mathbf{p}_i(t)$ and $\mathbf{p}_j(t)$ is decreasing, the derivative of the dihedral angle $\frac{d\alpha_{ij}(t)}{dt} \geq 0$. The result then follows from (2).

4. Integral formulas.


The last piece of information that we need before we get to the main result is a way of relating higher-dimensional volumes to lower-dimensional volumes. Let X be any integrable set in \mathbb{S}^n . Recall that we regard

$$X \subset \mathbb{S}^n = \mathbb{S}^n \times \{\mathbf{0}\} \subset \mathbb{E}^{n+1} \times \mathbb{E}^{k+1}.$$

Regard

$$\{\mathbf{0}\} \times \mathbb{S}^k \subset \mathbb{E}^{n+1} \times \mathbb{E}^{k+1}.$$

Let $X * \mathbb{S}^k$ be the subset of \mathbb{S}^{n+k+1} consisting of the union of the geodesic arcs from each point of X to each point of $\{\mathbf{0}\} \times \mathbb{S}^k$. (So, in particular, $\mathbb{S}^n * \mathbb{S}^k = \mathbb{S}^{n+k+1}$). See the Figure.



FIGURE

Theorem 3. *For any integrable subset X of \mathbb{S}^n ,*

$$\text{Vol}_{n+k+1}[X * \mathbb{S}^k] = \frac{\kappa_{n+k+1}}{\kappa_n} \text{Vol}_n[X],$$

where $\kappa_n = \text{Vol}_n[\mathbb{S}^n]$, and $\kappa_{n+k+1} = \text{Vol}_{n+k+1}[\mathbb{S}^n * \mathbb{S}^k] = \text{Vol}_{n+k+1}[\mathbb{S}^{n+k+1}]$.

Proof. Since the $*$ operation (a kind of spherical join) is associative, we only need to consider the case when $k = 0$. Regard $\{\mathbf{0}\} \times \mathbb{S}^0 = \{N, S\}$, the north pole and the south pole of \mathbb{S}^{n+1} . We use polar coordinates centered at N to calculate the $(n+1)$ -dimensional volume of $X * \mathbb{S}^0$. Let $X(z) = X * \mathbb{S}^0 \cap [\mathbb{E}^{n+1} \times \{z\}]$, and let θ be the angle that a point in \mathbb{S}^{n+1} makes with N , the north pole in \mathbb{S}^{n+1} . So $z = z(\theta) = \cos \theta$. Then the volume element for $\mathbb{S}^n(z)$ is $dV_n(z) = (\sin^n \theta) dV_n(0)$ because $\mathbb{S}^n(z)$ is obtained from $\mathbb{S}^n(0)$ by a dilation by $\sin \theta$. Then

$$\begin{aligned} \text{Vol}_{n+1}[X * \mathbb{S}^0] &= \int_{X * \mathbb{S}^0} dV_n(z) d\theta \\ &= \int_0^\pi \int_{X(z(\theta))} dV_n(z) d\theta \\ &= \int_0^\pi (\sin^n \theta) \text{Vol}_n[X] d\theta \\ &= \text{Vol}_n[X] \int_0^\pi (\sin^n \theta) d\theta \\ &= \text{Vol}_n[X] \frac{\kappa_{n+1}}{\kappa_n}. \end{aligned}$$

The last integral can be seen by taking $X = \mathbb{S}^n$, or by performing the integral explicitly.

5. Proof of Theorem 1.

Let the configuration $\mathbf{q} = (\mathbf{q}_1, \dots, \mathbf{q}_N)$ be a contraction of the configuration $\mathbf{p} = (\mathbf{p}_1, \dots, \mathbf{p}_N)$ in \mathbb{S}^n . By Corollary 4, there is an analytic motion $\mathbf{p}(t)$, in \mathbb{S}^{2n+1} for $0 \leq t \leq 1$, where $\mathbf{p}(0) = \mathbf{p}$, and $\mathbf{p}(1) = \mathbf{q}$, and all the pairwise distances between the points of $\mathbf{p}(t)$ vary monotonically in t .

Without loss of generality we may assume that $X^n(\mathbf{p}(0))$ is a convex spherical n -polytope in S^n . Since $\mathbf{p}(t)$ is analytic in t , the intersection of the hemispheres in \mathbb{S}^{2n+1} , $X^{2n+1}(\mathbf{p}(t))$, is a convex spherical $(2n+1)$ -polytope with a constant combinatorial structure, except for a finite number of points in the interval $[0, 1]$. By Corollary 5, $\text{Vol}_{2n+1}[X^{2n+1}(\mathbf{p}(t))]$, is monotone increasing in t .

Recall that $X^n(\mathbf{p})$ and $X^n(\mathbf{q})$ are the intersections of the hemispheres centered at the points of \mathbf{p} and \mathbf{q} in \mathbb{S}^n . From the definition of the spherical join $*$,

$$\begin{aligned} X^n(\mathbf{p}) * \mathbb{S}^n &= X^{2n+1}(\mathbf{p}) = X^{2n+1}(\mathbf{p}(0)) \\ X^n(\mathbf{q}) * \mathbb{S}^n &= X^{2n+1}(\mathbf{q}) = X^{2n+1}(\mathbf{p}(1)). \end{aligned}$$

Hence by Theorem 3,

$$\text{Vol}_n[X^n(\mathbf{p})] = \frac{\kappa_n}{\kappa_{2n+1}} \text{Vol}[X^{2n+1}(\mathbf{p}(0))] \leq \frac{\kappa_n}{\kappa_{2n+1}} \text{Vol}[X^{2n+1}(\mathbf{p}(1))] = \text{Vol}_n[X^n(\mathbf{q})].$$

This finishes the proof of Theorem 1.

6. Comments.

We present some possible generalizations of the above theorems.

By replacing Theorem 2 of [2] with Theorem 5.1 of Csikós's paper [5] one can rewrite the proof of Theorem 1 of our paper [2] in a rather straightforward way such that it leads to the following relative of Theorem 1 of this paper. (We phrase the theorem for intersection of balls only, however, a reversed inequality for unions of balls can be proved in the same way.)

Theorem 4. *Let $\mathbf{p} = (\mathbf{p}_1, \dots, \mathbf{p}_N)$ and $\mathbf{q} = (\mathbf{q}_1, \dots, \mathbf{q}_N)$ be two configurations in \mathbb{S}^n such that \mathbf{q} is a piecewise-analytic contraction of \mathbf{p} in \mathbb{S}^{n+2} . If $X(\mathbf{p})$ (resp., $X(\mathbf{q})$) denotes the finite intersection of congruent balls of radius r , $0 < r < \pi$ in \mathbb{S}^n whose configuration of centers is \mathbf{p} (resp., \mathbf{q}), then*

$$\text{Vol}_n[X(\mathbf{p})] \leq \text{Vol}_n[X(\mathbf{q})].$$

Very recently, independently, Csikós [6] extended the above theorem for not necessarily congruent balls as well. These theorems and our proof [2] of the Kneser-Poulsen conjecture in Euclidean plane leads us to the following conjecture.

Leapfrog Conjecture on S^2 . *Suppose that $\mathbf{p} = (\mathbf{p}_1, \dots, \mathbf{p}_N)$ and $\mathbf{q} = (\mathbf{q}_1, \dots, \mathbf{q}_N)$ are two configurations in \mathbb{S}^2 . Then there is a monotone piecewise-analytic motion from $\mathbf{p} = (\mathbf{p}_1, \dots, \mathbf{p}_N)$ to $\mathbf{q} = (\mathbf{q}_1, \dots, \mathbf{q}_N)$ in \mathbb{S}^4 .*

Obviously, our Leapfrog Conjecture and the above mentioned theorems imply the Kneser-Poulsen conjecture for (congruent) circles in \mathbb{S}^2 .

Finally, we remark that Theorem 1 as well as Theorem 4 extend to flowers of balls as well. The notion of flowers of balls has been introduced in [7] by Gordon and Meyer (in set theoretic sense a flower of balls is a finite union of finite intersections of balls) and the proper technology to treat them analogously to intersections (resp., unions) of balls has been developed by Csikós in [5]. For more information on this see also our paper [2].

REFERENCES

1. R. Alexander, *Lipschitzian mappings and the total mean curvature of polyhedral surfaces, no., I.*, Trans. Amer. Math. Soc. **288** (1985), no. 2, 661–678, MR86c:52004.
2. K. Bezdek and R. Connelly, *Pushing disks apart - the Kneser-Poulsen conjecture in the plane*, J. reine angew. Math. **553** (2002), 221–236.
3. V. Capoyreas; J. Pach, *On the perimeter of a point set in the plane*, Discrete and computational geometry (New Brunswick, NJ) DIMACS Ser. Discrete Math. Theoret. Comput. Sci., Amer. Math. Soc., Providence, RI, **6** (1991), 67–76, MR92k:52013.
4. B. Csikós, *On the volume of the union of balls*, Discrete Comput. Geom. **20** (1998), no. 4, 449–461., MR99g:52008.
5. B. Csikós, *On the volume of flowers in space forms*, Geometriae Dedicata **1-3** (2001), no. 86, 57–59.
6. B. Csikós, *A Schläfli-type formula and its application to the Kneser-Poulsen problem*, Lecture at the 5th Geometry Festival, Budapest, Hungary (Nov. 28-30, 2002).
7. Y. Gordon; M. Meyer, *On the volume of unions and intersections of balls in Euclidean space*, Operator Theory: Advances and Applications **77** (1995), 91-101, MR96h:52008.
8. H. Kneser, *Der Simplexinhalt in der nichteuklidischen Geometrie*, Deutsche Math. **1** (1936), 337–340.
9. M. Kneser, *Einige Bemerkungen über das Minkowskische Flächenmass*, Arch. Math. **6** (1955), 382–390, MR17,469e.
10. E. T. Poulsen, *Problem 10*, Mathematica Scandinavica **2** (1954), 346.
11. V. N. Sudakov, *Gaussian random processes, and measures of solid angles in Hilbert space*, (Russian), Dokl. Akad. Nauk SSSR **197** (1971), 43–45, MR#6027.

EÖTVÖS UNIVERSITY DEPARTMENT OF GEOMETRY H-1117 BUDAPEST PÁZMÁNY PÉTER SÉTÁNY
1/C HUNGARY

E-mail address: `kbezdek@ludens.elte.hu`

DEPARTMENT OF MATHEMATICS MALOTT HALL CORNELL UNIVERSITY ITHACA, NY 14853 USA

E-mail address: `connelly@math.cornell.edu`