

# REVIEW

§5.7 (Substitution Methods)

MATH 1910 Recitation

September 13, 2016

- Try the **Substitution Method** when the integrand has the form  $f(u(x))u'(x)$ . If  $F$  is an antiderivative of  $f$ , then

$$\int f(u(x))u'(x) dx = \boxed{\phantom{000}}^{(1)} + C$$

- The differential of  $u(x)$  is related to  $dx$  by  $du = \boxed{\phantom{000}}^{(2)}$ .

- The **Change of Variables Formula** says that

- For indefinite integrals:  $\int f(u(x))u'(x) dx = \boxed{\phantom{000}}^{(3)}$

- For definite integrals:  $\int_a^b f(u(x))u'(x) dx = \boxed{\phantom{000}}^{(4)}$

# PRACTICE PROBLEMS

§5.7 (Substitution Methods)

MATH 1910 Recitation

September 13, 2016

(1) Evaluate the indefinite integral.

(a)  $\int x(x+1)^9 dx$

(b)  $\int \sin(2x-4) dx$

(c)  $\int \frac{x^3}{(x^4+1)^4} dx$

(d)  $\int \sqrt{4x-1} dx$

(e)  $\int x \cos(x^2) dx$

(f)  $\int \sin^5 x \cos x dx$

(g)  $\int \sec^2 x \tan^4 x dx$

(h)  $\int \frac{dx}{(2+\sqrt{x})^3}$

(2) Evaluate the definite integral.

(a)  $\int_0^1 \frac{x}{(x^2+1)^3} dx$

(b)  $\int_{10}^{17} (x-9)^{-2/3} dx$

(c)  $\int_{-8}^8 \frac{x^{15}}{3+\cos^2 x} dx$

(d)  $\int_0^{\pi/2} \sec^2(\cos \theta) \sin \theta d\theta$

(e)  $\int_{-4}^{-2} \frac{12x dx}{(x^2+2)^3}$

(f)  $\int_1^8 \sqrt{t+8} dt$

(g)  $\int_0^{\pi/3} \frac{\sin \theta}{\cos^{2/3} \theta} d\theta$

(h)  $\int_{-2}^4 |(x-1)(x-3)| dx$