

## REVIEW

- (1) For small small changes
- $\Delta x$
- ,
- $\Delta y$
- we have:

$$f(a + \Delta x, b) \approx f(a, b) + f_x(a, b)\Delta x$$

$$f(a, b + \Delta y) \approx f(a, b) + f_y(a, b)\Delta y$$

- (2) Clairaut's theorem states that mixed partials are equal as long as all functions we are dealing with are continuous. Hence, we can take higher partial derivatives in any order we please.

- (3) The linearization of
- $f$
- in two and three variables:

$$L(x, y) = f(a, b) + f_x(a, b)(x - a) + f_y(a, b)(y - b)$$

$$L(x, y, z) = f(a, b, c) + f_x(a, b, c)(x - a) + f_y(a, b, c)(y - b) + f_z(a, b, c)(z - c)$$

- (4) If
- $f_x$
- and
- $f_y$
- exist and are continuous in a disk containing
- $(a, b)$
- , then
- $f$
- is differentiable at
- $(a, b)$
- .

- (5)
- Equation for the tangent plane**
- to
- $z = f(x, y)$
- at
- $(a, b)$
- .

$$z = f(a, b) + f_x(a, b)(x - a) + f_y(a, b)(y - b)$$

- (6) The
- gradient**
- of a function
- $f$
- is given by
- $\nabla f = \left\langle \frac{\partial}{\partial x} f, \frac{\partial}{\partial y} f, \frac{\partial}{\partial z} f \right\rangle$
- .

- (7)
- Chain rule for paths:**
- $\frac{d}{dt} f(\mathbf{r}(t)) = \nabla f_{\mathbf{r}(t)} \cdot \mathbf{r}'(t)$

- (8) The directional derivative with respect to
- $\mathbf{u}$
- a unit vector is given by
- $D_{\mathbf{u}} f = \nabla f \cdot \mathbf{u}$

- (9) If the angle between
- $\mathbf{u}$
- and
- $\nabla f$
- is
- $\theta$
- , then
- $D_{\mathbf{u}} f = \|\nabla f\| \|\mathbf{u}\| \cos(\theta)$

- (10) The equation of the tangent plane to the level surface
- $F(x, y, z) = k$
- at point
- $P = (a, b, c)$
- is

$$\nabla F_P \cdot \langle x - a, y - b, z - c \rangle = 0$$

## PROBLEMS

1. The volume of a right-circular cone of radius  $r$  and height  $h$  is  $V = \frac{\pi}{3}r^2h$ . Use linear approximation to estimate the percentage change in volume of a right-circular cone of radius  $r = 40\text{cm}$  if the height is increased from 40 to 41cm.
2. Find  $f$  such that:
  - (a)  $\frac{\partial}{\partial x}f = 6x^2y$ ,  $\frac{\partial}{\partial y}f = 2x^3 - 3$
  - (b)  $\frac{\partial}{\partial x}f = e^x - y \sin(xy)$ ,  $\frac{\partial}{\partial y}f = -x \sin(xy) + 5y^4$
3. Find the points on the graph of  $z = 3x^2 - 4y^2$  at which the vector  $\mathbf{n} = \langle 3, 2, 2 \rangle$  is normal to the tangent plane.
4. A bug located at  $P = (3, 9, 4)$  begins walking in a straight line toward  $(5, 7, 3)$ . At what rate is the bug's temperature changing at  $P$  if the temperature is  $T(x, y, z) = xe^{y-z}$ ? Units are in meters and degrees in Celcius.
5. Determine the derivative of the function along the path given:
  - (a)  $f(x, y, z) = yx^2 - e^{xy} + \ln(x)$   $\mathbf{r}(t) = \langle t^2, \ln(t), \sqrt{t} \rangle$
  - (b)  $f(x, y, z) = \sin(xyz)$   $\mathbf{r}(t) = \langle e^t, \cos(4t), -t \rangle$
6. Find a function such that  $\nabla f = \langle 2xe^y - z, y^3 + x^2e^y, -x \rangle$