

REVIEW

- (1) **Green's Theorem:** \mathcal{D} is a domain in the plane and $\partial\mathcal{D}$ is its boundary. Then:

$$\oint_{\partial\mathcal{D}} F_1 dx + F_2 dy = \iint_{\mathcal{D}} \left(\frac{\partial F_2}{\partial x} - \frac{\partial F_1}{\partial y} \right) dA$$

or

$$\oint_{\partial\mathcal{D}} \mathbf{F} \cdot d\mathbf{r} = \iint_{\mathcal{D}} \text{curl}_z(\mathbf{F}) dA.$$

- (2) **Stokes' Theorem:** \mathcal{S} is an oriented surface and $\partial\mathcal{S}$ is its boundary. Then:

$$\oint_{\partial\mathcal{S}} \mathbf{F} \cdot d\mathbf{r} = \iint_{\mathcal{S}} \text{curl}(\mathbf{F}) \cdot d\mathbf{S}.$$

- (3) **Divergence Theorem:** Let \mathcal{S} be a closed surface that encloses a region \mathcal{W} in \mathbb{R}^3 . Assume that \mathcal{S} is piecewise smooth and is oriented by normal vectors pointing to the outside of \mathcal{W} . Let \mathbf{F} be a vector field whose domain contains \mathcal{W} . Then

$$\iint_{\mathcal{S}} \mathbf{F} \cdot d\mathbf{S} = \iiint_{\mathcal{W}} \text{div}(\mathbf{F}) dV.$$

PROBLEMS

- (1) Use Green's Theorem to evaluate the line integral around the given closed curve.
- (a) $\oint_C xy^3 dx + x^3y dy$, where C is the rectangle $-1 \leq x \leq 2$, $-2 \leq y \leq 3$, oriented counterclockwise.
 - (b) $\oint_C y^2 dx - x^2 dy$, where C consists of the arcs $y = x^2$ and $y = \sqrt{x}$, $0 \leq x \leq 1$, oriented clockwise.
- (2) Let I be the flux of $\mathbf{F} = \langle e^y, 2xe^{x^2}, z^2 \rangle$ through the upper hemisphere \mathcal{S} of the unit sphere.
- (a) Let $\mathbf{G} = \langle e^y, 2xe^{x^2}, 0 \rangle$. Find a vector field \mathbf{A} such that $\text{curl}(\mathbf{A}) = \mathbf{G}$. circulation of \mathbf{A} around $\partial\mathcal{S}$.
 - (b) Use Stokes' theorem to show that the flux of \mathbf{G} through \mathcal{S} is zero. *Hint:* Calculate the (c) Calculate I . *Hint:* Use (b) to show that I is equal to the flux of $\langle 0, 0, z^2 \rangle$ through \mathcal{S} .
- (3) Let \mathcal{S} be the portion of the plane $z = x$ contained in the half-cylinder of radius R . Use Stokes' theorem to calculate the circulation of $\mathbf{F} = \langle z, x, y + 2z \rangle$ around the boundary of \mathcal{S} (a half-ellipse) in the counterclockwise direction when viewed from above.
- (4) Show that the circulation of $\mathbf{F}(x, y, z) = \langle x^2, y^2, z(x^2 + y^2) \rangle$ around any curve C on the surface of the cone $z^2 = x^2 + y^2$ is equal to zero.
- (5) Compute the flux of $\mathbf{F} = \langle xyz + xy, \frac{1}{2}y^2(1 - z) + e^x, e^{x^2+y^2} \rangle$, \mathcal{S} is the boundary of the solid bounded by the cylinder $x^2 + y^2 = 16$ and the planes $z = 0$ and $z = y - 4$.
- (6) Compute the flux of $\mathbf{F} = \langle \sin(yz), \sqrt{x^2 + z^4}, x \cos(x - y) \rangle$, \mathcal{S} is any smooth closed surface that is the boundary of a region in \mathbb{R}^3 .




Figure 1: Problem 5.




Figure 2: Problem 4.