

Enumeration of Symmetry Classes of Convex Polyominoes in the Square Lattice*

P. LEROUX, E. RASSART, A. ROBITAILLE

March 30, 1998

*LaCIM, Department of Mathematics
Université du Québec à Montréal (UQAM)*

Abstract

This paper concerns the enumeration of rotation-type and congruence-type convex polyominoes on the square lattice. These can be defined as orbits of the groups \mathfrak{C}_4 , of rotations, and \mathfrak{D}_4 , of symmetries, of the square, acting on (translation-type) polyominoes. In virtue of Burnside's Lemma, it is sufficient to enumerate the various symmetry classes (fixed points) of polyominoes defined by the elements of \mathfrak{C}_4 and \mathfrak{D}_4 . Using the Temperley–Bousquet–Mélou methodology, we solve this problem and provide explicit or recursive formulas for their generating functions according to width, height and area. We also enumerate the class of asymmetric convex polyominoes, using Möbius inversion, and prove that their number is asymptotically equivalent to the number of convex polyominoes, a fact which is empirically evident.

Résumé

Cet article porte sur l'énumération de polyominos à rotations et réflexions près dans un réseau carré. Ces polyominos peuvent être considérés comme des orbites dans l'action du groupe diédral \mathfrak{D}_4 du carré et de son sous-groupe \mathfrak{C}_4 , le groupe des rotations du carré, sur l'ensemble de polyominos convexes (à translations près). Le lemme de Burnside réduit alors le problème à l'énumération des ensembles de points fixes (classes de symétrie) de l'action. Utilisant la méthodologie de Temperley–Bousquet–Mélou, nous donnons des formules explicites ou récurrentes pour les séries génératrices de ces types de polyominos selon la largeur, la hauteur et l'aire. Nous énumérons aussi, à l'aide de l'inversion de Möbius, la classe des polyominos convexes asymétriques et prouvons que ce nombre est asymptotique à celui des polyominos convexes, un fait mis en évidence expérimentalement.

1 Introduction

The enumeration of polyominoes and of self-avoiding walks of a square lattice is a deep and difficult combinatorial problem. These structures are important in statistical physics, as discrete models for polymers, clusters, percolation and phase transitions. See, for instance, Privman and Švrakić [31], Viennot [35], Bousquet–Mélou [5], Guttmann, Prellberg and Owczarek [18], Langlands, Pouliot and Saint-Aubin [25], Prellberg and Brak [29] and Hughes [19] and [20].

Recent research in this area has mainly dealt with polyominoes up to translation (where two polyominoes are equivalent if some translation of the plane maps one onto the other). The

*Work partially supported by NSERC (Canada) and FCAR (Québec).

equivalence classes are then called *translation-type* polyominoes, or simply polyominoes when the context is clear. The exact enumeration of polyominoes is an open problem, but considerable progress has been made for classes of polyominoes defined by special properties of convexity or directedness (see [6]). A polyomino is said to be *convex* when its intersection with any vertical or horizontal line is connected. A polyomino P is (North-East) *directed* if each of its cells can be reached from some distinguished cells (the source cells) by a path that is contained in P and has only North and East steps (see Figure 1). Formulas have been published for the generating function of convex polyominoes according to perimeter ([11], [22] and [16]), to height and width ([27]) and joint parameters area, height and width ([26], [7] and [4]).




Figure 1: Polyominoes: (a) convex; (b) directed (one source) convex; (c) directed (diagonal source).

In this paper, we consider *rotation-type* polyominoes, that is equivalence classes of polyominoes under rotations, and *congruence-type* polyominoes, where rotations and reflections are permitted. They occur naturally as pieces that can freely move in space, as used in plane packing problems (see [23] and [14]). For example, there are 19 (translation-type) tetrominoes, 7 rotation-type tetrominoes and 5 congruence-type tetrominoes (tetrominoes are polyominoes with area 4).




Figure 2: The 19 tetrominoes and their equivalence classes.

As Figure 2 illustrates, congruence-type polyominoes can be seen as orbits of the action of the group \mathfrak{D}_4 of symmetries of the square on the set of translation-type polyominoes. Similarly,

rotation-type polyominoes are orbits of the subgroup \mathfrak{C}_4 of \mathfrak{D}_4 containing the rotations. In virtue of Burnside's Lemma, the enumeration of rotation-type or congruence-type polyominoes reduces to the enumeration of the various symmetry classes, i.e. the sets of fixed polyominoes determined by the group elements of \mathfrak{C}_4 and \mathfrak{D}_4 . Figure 3 illustrates the class of vertically symmetric tetrominoes, i.e. tetrominoes left fixed by a reflection about a vertical axis.




Figure 3: Vertically symmetric tetrominoes.

The problem of enumerating rotation-type and congruence-type polyominoes is mentioned in Golomb [14] and Klarner [23], but only experimental results, for polyominoes of area up to 24, are reported, together with asymptotic estimates. Here we solve this problem for the class of convex polyominoes. The study of convex polyominoes was suggested by D. Knuth (see [24]). Convex polyominoes also appear naturally in statistical mechanics as self-avoiding polygons whose number of steps is equal to the perimeter of their minimal bounding rectangle (see [16]). Using elementary methods and the Temperley–Bousquet–Mélou methodology [4], we give explicit or recursive formulas for the generating functions according to perimeter and area of each of their symmetry classes. These results are then combined to enumerate rotation-type and congruence-type convex polyominoes. Using Möbius inversion in the lattice of subgroups of \mathfrak{D}_4 , we also count asymmetric convex polyominoes. Since the symmetry class defined by each subgroup of \mathfrak{D}_4 (except the trivial one) is significantly smaller than the whole class, we conclude that the number of asymmetric convex polyominoes, as the area goes to infinity, is asymptotic to the number of convex polyominoes. In other words, almost all convex polyominoes are asymmetric.

The paper is organized as follows. In section 2, we review some basic results that are used in the sequel on group actions and convex polyomino enumeration, according to width, height and area. Polyomino classes that are covered include partitions, stacks, in particular the new class of shifted stacks, directed convex and convex polyominoes. We then treat successively the symmetry classes of convex polyominoes in sections 3 (rotational symmetries), 4 (horizontal and vertical symmetries) and 5 (diagonal symmetries). The enumeration of diagonally symmetric convex polyominoes requires the determination of a subsidiary class of diagonal source (shifted) directed convex polyominoes, in subsection 5.2. These results are then combined to enumerate rotation-type polyominoes (subsection 3.3) and congruence-type polyominoes (section 6) according to half-perimeter and area. Finally section 7 is devoted to asymmetric convex polyominoes. Some numerical results are found in Appendices A and B.

Acknowledgements

The first author would like to thank Mireille Bousquet-Mélou, Richard Brak, Dominique Gouyou-Beauchamps and Tony Guttmann for stimulating and useful discussions.

2 Preliminaries

2.1 Review of convex polyomino enumeration

We recall here some basic and known results on the enumeration of some classes of translation-type polyominoes that will be used in this paper, namely Ferrers diagrams or partition polyominoes (P), partitions with empty parts allowed (P_0), (shifted) partitions with distinct parts (P_S), varieties of stack or pyramid polyominoes (T , T_0 and T_S), directed (one source) convex (D), and convex (C) polyominoes. For each of these classes, we give the generating series $F(x, y, q)$ where the variable x marks the width of the polyominoes, y marks the height and q the area. For example, the weight of the convex polyomino of Figure 1(a) according to these parameters is $x^7 y^6 q^{20}$. We will often use the same letters P , T , D , \dots to denote a class of polyominoes, its generating function, or a generic polyomino in the class.

2.1.1 Partition polyominoes

A partition polyomino is the Ferrers diagram of a non-empty partition (see Figure 4(a)). It can be viewed as a left justified pile of weakly decreasing rows (from top to bottom). Let $P(x, y, q)$ denote their generating series. In this context, the variable x marks the size of the maximum part and y the number of parts. Let also $P_0(x, y, q)$ denote the generating series of partitions where empty rows (contributing to the height) are permitted (see Figure 4(b)), and $P_S(x, y, q)$, of partitions with distinct parts (diagonally shifted) (see Figure 4(c)).




Figure 4: Partition polyominoes. (a) P : general; (b) P_0 : with empty parts allowed; (c) P_S : shifted.

We then have the following classical results (see [1]), for which direct counting arguments can be given:

$$\begin{aligned}
 P(x, y, q) &= \sum_{m \geq 1} \frac{x^m y q^m}{(yq)_m} \\
 &= \sum_{m, k \geq 1} x^m y^k q^{m+k-1} \begin{bmatrix} m+k-2 \\ k-1 \end{bmatrix}_q, \tag{1}
 \end{aligned}$$

$$\begin{aligned}
 P_0(x, y, q) &= \sum_{m \geq 0} \frac{x^m y q^m}{(y)_{m+1}} \\
 &= \sum_{m \geq 0, k \geq 1} x^m y^k q^m \begin{bmatrix} m+k-1 \\ k-1 \end{bmatrix}_q, \tag{2}
 \end{aligned}$$

$$P_S(x, y, q) = \sum_{m \geq 1} x^m y q^m (-yq)_{m-1}$$

$$= \sum_{m \geq 1} \sum_{k=1}^m x^m y^k q^{m + \binom{k}{2}} \begin{bmatrix} m-1 \\ k-1 \end{bmatrix}_q, \quad (3)$$

where we have used the familiar notation

$$(a)_n = (a; q)_n = (1-a)(1-aq) \dots (1-aq^{n-1}), \text{ for } n \geq 0 \quad (4)$$

so that in particular $(-yq)_m = (1+yq)(1+yq^2) \dots (1+yq^m)$, and where, for $0 \leq k \leq n$,

$$\begin{bmatrix} n \\ k \end{bmatrix}_q = \frac{(q)_n}{(q)_k (q)_{n-k}} \quad (5)$$

is the q -binomial (Gaussian) polynomial.

2.1.2 Stack polyominoes

A *stack* or *pyramid* polyomino is defined as a left justified unimodal (i.e. weakly increasing, then weakly decreasing) pile of rows (see Figure 5(a)). Let $T(x, y, q)$ (T for French “tas”) denote the generating series of stacks, and $T_0(x, y, q)$, the generating series of stacks with empty rows allowed at either end (see Figure 5(b)).




Figure 5: Stack polyominoes. (a) T : general; (b) T_0 : with empty parts allowed; (c) T_S : shifted stacks.

By extracting coefficients, define, for $n \geq 1$,

$$T_n(x, q) = [y^n] T(x, y, q) \quad \text{and} \quad T_{0,n}(x, q) = [y^n] T_0(x, y, q), \quad (6)$$

as the width and area generating series for stack polyominoes having height equal to n . It is clear that $T(x, y, q) = xT_0(x, yq, q)$ and that $T_n(x, q) = xq^n T_{0,n}(x, q)$. Hence any enumerative result for the class T_0 will yield a result for T . By concentrating on the width of the largest row, we obtain

$$T_0(x, y, q) = \sum_{m \geq 0} \frac{x^m y q^m}{(y)_{m+1} (y)_m} \quad (7)$$

and

$$T_{0,n}(x, q) = \sum_{m \geq 1} \sum_{j=1}^n x^m q^m \begin{bmatrix} m+j-1 \\ m \end{bmatrix}_q \begin{bmatrix} m+n-j-1 \\ m-1 \end{bmatrix}_q. \quad (8)$$

The following simple expression is due to Bousquet-Mélou and Viennot [9].

Proposition 1 *The width and area generating series for stacks having n (possibly empty) rows is given by*

$$T_{0,n}(x, q) = \frac{V_n}{(xq)_n}, \quad n \geq 1, \quad (9)$$

where $V_n = V_n(x, q)$ are polynomials in x and q defined by the recurrence

$$V_n = 2V_{n-1} + (xq^{n-1} - 1)V_{n-2}, \quad n \geq 2, \quad (10)$$

with $V_0 = V_1 = 1$. Moreover, V_n can be given explicitly as

$$V_n = 1 + \sum_{1 \leq k \leq n/2} \left(x^k q^{k^2} \sum_{m=k}^{n-k} \begin{bmatrix} m \\ k \end{bmatrix}_q \begin{bmatrix} n-m-1 \\ k-1 \end{bmatrix}_q \right). \quad (11)$$

Proof. Setting $T_{0,0}(x, q) = 1$ by convention (the empty stack), it is easy to see that for $n \geq 2$,

$$T_{0,n} = xq^n T_{0,n} + 2T_{0,n-1} - T_{0,n-2}, \quad (12)$$

so that $(1 - xq^n)T_{0,n} = 2T_{0,n-1} - T_{0,n-2}$. Since $T_{0,1} = 1/(1 - xq) = 1/(xq)_1$, we see that indeed $T_{0,n} = V_n/(xq)_n$ where V_n satisfies $V_0 = V_1 = 1$ and recurrence (10). This recurrence can also be used to prove that for $n \geq 1$, V_n is the generating polynomial for T_0 -stacks of height $n - 1$ having the additional property that any column exceeds strictly its right neighbor both at the top and at the bottom. Formula (11) then follows easily. \square

2.1.3 Shifted stacks

In analogy with shifted partitions P_S , we introduce the class T_S of (diagonally) shifted stacks (Figure 5(c)), and we denote by $T_S(u, x, y, q)$ their generating series, where u is an extra variable which marks the top row width. For example, the shifted stack of Figure 5(c) has global weight $u^2 x^8 y^7 q^{20}$. This generating series will be used in Section 5.

Define the *South-East corner* of a shifted stack T_S to be the bottommost cell of the rightmost column of T_S . The *base* of T_S consists of all rows below (and including) the South-East corner row. It is a shifted partition P_S with the additional property that the first row exceeds the other rows, i.e. that the first part of the partition exceeds the others by at least 2. Let $P_1(u, x, y, q)$ denote the generating series of all such possible bases of shifted stacks. We obviously have

$$P_1(u, x, y, q) = uxyq + y \sum_{k \geq 2} (uxq)^k (-yq)_{k-2}. \quad (13)$$

Proposition 2 *The generating function $T_S(u) = T_S(u, x, y, q)$ of shifted stacks satisfies the functional equation*

$$T_S(u) = P_1(u, x, y, q) + \frac{xyu^2 q^2}{1 - uq} (T_S(1) - T_S(uq)) \quad (14)$$

Proof. Equation (14) states that a shifted stack is either a P_1 -polyomino or is obtained from a P_1 -polyomino by iterating the procedure that adds a row on top of a shifted stack. At each step, the added row should be shifted one cell to the left and should not exceed the previous row on the right hand side. Now the term $xyu^2 q^2 (1 - uq)^{-1} T_S(1)$ enumerates the cases where a shifted row of any length (≥ 2) is added on top of a shifted stack, and $xyu^2 q^2 (1 - uq)^{-1} T_S(uq)$ counts the cases where the added row does exceed on the right, so that this term should in fact be subtracted. \square




Figure 6: A variation of construction B of Bousquet-Mélou.

The above procedure of adding a left shifted non-exceeding row (see Figure 6) can be seen as a variation of construction B of Bousquet-Mélou [4], where Temperley's methodology is translated into linear q -equations. (see also [12]). Using Lemma 2.3 of [4] which solves (14) by iteration, we find the following closed form:

Corollary 3 *The generating series $T_S(u) = T_S(u, x, y, q)$ for shifted stacks is given by*

$$T_S(u) = \frac{E(u) + E(1)F(u) - E(u)F(1)}{1 - F(1)}, \quad (15)$$

where

$$E(u) = \sum_{n \geq 0} \frac{(-1)^n x^{n+1} y^n u^{2n} q^{n^2+n} P_1(uq^n, x, y, q)}{(uq)_n}, \quad (16)$$

$$F(u) = \sum_{n \geq 0} \frac{(-1)^n x^{n+1} y^{n+1} u^{2n+2} q^{n^2+3n+2}}{(uq)_{n+1}}. \quad (17)$$

□

Note that the functional equation (14) and its solution (15) remain valid for other possible classes P_1 of polyominoes as bases. The method can then be adapted to enumerate Dyck words (P_1 is a single cell) and left factors of Dyck words (P_1 is a single row) according to length, area and size of the first stretch of x 's (marked by u).

2.1.4 Directed convex and convex polyominoes

We now give without proof the generating function $D(s, x, y, q)$ for directed convex polyominoes, where the variable s marks the leftmost column height, borrowed from Bousquet-Mélou [4] (Theorem 3.4, with $t = 1$).

Proposition 4 *The generating function for directed convex polyominoes is given by*

$$D(s, x, y, q) = y \frac{M_1(s)J_0(1) - M_1(1)J_0(s) + M_1(1)}{J_0(1)}, \quad (18)$$

where

$$J_0(s) = \sum_{n \geq 0} \frac{(-1)^n x^n s^n q^{\binom{n+1}{2}}}{(sq)_n (syq)_n} \quad (19)$$

and

$$M_1(s) = \sum_{n \geq 1} \left(\frac{x^n q^n}{(syq)_n} \left(\frac{s}{(syq)_{n-1}} + \sum_{m=1}^{n-1} \frac{(-1)^m s^m q^{\binom{m}{2}}}{(sq)_{n-1} (syq^{m+1})_{n-m-1}} \right) \right). \quad (20)$$

□

We also extract from [4] (Theorem 4.4, with $t = 1$) the generating function $C(x, y, q)$ for convex polyominoes. We introduce the series $E(s) = E(s, x, y, q)$ by

$$E(s) = 2y^2 \frac{M_1(1)}{J_0(1)} \alpha(s) - ya(s), \quad (21)$$

where J_0 and M_1 are given by (19) and (20),

$$\alpha(s) = \sum_{m \geq 1} \left(\frac{(-1)^m x^m s^m q^{\binom{m+1}{2}}}{(sq)_m} \sum_{n=1}^m \frac{(-1)^n s^n q^{\binom{n+1}{2}}}{(sq)_{n-1} (syq^n)_{m-n+1}} \right) \quad (22)$$

and

$$\begin{aligned} a(s) = & \sum_{m \geq 1} \left(\frac{x^m q^m}{1 - syq^m} \left(-\frac{s^{2m-1} q^{m(m-1)}}{(sq)_{m-1}^2} + y \sum_{n=1}^{m-1} \frac{s^{2n} q^{n^2} (syq^n - 2)}{(sq)_{n-1}^2 (syq^n)_{m-n}^2} \right. \right. \\ & \left. \left. + 2y \sum_{1 \leq n \leq k < m} \frac{(-1)^{n+k} s^{n+k} q^{\binom{n+1}{2}} q^{\binom{k}{2}}}{(sq)_{n-1} (syq^n)_{m-n} (sq)_k (syq^{k+1})_{m-k-1}} \right) \right). \end{aligned} \quad (23)$$

Proposition 5 *The generating series $C(x, y, q)$ for convex polyominoes is given by*

$$C(x, y, q) = J_1 E'(1) - K_1 E(1), \quad (24)$$

where $E(s)$ is given by (21), the derivative E' is taken with respect to s ,

$$J_1 = \sum_{n \geq 1} \frac{x^n q^n}{(q)_{n-1} (q)_n}, \quad (25)$$

and

$$K_1 = -1 + \sum_{n \geq 1} \left(\frac{x^n q^n}{(q)_{n-1} (q)_n} \left(\sum_{k=1}^{n-1} \frac{2q^k}{1 - q^k} + \frac{q^n}{1 - q^n} \right) \right). \quad (26)$$

□

2.2 The action of the dihedral group \mathfrak{D}_4 on polyominoes

The dihedral group \mathfrak{D}_4 is the symmetry group of the square. It has eight elements, usually represented as $1, r, r^2, r^3, h, v, d_1$ and d_2 , where 1 denotes the identity element, r denotes a rotation by a right angle, h and v reflections with respect to the horizontal and vertical axes respectively, and d_1 and d_2 reflections about the two diagonal axes of the square. The (nontrivial) subgroups of \mathfrak{D}_4 are the following:

- The cyclic subgroups of order 2: $\langle r^2 \rangle, \langle h \rangle, \langle v \rangle, \langle d_1 \rangle, \langle d_2 \rangle$
- The cyclic subgroup of order 4: $\mathfrak{C}_4 = \langle r \rangle = \{1, r, r^2, r^3\}$
- The (non-cyclic) subgroups of order 4: $\langle h, v \rangle = \{1, h, v, r^2\}$
 $\langle d_1, d_2 \rangle = \{1, d_1, d_2, r^2\}$

The group \mathfrak{D}_4 acts on (translation-type) polyominoes in a natural way, by rotating or reflecting them. Let \mathcal{P} be a finite set of polyominoes closed under the action of \mathfrak{D}_4 . We could take, for example, the set of convex polyominoes with a given area and/or perimeter. Then any subgroup G of \mathfrak{D}_4 acts on \mathcal{P} and the G -orbits are the equivalence classes of polyominoes modulo the transformations in G . The set of G -orbits of polyominoes in \mathcal{P} is denoted by \mathcal{P}/G . In particular, the set of rotation-type polyominoes is precisely $\mathcal{P}/\mathfrak{C}_4$, and the set of congruence-type polyominoes, $\mathcal{P}/\mathfrak{D}_4$.

The number of distinct G -orbits of polyominoes in \mathcal{P} is given by the Cauchy-Frobenius formula (alias Burnside's Lemma):

$$|\mathcal{P}/G| = \frac{1}{|G|} \sum_{g \in G} |\text{Fix}_{\mathcal{P}}(g)|, \quad (27)$$

where $\text{Fix}_{\mathcal{P}}(g)$ is the subset of \mathcal{P} consisting of the invariant polyominoes under the transformation g . The $\text{Fix}_{\mathcal{P}}(g)$ is called the *symmetry class* of g , and we call its elements *g -symmetric polyominoes*. Formula (27) extends to infinite classes of polyominoes by taking a weighted cardinality, e.g. the generating series with respect to half-perimeter (variable t) and area (variable q).

In the following sections we compute the generating series of the symmetry classes $\text{Fix}_{\mathcal{P}}(g)$ for the set \mathcal{C} of all convex (translation-type) polyominoes, and for $g \in \mathfrak{D}_4$. We use the notation

$$F_g(t, q) = \text{generating series of } \text{Fix}_{\mathcal{C}}(g).$$

Setting

$$(C/G)(t, q) = \text{generating series of } G\text{-orbits of convex polyominoes}$$

we have

$$(C/G)(t, q) = \frac{1}{|G|} \sum_{g \in G} F_g(t, q). \quad (28)$$

The concept of symmetry classes of polyominoes is extended to any subgroup G of \mathfrak{D}_4 by setting

$$\begin{aligned} S_{\geq G} &= \bigcap_{g \in G} \text{Fix}_{\mathcal{P}}(g) = \{P \in \mathcal{P} \mid \text{Stab}(P) \supseteq G\}, \\ S_{=G} &= \{P \in \mathcal{P} \mid \text{Stab}(P) = G\}, \end{aligned}$$

where $\text{Stab}(P)$ denotes the stabilizer of P , that is the subgroup of \mathfrak{D}_4 containing the transformations which leave P unchanged. Observe in particular that $S_{=0}$ is the set of asymmetric polyominoes. We let $F_{\geq G}$ and $F_{=G}$ denote the generating series of the sets $S_{\geq G}$ and $S_{=G}$ respectively. The series $F_{\geq G}$ for the subgroups G of \mathfrak{D}_4 are easier to compute directly than $F_{=G}$. We obviously have

$$F_{\geq G} = \sum_{H \supseteq G} F_{=H}, \quad (29)$$

where H ranges over the lattice of subgroups of \mathfrak{D}_4 , denoted by $\mathcal{S}(\mathfrak{D}_4)$.

Using Möbius inversion (see Rota [32]) in the lattice $\mathcal{S}(\mathfrak{D}_4)$, we then find

$$F_{=G} = \sum_{H \supseteq G} \mu(G, H) F_{\geq H}, \quad (30)$$

where $\mu(G, H)$ is the value of the Möbius function in $\mathcal{S}(\mathfrak{D}_4)$. In particular, taking $G = 0 = \{1\}$, the minimum element in $\mathcal{S}(\mathfrak{D}_4)$, we obtain the generating series for the asymmetric convex polyominoes:

$$F_{=0} = \sum_{H \in \mathcal{S}(\mathfrak{D}_4)} \mu(0, H) F_{\geq H}. \quad (31)$$




Figure 7: Lattice of the subgroups of \mathfrak{D}_4 and the Möbius function $\mu(0, H)$ on the lattice.

The value of the Möbius function $\mu(0, H)$ is easy to compute recursively (see Rota [32]), using the formulas

$$\mu(0, y) = - \sum_{x < y} \mu(0, x), \quad \mu(0, 0) = 1. \quad (32)$$

These values of μ on $\mathcal{S}(\mathfrak{D}_4)$ are given in Figure 7.

Note that for a cyclic subgroup $\langle g \rangle$ of \mathfrak{D}_4 , $F_{\geq \langle g \rangle}$ is precisely $F_g(t, q)$. The series $F_{hv}(t, q) = F_{\geq \langle h, v \rangle}$ and $F_{d_1 d_2}(t, q) = F_{\geq \langle d_1, d_2 \rangle}$ are also computed in the following sections.

3 Rotational symmetry classes

We now study the convex polyominoes admitting a rotational symmetry. Observe that if a polyomino is r -symmetric, where r denotes a 90° rotation, then it is also r^3 -symmetric and conversely, since $r^3 = r^{-1}$. In other words, $\text{Fix}_{\mathcal{P}}(r) = \text{Fix}_{\mathcal{P}}(r^3)$ and $F_r(t, q) = F_{r^3}(t, q)$. Hence it will be sufficient to determine the two generating series $F_r(t, q)$ and $F_{r^2}(t, q)$ of r -symmetric and of r^2 -symmetric convex polyominoes, according to half-perimeter and area.

3.1 r -Symmetric convex polyominoes

Observe that the minimal bounding rectangle of an r -symmetric convex polyomino is a square (see Figure 8(b)). In particular, the half-perimeter is always even. There is also a maximal square fitting inside the polyomino (see Figure 8(a)).

A first way to construct an r -symmetric convex polyomino P is to start with its maximal fitting square S and to glue the same stack polyomino T , after appropriate rotations, on each of its sides. The added stack can have empty parts on each side. The generating series $T_0(x, y, q)$ of such stacks is given in Subsection 2.1.2 (see formulas (7), (9) and (8)).




Figure 8: Two ways of constructing a convex r -symmetric polyomino.

Proposition 6 *The generating function for convex r -symmetric polyominoes $F_r(t, q)$ is given by*

$$F_r(t, q) = T_0(t^4, (y^m \mapsto t^{2m} q^{m^2})_{m \geq 1}, q^4), \quad (33)$$

Proof. Suppose that the stack T has height m , width n and area a . Then the polyomino P will have half-perimeter $2m + 4n$ and area $m^2 + 4a$. The “substitution” (33) gives exactly this. \square

A second way to construct an r -symmetric convex polyomino P , of half-perimeter $2m$, say, is to start with the minimal bounding square, of side m , and to remove from each corner a given (appropriately rotated) Ferrers diagram F with half-perimeter at most $m - 1$ (see Figure 8(b)). Denote by $D_m(q)$ the area generating series of Ferrers diagrams with half-perimeter less than or equal to m . If the perimeter of a diagram is less than m , then we can force it to be m by adding empty columns to the diagram (see Figure 9). We will call such diagrams, *extended* Ferrers diagrams. Thus $D_m(q)$ is the area generating function of the extended Ferrers diagrams with half-perimeter m . By convention $D_0(q) = 1$, and $D_1(q) = 1$ (one empty column).




Figure 9: Extended Ferrers diagram.

Proposition 7 *For $m \geq 1$,*

$$D_m(q) = \sum_{i=0}^{m-1} q^i \begin{bmatrix} m-1 \\ i \end{bmatrix}_q. \quad (34)$$

Proof. Let i be the height of the first column on the right. Then this first column contributes for q^i to the area. Removing that column yields a binomial path from A to B , of length $m - 1$ with i vertical steps (see Figure 9). The area generating function for such paths is given by the q -binomial coefficient $\begin{bmatrix} m-1 \\ i \end{bmatrix}_q$. Since i can vary from 0 to $m - 1$, the conclusion follows. \square

Proposition 10 *The area generating polynomial for r -symmetric convex polyominoes with half-perimeter $2m$ is given by*

$$a_{2m}(q) = q^{m^2} \left[\sum_{i=0}^{m-2} q^i \begin{bmatrix} m-2 \\ i \end{bmatrix}_q \right]_{q \mapsto q^{-4}}, \quad (39)$$

for $m \geq 2$. □

Corollary 11 *Let $a_{2m} = a_{2m}(1)$, the number of convex r -symmetric polyominoes with half-perimeter $2m$. Then $a_2 = 1$ and*

$$a_{2m} = 2^{m-2} \quad (40)$$

for $m \geq 2$. □

The following recurrences are useful in order to compute explicitly the polynomials $a_{2m}(q)$.

Proposition 12 *The polynomials $a_{2m}(q)$ satisfy the recurrences*

$$(a) \quad a_{2m}(q) = q^{2m-1} a_{2m-2}(q) + \sum_{j=1}^{m-2} q^{(m-2)^2+4-j^2} a_{2j}(q), \quad m \geq 1, \quad (41)$$

$$(b) \quad a_{2m}(q) = q^{2m-5} (1 + q^4) a_{2m-2}(q) + q^4 (1 - q^{4m-12}) a_{2m-4}(q), \quad m \geq 3, \quad (42)$$

with initial conditions $a_0(q) = 1$, $a_2(q) = q$ and $a_4(q) = q^4$.

Proof. This follows immediately from Proposition 8. □

Here are the first few terms of the generating series $F_r(t, q)$ for convex r -symmetric polyominoes:

$$F_r(t, q) = t^2 q + t^4 q^4 + t^6 q^5 + t^6 q^9 + 2t^8 q^8 + t^{10} q^9 + t^8 q^{12} + \dots \quad (43)$$

3.2 r^2 -Symmetric convex polyominoes

r^2 -symmetric convex polyominoes fall into two categories: those with even width and those with odd width. Thus, if $F_{r^2}(x, y, q)$ denotes the generating series for r^2 -symmetric convex polyominoes according to width, height and area, we have

$$F_{r^2}(x, y, q) = F_{r^2,e}(x, y, q) + F_{r^2,o}(x, y, q), \quad (44)$$

where $F_{r^2,e}$ (resp. $F_{r^2,o}$) is the generating series for r^2 -symmetric convex polyominoes of even width (resp. odd width). We calculate these series in the next two propositions. Recall that $T(x, y, q)$ is the generating function of stack polyominoes and $D(s, x, y, q)$, that of directed (one source) convex polyominoes, where s is an extra variable which marks the length of the leftmost column (see Subsection 2.1.4).

Proposition 13 *Let $F_{r^2,e}(x, y, q)$ be the generating series for r^2 -symmetric convex polyominoes of even width. Then we have*

$$F_{r^2,e}(x, y, q) = \frac{2}{1-y} \left(D\left(\frac{1}{y}, x^2, y^2, q^2\right) - D(1, x^2, y^2, q^2) \right) - T(x^2, y, q^2). \quad (45)$$




Figure 11: An r^2 -symmetric convex polyomino P and its fundamental region Q .

Proof. Let P be an r^2 -symmetric convex polyomino of even width. We define the fundamental region of P to be the right half of P . Call this polyomino Q (see Figure 11). Three cases can occur according to the privileged direction taken by Q with respect to P .

In the first case, the fundamental region Q is directed North-East (Figure 11). To recover P , we rotate a copy of Q by 180° and glue the result \bar{Q} to Q along the leftmost column. If this column has length equal to k , there will be k possible positions for \bar{Q} relative to Q . The substitution $s \mapsto 1/y$, $x \mapsto x^2$, $y \mapsto y^2$ and $q \mapsto q^2$ in the generating series $D(s, x, y, q)$ of directed convex polyominoes corresponds to the highest position of \bar{Q} , which minimizes the overall height of P . All the possible positions will be accounted for by multiplying by $(1 + y + \dots + y^{k-1})$. In other words, the substitution to make in $D(s, x^2, y^2, q^2)$ is

$$s^k \mapsto \frac{1 + y + \dots + y^{k-1}}{y^k} = \frac{1}{1 - y} \left(\frac{1}{y^k} - 1 \right). \quad (46)$$

Summing over all possible k 's, we find that the generating series for r^2 -symmetric convex polyominoes in the first class is

$$\frac{1}{1 - y} \left(D\left(\frac{1}{y}, x^2, y^2, q^2\right) - D(1, x^2, y^2, q^2) \right) \quad (47)$$

In the second case, the fundamental region Q is directed South-East. Using an horizontal reflection, we see that the r^2 -symmetric convex polyominoes in this class are also enumerated by (47).

In the third case, the fundamental region Q does not take a definite North or South direction. Q is then a stack polyomino, and the rotated image \bar{Q} is glued to Q in the minimum height position (see Figure 12). Hence these r^2 -symmetric convex polyominoes are enumerated by $T(x^2, y, q^2)$. However, this class is already contained both in the first and the second classes, so that this term must in fact be subtracted from the others. \square

Proposition 14 *The generating series $F_{r^2, o}(x, y, q)$ for convex r^2 -symmetric polyominoes with odd width is given by*

$$F_{r^2, o}(x, y, q) = \frac{2}{x} D\left(\frac{1}{yq}, x^2, y^2, q^2\right) - \frac{1}{x} T\left(x^2, \frac{y}{q}, q^2\right). \quad (48)$$




Figure 12: Convex r^2 -symmetric polyomino whose fundamental region is a stack.

Proof. The proof is similar to the previous one. The main difference is that only one glueing position of \bar{Q} to Q is admissible and that furthermore the leftmost column of Q and its rotated image in \bar{Q} are superimposed to yield an odd width (see Figure 13). Details are left to the reader.

□




Figure 13: Convex r^2 -symmetric polyomino with odd width.

We give here the first few terms of the generating series $F_{r,2}(t, t, q)$ according to half-perimeter and area for r^2 -symmetric convex polyominoes:

$$F_{r,2}(t, t, q) = t^2q + 2t^3q^2 + 2t^4q^3 + t^4q^4 + 6t^5q^4 + 2t^5q^6 + 7t^6q^5 + \dots \quad (49)$$

3.3 Rotation-type convex polyominoes

We can now apply formula (27) to the case where \mathcal{P} is the class C of convex (translation-type) polyominoes and G is the subgroup \mathfrak{C}_4 of \mathfrak{D}_4 , of rotations, since rotation-type polyominoes are precisely defined as orbits of polyominoes relative to \mathfrak{C}_4 . We thus get

$$|C/\mathfrak{C}_4| = \frac{1}{4} (|\text{Fix}(1)| + |\text{Fix}(r)| + |\text{Fix}(r^2)| + |\text{Fix}(r^3)|) \quad (50)$$

Note that $|\text{Fix}(1)|$ is the generating series $C(t, t, q)$ of convex polyominoes (see Subsection 2.1.4). $|\text{Fix}(r)|_w = |\text{Fix}(r^3)|_w = F_r(t, q)$ is given by equation (33) or equation (37) and $|\text{Fix}(r^2)|_w = F_{r,2}(t, t, q)$ is given by equation (44).

Proposition 15 *The half-perimeter and area generating series for convex rotation-type polyominoes $(C/\mathfrak{C}_4)(t, q)$ is given by*

$$(C/\mathfrak{C}_4)(t, q) = \frac{1}{4} (C(t, t, q) + 2F_r(t, q) + F_{r,2}(t, t, q)) . \quad (51)$$

□

The first few terms of $(C/\mathfrak{C}_4)(t, q)$ are given in Appendix B.

4 Horizontal and vertical symmetry classes

4.1 Horizontal or vertical symmetry

We first note that there is an obvious bijection between the horizontally and vertically symmetric polyominoes: rotation by a right angle. We can thus restrict our study to the v -symmetric polyominoes (those left unchanged by a reflection about some vertical axis).

Define the fundamental region of a v -symmetric convex polyomino P to be the part Q of P on the right hand side of the symmetry axis (including the central column in the odd width case). This region is a stack polyomino (see Figure 14).

Proposition 16 *Let $F_v(t, q)$ be the generating series for the v -symmetric convex polyominoes. Then*

$$F_v(t, q) = T(t^2, t, q^2) + \frac{1}{t}T(t^2, \frac{t}{q}, q^2), \quad (52)$$

where $T(x, y, q)$ is the generating series for stack polyominoes.

Proof. Let P be a convex v -symmetric polyomino. There are two possibilities:

1. The width of P is even (see Figure 14(a)), in which case the symmetry axis lies between two columns of P . Consider the stack polyomino Q right of the axis. Reflecting it about the axis doubles the width and the area, but the height remains unchanged. Hence the generating series in this case is given by $T(t^2, t, q^2)$.




Figure 14: Convex v -symmetric polyomino with: (a) even width, (b) odd width.

2. The width of P is odd (see Figure 14(b)), in which case the symmetry axis passes through the central column. To construct P , reflect the fundamental region and glue the two parts together, taking care to superimpose the two central columns (identical by symmetry) to get an odd-width polyomino. The correct generating series for this case is then $\frac{1}{t}T(t^2, \frac{t}{q}, q^2)$.

□

Here are the first few terms of the series $F_v(t, q)$:

$$F_v(t, q) = t^2q + 2t^3q^2 + 2t^4q^3 + t^4q^4 + 4t^5q^4 + 2t^5q^6 + 5t^6q^5 + \dots \quad (53)$$

4.2 Horizontal and vertical symmetry

Let us examine now the set of convex polyominoes that are both horizontally and vertically symmetric. Such a polyomino P will be called a hv -symmetric polyomino. Define the fundamental region Q of P to be the bottom-right part of P , that is the part right of the vertical symmetry axis and down the horizontal symmetry axis. The region Q is a partition polyomino (see Figure 15 and Subsection 2.1.1).

Proposition 17 *The generating series $F_{hv}(t, q)$ for the convex hv -symmetric polyominoes is given by*

$$F_{hv}(t, q) = P(t^2, t^2, q^4) + 2\frac{1}{t}P\left(\frac{t^2}{q^2}, t^2, q^4\right) + \frac{q}{t^2}P\left(\frac{t^2}{q^2}, \frac{t^2}{q^2}, q^4\right), \quad (54)$$

where $P(x, y, q)$ is the generating series for partition polyominoes.

Proof. Let P be a convex hv -symmetric polyomino. P falls into one of four disjoint cases, according to the parities of its height and width.

1. P has even height and even width. Then we simply have to glue together four copies of the fundamental region Q (see Figure 15). Replacing x and y by t^2 we will get the correct




Figure 15: Convex hv -symmetric polyomino with even width and even height.

perimeter value for the whole object, and replacing q by q^4 will give the right area. So the generating series $F_{hv,ee}(t, q)$ for this type of polyominoes is

$$F_{hv,ee}(t, q) = P(t^2, t^2, q^4). \quad (55)$$

2. P has even width and odd height. Then the horizontal symmetry axis passes through the middle row (see Figure 16(a)). We first duplicate four times the fundamental region, glue them together as above, but merge the two central rows. The correct adjustment requires replacing x by t^2/q^2 and dividing by t . The generating series $F_{hv,eo}(t, q)$ for these polyominoes is thus

$$F_{hv,eo}(t, q) = \frac{1}{t}P\left(\frac{t^2}{q^2}, t^2, q^4\right). \quad (56)$$

3. P has odd width and even height. Using a 90° rotation, we see that this case reduces to the preceding one by interchanging x and y in $P(x, y, q)$. Since $P(x, y, q)$ is symmetric in x and y , the generating series $F_{hv,oe}(t, q)$ for this case is also given by

$$F_{hv,oe}(t, q) = \frac{1}{t}P\left(\frac{t^2}{q^2}, t^2, q^4\right). \quad (57)$$




Figure 16: Convex hv -symmetric polyomino with odd height and: (a) even width, (b) odd width.

4. P has odd width and odd height (see Figure 16(b)). The generating series $F_{hv,oo}(t, q)$ for this class of polyominoes is given by

$$F_{hv,oo}(t, q) = \frac{q}{t^2} P\left(\frac{t^2}{q^2}, \frac{t^2}{q^2}, q^4\right). \quad (58)$$

Summing the four series yields (54).

□

Let $f_n^{ee}(q)$, $f_n^{eo}(q)$, $f_n^{oe}(q)$ and $f_n^{oo}(q)$ denote the area generating polynomials for these classes of hv -symmetric polyominoes with half-perimeter equal to n . In other words,

$$F_{hv,ee}(t, q) = \sum_{n \geq 1} f_n^{ee}(q) t^n, \quad F_{hv,eo}(t, q) = \sum_{n \geq 1} f_n^{eo}(q) t^n,$$

$$F_{hv,oe}(t, q) = \sum_{n \geq 1} f_n^{oe}(q) t^n, \quad F_{hv,oo}(t, q) = \sum_{n \geq 1} f_n^{oo}(q) t^n.$$

Obviously, we have, for $n \geq 1$,

$$f_{n+2}^{ee}(q) = q^{2n-1} f_n^{oo}(q), \quad (59)$$

$$f_n^{oe}(q) = f_n^{eo}(q), \quad (60)$$

so that we can concentrate our attention on the polynomials $f_n^{oo}(q)$ and $f_n^{eo}(q)$.

Proposition 18 *The area generating polynomials $f_n^{oo}(q)$ and $f_n^{eo}(q)$ are given by*

$$f_n^{oo}(q) = \begin{cases} q^{n-1} \sum_{i=0}^{(n-2)/2} \begin{bmatrix} (n-2)/2 \\ i \end{bmatrix}_{q^4} & \text{if } n \text{ is even,} \\ 0 & \text{if } n \text{ is odd,} \end{cases} \quad (61)$$

$$f_n^{eo}(q) = \begin{cases} q^{n-1} \sum_{i=0}^{(n-3)/2} q^{2i} \begin{bmatrix} (n-3)/2 \\ i \end{bmatrix}_{q^4} & \text{if } n \text{ is odd,} \\ 0 & \text{if } n \text{ is even.} \end{cases} \quad (62)$$

Moreover they satisfy the following recurrences: for $n \geq 5$,

$$f_n^{oo}(q) = 2q^2 f_{n-2}^{oo}(q) - q^4(1 - q^{2n-8}) f_{n-4}^{oo}(q), \quad (63)$$

$$f_n^{eo}(q) = q^2(1 + q^2) f_{n-2}^{eo}(q) - q^6(1 - q^{2n-10}) f_{n-4}^{eo}, \quad (64)$$

with initial conditions

$$f_1^{oo}(q) = 0, \quad f_2^{oo}(q) = q, \quad f_3^{oo}(q) = 0, \quad f_4^{oo}(q) = 2q^3,$$

$$f_1^{eo}(q) = 0, \quad f_2^{eo}(q) = 0, \quad f_3^{eo}(q) = q^2, \quad f_4^{eo}(q) = 0.$$

Proof. The explicit formulas (61) and (62) can be easily derived from the generating series or from a direct combinatorial argument. The recurrences (63) and (64) can be checked by substituting the explicit formulas. \square

Corollary 19 *The number f_n of hv -symmetric convex polyominoes with half-perimeter n is 0 if $n = 1$, 1 if $n = 2$, and for $n \geq 3$, is given by*

$$f_n = f_n^{oo}(1) + f_n^{ee}(1) = 3 \cdot 2^{(n/2)-2} \quad \text{for } n \text{ even}, \quad (65)$$

$$f_n = f_n^{eo}(1) + f_n^{oe}(1) = 2^{(n-1)/2} \quad \text{for } n \text{ odd}. \quad (66)$$

Furthermore, we have

$$F_{hv}(t, 1) = \sum_{n \geq 1} f_n t^n = \frac{t(t+1)^2}{1-2t^2}. \quad (67)$$

\square

We give here the first few terms of the series $F_{hv}(t, q)$:

$$F_{hv}(t, q) = t^2 q + 2t^3 q^2 + 2t^4 q^3 + t^4 q^4 + 2t^5 q^4 + 2t^5 q^6 + 3t^6 q^5 + \dots \quad (68)$$

5 Diagonal symmetry classes

5.1 Diagonally symmetric convex polyominoes

Our next goal is to compute the generating series $F_d(t, q)$ of d -symmetric convex polyominoes. There is a simple bijection between d_1 -symmetric and d_2 -symmetric polyominoes, given by the 90° rotation r , so that $F_{d_1}(t, q) = F_{d_2}(t, q)$. We choose $d = d_1$, the diagonal along the line $y = -x$ and define the *fundamental region* Q of a d -symmetric convex polyomino P as the part of P located on the North-East side of the diagonal symmetry line, including the cells on the diagonal (see Figure 17).

Observe that the fundamental region Q of a d -symmetric convex polyomino P is a (diagonally) shifted directed (diagonal source) convex polyomino, called D_S -polyomino for short. These should not be confused with the neighboring class of directed diagonally convex polyominoes, also known as fully directed compact lattice animals (see [30] and [31]), and a different enumerating method has to be used.

We introduce an extra variable z to mark the length of the diagonal source. For example, the D_S -polyomino in Figure 17(b) has global weight $x^6 y^5 z^3 q^{17}$. Let $D_S(x, y, z, q)$ denote the generating series of D_S -polyominoes. Before computing this series, we can state the following:




Figure 17: Diagonally symmetric convex polyomino P and its fundamental region Q .

Proposition 20 *The half-perimeter and area generating series $F_d(t, q)$ of d -symmetric convex polyominoes is given by*

$$F_d(t, q) = D_S(t^2, t^2, \frac{1}{t^2q}, q^2), \quad (69)$$

where $D_S(x, y, z, q)$ is the generating series of shifted directed convex polyominoes.

Proof. Observe that if the fundamental region Q of a d -symmetric polyomino P has width m , height n , diagonal source length k and area a , then P has half-perimeter equal to $2m + 2n - 2k$ and area equal to $2a - k$. This is achieved precisely by the substitution $x = t^2$, $y = t^2$, $z = 1/t^2q$ and $q = q^2$ as in formula (69). \square

Here are the first few terms:

$$F_d(t, q) = t^2q + 2t^4q^3 + t^4q^4 + 5t^6q^5 + 4t^6q^6 + 2t^6q^7 + 2t^6q^8 + \dots \quad (70)$$

5.2 Shifted directed convex (D_S) polyominoes

We now outline some basic enumerative results concerning the class D_S of shifted directed convex polyominoes. This class is studied in more detail in a separate paper [17].

Let Q be a D_S -polyomino. Imagining Q as a kind of floating device, we define the *flotation* row of Q as the (horizontal) row containing the top diagonal source cell. The two bounding lines, below and above the flotation row are called respectively the *flotation line* and the *security line* of Q . We also define the *North-East corner* of Q as the topmost cell of the rightmost column of Q (see Figure 18).

We now partition the class of D_S -polyominoes into two subclasses denoted by Y_1 and Y_2 , according to the relative position of the North-East corner with respect to the flotation line: above for Y_1 and below for Y_2 . We also denote their generating series by the same names and we have

$$D_S(x, y, z, q) = Y_1(x, y, z, q) + Y_2(x, y, z, q). \quad (71)$$

In order to enumerate Y_1 -polyominoes, we introduce a new variable v to mark the width of the bottom row. For example, the polyomino Q of Figure 17(b) has bottom width equal to 4. Let $Y_1(v, x, y, z, q)$ denote the corresponding generating function. Observe that the (one source) directed convex polyominoes are a special case. The corresponding generating function is obtained

from $D(s, x, y, q)$ (see Subsection 2.1.4 and [4]) by setting $s = v$, applying a diagonal reflection. Extracting the coefficient of z in Y_1 gives

$$D(v, x, y, q) = [z] Y_1(v, x, y, z, q). \quad (72)$$

Now any D_S -polyomino Q in the class Y_1 can be constructed by glueing together along the flotation line a directed convex polyomino D , of bottom width $m \geq 1$, with a (possibly empty) diagonally decreasing stack of rows, i.e. a shifted partition P_S with parts of length less than m (see Figure 18(a)). This gives the following:

Proposition 21 *Let $D_m(x, y, q) = [v^m] D(v, x, y, q)$ be the generating series for directed convex polyominoes of bottom width m . We then have*

$$Y_1(x, y, z, q) = \sum_{m \geq 1} z D_m(x, y, q) (1 + yzq) \cdots (1 + yzq^{m-1}). \quad (73)$$

□

It is possible to get a more closed form for this generating function, using the following functional equation:

Proposition 22 *The generating function $Y_1(v) = Y_1(v, x, y, z, q)$ of Y_1 -polyominoes is characterized by the equation*

$$Y_1(v) = zD(v, x, y, q) + \frac{yz}{1 - vq} (vqY_1(1) - Y_1(vq)). \quad (74)$$

Proof. This equation says that any Y_1 -polyomino can be obtained as follows: start with a directed convex polyomino D and then successively glue, on the bottom, rows that follow the diagonal on the left hand side but do not exceed the previous row on the right hand side. The term $yzvq(1 - vq)^{-1}Y_1(1)$ accounts for the addition of a row of arbitrary length whereas $yz(1 - vq)^{-1}Y_1(vq)$ enumerates the cases where the added row does in fact exceed on the right hand side, and thus should be subtracted. □

Corollary 23 *The generating series for Y_1 -polyominoes is given by*

$$Y_1(x, y, z, q) = \frac{\sum_{n \geq 0} zD(q^n, x, y, q) \frac{(-yz)^n}{(q)_n}}{1 + \sum_{n \geq 1} q^{n-1} \frac{(-yz)^n}{(q)_n}}. \quad (75)$$

Proof. This is the solution $Y_1(1)$ of equation (74) obtained by iteration. See Lemma 2.3 of [4]. □

We now study the subclass Y_2 of D_S -polyominoes. Let us introduce a variable u to mark the top row width and let $Y_2(u, x, y, z, q)$ be the corresponding generating series. Cutting a Y_2 -polyomino P along the security line yields a stack polyomino T on top and a shifted stack polyomino T_S (see Subsection 2.1.3) below the security line (see Figure 18(b)). Note that T_S is not a general diagonal row stack polyomino since the North-East corner is not allowed to be on the top row. Call R the subclass of T_S satisfying this restriction and observe that T_S -polyominoes whose North-East corner is on the top row are precisely shifted partitions (see Subsection 2.1.1). We then have

$$R(u, x, y, q) = T_S(u, x, y, q) - P_S(xu, y, q). \quad (76)$$




Figure 18: D_S -polyominoes of types Y_1 (case (a)) and Y_2 (case (b)).

Let $R_m(x, y, q)$ be the generating series of R -polyominoes having top width equal to m , i.e. set

$$R(u, x, y, q) = \sum_{m \geq 1} R_m(x, y, q) u^m. \quad (77)$$

Also denote by $T_{0,m}(y, q)$ the generating series of stack polyominoes of width m and having possibly empty columns (see Subsection 2.1.2; note that these stacks have been rotated by 90°). We then have

Proposition 24 *The generating series for Y_2 -polyominoes is given by*

$$Y_2(x, y, z, q) = \sum_{m \geq 1} R_m(x, yz, q) T_{0,m}(y, q). \quad (78)$$

□

An alternate expression can be obtained for Y_2 by solving the following functional equation for $Y_2(u, x, y, z, q)$, with the help of Lemma 2.2 of [4].

Proposition 25 *The series $Y_2(u) = Y_2(u, x, y, z, q)$ satisfies the functional equation*

$$Y_2(u) = R(u, x, yz, q) + \frac{yuq}{1 - uq} Y_2'(1) - \frac{yu^2 q^2}{(1 - uq)^2} (Y_2(1) - Y_2(uq)), \quad (79)$$

where the derivative $Y_2'(u)$ is taken with respect to u .

Proof. This functional equation states in fact that a Y_2 -polyomino can be obtained as follows: start with a R -polyomino and then glue successively rows to the top. Each added row should not exceed the previous row on either side. This corresponds to the construction C of [4] where the general functional equation is stated and solved. □

Here are the first few terms of D_S :

$$D_S(x, y, z, q) = xyzq + x^2 yzq^2 + xy^2 zq^2 + x^3 yzq^3 + x^2 y^2 z^2 q^3 + 3x^2 y^2 zq^3 + xy^3 zq^3 + \dots \quad (80)$$

5.3 Symmetry with respect to both diagonals

A polyomino which is symmetric with respect to both diagonals is called d_1d_2 -symmetric. Let P be a d_1d_2 -symmetric convex polyomino. Observe that the height of P must be equal to its width. Two cases occur according to the parity of the width. Let $F_{d_1d_2,e}(t, q)$ and $F_{d_1d_2,o}(t, q)$ denote the half-perimeter and area generating functions for even and odd width d_1d_2 -symmetric convex polyominoes respectively. We then have

$$F_{d_1d_2}(t, q) = F_{d_1d_2,e}(t, q) + F_{d_1d_2,o}(t, q). \quad (81)$$

We define the *fundamental region* Q of P to be the part of P which is bounded by the two diagonals and extends in the East direction, including the bounding diagonal cells (see Figure 19). Observe that Q is a stack of rows with a double diagonal shift. We call these polyominoes *doubly shifted stacks* (T_{SS}) and qualify them as *even* in the even width case (Figure 19(a)) and *acute* in the odd case (Figure 19(b)). We use the variables z to mark the length of the d_1 -diagonal and w for the d_2 -diagonal, and introduce the generating series $E(x, z, w, q)$ and $A(x, z, w, q)$ of even and acute (respectively) doubly shifted stacks. Note that in the acute case, the extreme West cell contributes to both diagonal lengths. Before determining these series, we can state the following:

Proposition 26 *The generating series for even and odd d_1d_2 -symmetric convex polyominoes are given by*

$$F_{d_1d_2,e}(t, q) = E(t^4, \frac{1}{q^2}, \frac{1}{q^2}, q^4), \quad (82)$$

$$F_{d_1d_2,o}(t, q) = \frac{q}{t^2} A(t^4, \frac{1}{q^2}, \frac{1}{q^2}, q^4). \quad (83)$$




Figure 19: Convex d_1d_2 -symmetric polyomino with: (a) even width; (b) odd width.

Proof. Suppose that the doubly shifted stack Q has width m , d_1 and d_2 diagonal lengths n_1 and n_2 respectively, and area a . Then, in the even case, P will have half-perimeter $4m$ and area $4a - 2n_1 - 2n_2$, while in the odd case, the half-perimeter will be $4m - 2$ and the area, $4a - 2n_1 - 2n_2 + 1$. Since the correspondence between P and Q is bijective, we have (82) and (83). \square

We now concentrate on the computation of the generating series $E(x, z, w, q)$ and $A(x, z, w, q)$ of doubly shifted stacks. The concepts of flotation line, security line, North-East and South-East corners are defined similarly to the case of D_S -polyominoes. In the even case, we then introduce three subclasses of doubly shifted stacks as follows: E_1 if the South-East corner is below the security line; E_2 when the North-East corner is above the security line; and E_3 when both conditions are satisfied (see Figure 20). We obviously have

$$E(x, z, w, q) = E_1(x, z, w, q) + E_2(x, z, w, q) - E_3(x, z, w, q). \quad (84)$$




Figure 20: The three classes of even doubly shifted stacks.

Observe that under the security line of an E_1 -polyomino Q lies a (single) shifted stack polyomino T_S whereas on top of the security line is found a non-empty partition P_S in distinct parts (rows). If T_S has top width equal to m , then P_S has biggest row at most m .

Let $T_{S,m}(x, y, q) = [u^m]T_S(u, x, y, q)$ be the generating series for shifted stacks having top width equal to m (see Subsection 2.1.2). We then have

Proposition 27 *The generating series for even doubly shifted stacks E_1 , E_2 and E_3 are given by*

$$E_1(x, z, w, q) = \sum_{m \geq 1} T_{S,m}(x, z, q)((-wq)_m - 1), \quad (85)$$

$$E_2(x, z, w, q) = E_1(x, w, z, q), \quad (86)$$

$$E_3(x, z, w, q) = \sum_{m \geq 1} x^m z w q^{2m} (-zq)_{m-1} (-wq)_{m-1}. \quad (87)$$

Proof. Formula (85) follows from the previous discussion. To see (86), simply observe that an horizontal reflection will send any E_2 -polyomino into an E_1 -polyomino and vice-versa. Finally, the closed form (87) for $E_3(x, z, w, q)$ is obvious by construction. \square

Note that a more closed form can be given for E_1 , using the following functional equation. Let the variable u mark the top row width of any convex polyomino and let $E_1(u) = E_1(u, x, z, w, q)$ and $T_S(u) = T_S(u, x, z, q)$ be the corresponding generating series.

Proposition 28 *The generating series $E_1(u)$ of even doubly shifted stacks of the first class satisfies the functional equation*

$$E_1(u) = \frac{uw}{1-uw} (T_S(1) - T_S(uq)) + \frac{w}{1-uw} (uqE_1(1) - E_1(uq)) . \quad (88)$$

Proof. The first term on the right hand side of (88) has the effect of constructing the first row of the d_2 -diagonal, directly above the base T_S , while the second term iteratively constructs the other rows, following the diagonal. \square

In the odd case, we similarly introduce the following three subclasses of acute doubly shifted stacks: A_1 if the South-East corner is below the security line; A_2 if the North-East corner is above the flotation line; and A_3 if both conditions hold (see Figure 21). We obviously have

$$A(x, z, w, q) = A_1(x, z, w, q) + A_2(x, z, w, q) - A_3(x, z, w, q) \quad (89)$$




Figure 21: The three classes acute doubly shifted stacks.

The following results are similar to Propositions 27 and 28 above. Proofs are left to the reader.

Proposition 29 *The generating series for acute doubly shifted stacks A_1 , A_2 and A_3 are given by*

$$A_1(x, z, w, q) = \sum_{m \geq 1} T_{S,m}(x, z, q) (-wq)_{m-1} , \quad (90)$$

$$A_2(x, z, w, q) = A_1(x, w, z, q) , \quad (91)$$

$$A_3(x, z, w, q) = \sum_{m \geq 1} x^m z w q^m (-zq)_{m-1} (-wq)_{m-1} . \quad (92)$$

\square

Proposition 30 *The generating series $A_1(u) = A_1(u, x, z, w, q)$ of acute doubly shifted stacks of the first class satisfies the functional equation*

$$A_1(u) = wT_S(u, x, z, q) + \frac{w}{1-uw} (uqA_1(1) - A_1(uq)) . \quad (93)$$

\square

Finally, putting together formulas (81)–(92) yields $F_{d_1 d_2}(t, q)$. Here are the first few terms:

$$F_{d_1 d_2}(t, q) = t^2 q + t^4 q^4 + t^6 q^5 + 2 t^6 q^7 + t^6 q^9 + 2 t^8 q^8 + 2 t^8 q^{10} + \dots \quad (94)$$

6 Congruence-type convex polyominoes

We are now in a position to enumerate congruence-type convex polyominoes, i.e. convex polyominoes up to rotation and reflection using formula (27) of Section 2.2 with $G = \mathfrak{D}_4$ and $\mathcal{P} = C$, the class of all convex polyominoes:

$$|C/\mathfrak{D}_4|_w = \frac{1}{8} \sum_{g \in \mathfrak{D}_4} |\text{Fix}(g)|_w, \quad (95)$$

where $|\text{Fix}(g)|_w$ is the half-perimeter and area generating series of the convex g -symmetric polyominoes. Therefore,

Proposition 31 *The half-perimeter and area generating series $(C/\mathfrak{D}_4)(t, q)$ of congruence-type polyominoes is given by*

$$(C/\mathfrak{D}_4)(t, q) = |C/\mathfrak{D}_4|_w = \frac{1}{8} (C(t, t, q) + 2F_r(t, q) + F_{r^2}(t, t, q) + 2F_d(t, q) + 2F_v(t, q)). \quad (96)$$

□

The first few terms of $(C/\mathfrak{D}_4)(t, q)$ are given in Appendix B.

7 Asymmetric convex polyominoes

We can also enumerate asymmetric convex polyominoes, i.e. polyominoes having no symmetry at all, using formula (31) of Section 2.2 and the value of the Möbius function for the subgroups of \mathfrak{D}_4 given in Figure 7.

Proposition 32 *The half-perimeter and area generating series $\overline{C}(t, q) = F_{=0}$ of asymmetric convex polyominoes is given by*

$$\begin{aligned} \overline{C}(t, q) &= F_{\geq 0} - F_{\geq d_1} - F_{\geq d_2} - F_{\geq r^2} - F_{\geq h} - F_{\geq v} + 2F_{\geq \langle d_1, d_2 \rangle} + 2F_{\geq \langle h, v \rangle} \\ &= C(t, t, q) - 2F_d(t, q) - F_{r^2}(t, t, q) - 2F_v(t, q) + 2F_{d_1 d_2}(t, q) + 2F_{hv}(t, q). \end{aligned} \quad (97)$$

The first few terms of this series are given in Appendix B.

Note that the same method would allow us to enumerate the convex polyominoes having exactly the symmetries of any given subgroup of \mathfrak{D}_4 .

As an application we show the asymptotic result that for large area, almost all convex polyominoes are asymmetric. In other words, the probability for a convex polyomino to have at least one symmetry goes to zero as the area goes to infinity.

Proposition 33 (Bender [2]) *Let c_n be the number of convex polyominoes with area n . Then*

$$c_n \sim k \mu^n, \quad (98)$$

with

$$k = 2.67574\dots \quad \mu = 2.30913859330\dots$$

□

Lemma 34 *Let H be any non-trivial subgroup of \mathfrak{D}_4 and denote by $F_{H,n}$ the number of H -symmetric polyominoes with area n . Then,*

$$\lim_{n \rightarrow \infty} \frac{F_{H,n}}{c_n} = 0. \quad (99)$$

Proof. A same basic argument works for r^2 -, h -, v -, d_1 - and d_2 -symmetric convex polyominoes, using the fact that they are all constructed from two congruent subpolyominoes. A supplementary column or row has to be added in some cases, according to whether the height or the width of the initial polyomino is odd or even. These subpolyominoes are convex in every case, and they have at most half the area of the initial object.

- **r^2 -symmetric convex polyominoes :** An r^2 -symmetric convex polyomino with even width and area n is constructed from two congruent convex subpolyominoes with exactly half the area, which can be glued together in at most $n/2$ ways (the maximal height of the columns that get glued together). So $F_{r^2,n}^{even} \leq \frac{1}{2}nc_{n/2}$. Hence

$$\lim_{n \rightarrow \infty} \frac{F_{r^2,n}^{even}}{c_n} \leq \lim_{n \rightarrow \infty} \frac{\frac{1}{2}nk\mu^{n/2}}{k\mu^n} = 0.$$

Next consider an r^2 -symmetric convex polyomino with odd width and area n . This polyomino is constructed from a central column (n choices of height) and two congruent convex subpolyominoes of area at most $\lfloor n/2 \rfloor$. Then there are at most n possible positions where to glue the subpolyominoes to the central column (they are glued symmetrically). Thus $F_{r^2,n}^{odd} \leq n^2(1 + c_1 + c_2 + \dots + c_{\lfloor n/2 \rfloor}) < n^3c_{\lfloor n/2 \rfloor}$ and the result follows as above. Hence the result holds for the subgroup $\langle r^2 \rangle$ of \mathfrak{D}_4 ;

- **h - and v -symmetric convex polyominoes :** A v -symmetric convex polyomino of even width is constructed much in the same way as an r^2 -symmetric convex polyomino of even width, with the difference that there is only one glueing position for the two subpolyominoes. So $F_{v,n}^{even} \leq c_{n/2}$ and the limit will be zero. Similarly, the construction for v -symmetric convex polyominoes of odd width resembles that of r^2 -symmetric convex polyominoes of odd width, and we get $F_{v,n}^{odd} \leq n^2(1 + c_1 + c_2 + \dots + c_{\lfloor n/2 \rfloor}) < n^3c_{\lfloor n/2 \rfloor}$. So the result holds for the subgroups $\langle h \rangle$ and $\langle v \rangle$;
- **d_1 - and d_2 -symmetric convex polyominoes :** Let P be a d -symmetric convex polyomino and Q its fundamental region. Suppose that P has b cells on the diagonal symmetry axis. Then the minimum area P can have is $b + 2(b - 1)$. This gives a minimum area of $b + (b - 1)$ for Q . Hence

$$\frac{\text{Area of } Q_{min}}{\text{Area of } P_{min}} = \frac{2b - 1}{3b - 2}.$$

Then if we add to Q a cell not on the diagonal symmetry axis, two cells get added to P , and thus we conclude that the ratio can only decrease as we make P into a larger d -symmetric convex polyomino with the same number of cells on the diagonal axis. For $b \geq 2$, the ratio will be smaller than or equal to $3/4$. As a loose approximation, we can take Q to be any convex polyomino. This gives $F_{d,n} \leq c_{\lfloor 3n/4 \rfloor} + c_{\lfloor n/2 \rfloor}$. The $c_{\lfloor n/2 \rfloor}$ term corresponds to those d -symmetric convex polyominoes having only one cell on the diagonal axis (in which case the fundamental region is a convex polyomino of area $(n + 1)/2$). Hence the result will also hold for the subgroups $\langle d_1 \rangle$ and $\langle d_2 \rangle$.

Then since r -symmetric polyominoes are a subclass of r^2 -symmetric polyominoes, hv -symmetric polyominoes a subclass of h - or v -symmetric polyominoes and d_1d_2 -symmetric polyominoes a subclass of d_1 - or d_2 -symmetric polyominoes, the results extends to all of the symmetry classes we considered. \square

Proposition 35 *If we denote by \bar{c}_n the number of asymmetric convex polyominoes of area n , then*

$$\bar{c}_n \sim c_n. \quad (100)$$

Proof. We get the result from equation (97) and the previous lemma. \square

References

- [1] Andrews, G. E., *The Theory of Partitions*, Encyclopedia of Mathematics and its Applications, Addison Wesley Pub. Comp., Reading, Mass., **1976**.
- [2] Bender, E. A., *Convex n -ominoes*, Discrete Math., 8, **1974**, 219–226.
- [3] Bergeron, F., G. Labelle and P. Leroux, *Combinatorial Species and Tree-like Structures*, Cambridge University Press, Series Encyclopedia of Mathematics and its Applications, 67, **1997**.
- [4] Bousquet-Mélou, M., *A Method for the Enumeration of Various Classes of Column-convex Polygons*, Discrete Math., 154, **1996**, 1–25.
- [5] Bousquet-Mélou, M., *Percolation Models and Animals*, European J. Combin., 17, **1996**, 343–369.
- [6] Bousquet-Mélou, M., *Rapport scientifique d’habilitation*, LaBRI, Université Bordeaux 1, **déc. 1996**.
- [7] Bousquet-Mélou, M. and J.-M. Fédo, *The generating function of convex polyominoes: the resolution of a q -differential system*, Discrete Math, 137, **1995**, 53–75.
- [8] Bousquet-Mélou, M. and A. J. Guttmann, *Enumeration of three-dimensional convex polygons*, Annals of Combinatorics, 1, **1997**, 27–53.
- [9] Bousquet-Mélou, M. and X. G. Viennot, *Empilements de segments et q -énumération de polyominos convexes dirigés*, J. Combin. Theory Ser A, 60, **1992**, 196–224.
- [10] Delest, M.-P., *Generating Functions for Column-convex Polyominoes*, J. Combin. Theory Ser A, 48, **1988**, 12–31.
- [11] Delest, M.-P. and G. Viennot, *Algebraic languages and polyomino enumeration*, Theoret. Comput. Sci., 34, **1984**, 169–206.
- [12] Feretić, S. and D. Svrtan, *On the number of column-convex polyominoes with given perimeter and number of columns*, in: *Proceedings of the 5th Conference on Formal Power Series and Algebraic Combinatorics*, editors: A. Barlotti, M. Delest, and R. Pinzani, Florence, june **1993**, 201–214.
- [13] Gessel, I., *On the number of convex polyominoes*, Manuscript, **1990**.
- [14] Golomb, S. W., *Polyominoes: Puzzles, Patterns, Problems and Packings, Revised Edition*, Princeton University Press, Princeton, **1994**.
- [15] Guttmann, A. J., *On the Number of Lattice Animals Embeddable in the Square Lattice*, J. Phys. A: Math. Gen., 15, **1982**, 1987–90.
- [16] Guttmann, A. J. and I. G. Enting, *The number of convex polygons on the square and honeycomb lattices*, J. Phys. A: Math. Gen., 21, **1988**, L467–L474.

- [17] Guttmann, A. J. and P. Leroux, *Diagonal source directed convex polyominoes*. (In preparation).
- [18] Guttmann, A. J., T. Prellberg and A. L. Owczarek, *On the symmetry classes of planar self-avoiding walks*, J. Phys. A: Math. Gen., 26, **1993**, 6615–6623.
- [19] Hughes, B. D., *Random Walks and Random Environments. Volume 1: Random Walks*. Clarendon Press, Oxford, **1995**.
- [20] Hughes, B. D., *Random Walks and Random Environments. Volume 2: Random Environments*. Clarendon Press, Oxford, **1996**.
- [21] Kerber, A., *Enumeration under Finite Group Action: Symmetry Classes of Mappings*, in: *Combinatoire énumérative*, editors: G. Labelle et P. Leroux, Lectures Notes in Mathematics, 1234, Springer-Verlag, **1986**, 160–176.
- [22] Kim, D., *The number of convex polyominoes with given perimeter*, Discrete Math, 70, **1988**, 47–51.
- [23] Klarner, D. A., *My Life Among the Polyominoes*, in: *The Mathematical Gardner*, editor: D. A. Klarner, Prindle, Weber and Schmidt, **1981**.
- [24] Klarner, D. A. and R. L. Rivest, *Asymptotic bounds for the number of convex n -ominoes*, Discrete Math, 8, **1974**, 31–40.
- [25] Langlands, R., P. Pouliot and Y. Saint-Aubin, *Conformal invariance in two-dimensional percolation*, Bulletin of the AMS, 30 #1, **1994**, 1–61.
- [26] Lin, K. Y., *Exact solution of the convex polygon perimeter and area generating function*, J. Phys. A: Math. Gen., 24, **1991**, 2411–2417.
- [27] Lin, K. Y. and S. J. Chang, *Rigorous results for the number of convex polygons on the square and honeycomb lattices*, J. Phys. A: Math. Gen., 21, **1988**, 2635–2642.
- [28] Pólya, G., *On the Number of Certain Lattice Polygons*, J. Combin. Theory, 6, **1969**, 102–105.
- [29] Prellberg, T. and R. Brak, *Critical Exponents from Nonlinear Functional Equations for Partially Directed Cluster Models*, J. Statist. Phys., 78, **1995**, 701–730.
- [30] Privman, V. and N. M. Švrakić, *Exact generating function for fully directed compact lattice animals*, Physical Review Letters, 60, **1988**, 1107–1109.
- [31] Privman, V. and N. M. Švrakić, *Directed Models of Polymers, Interfaces, and Clusters: Scaling and Finite-size Properties*, Lectures Notes in Physics, 338, Springer-Verlag, Berlin, **1989**.
- [32] Rota, G.-C., *On the Foundations of Combinatorial Theory; I. Theory of Möbius Functions*, Zeitschrift für Wahrscheinlichkeitstheorie, Band 2, Heft 4, **1964**, 340–368, reprinted in: Gian-Carlo Rota on Combinatorics, Introductory Papers and Commentaries, editor: J. Kung, Birkhäuser, **1995**, 3–31.
- [33] Temperley, H. N. V., *Statistical Mechanics and the Partitions of Numbers II: The Form of Crystal Surfaces*, Proc. Cambridge Phil. Soc., 48, **1952**, 683–697.
- [34] Temperley, H. N. V., *Combinatorial Problems Suggested by the Statistical Mechanics of Domains and Rubber-like Molecules*, Phys. Rev., 103, **1956**, 1–16.
- [35] Viennot, X. G., *A Survey of Polyomino Enumeration*, in: *Proceedings of the 4th Conference on Formal Power Series and Algebraic Combinatorics*, editors: P. Leroux and C. Reutenauer, Publications du LaCIM, 11, **1992**, 399–420.

Appendix A

The following two tables present the numbers of convex polyominoes according to their symmetry types and their perimeter or area. The columns indexed by subgroups of \mathfrak{D}_4 give the numbers of convex polyominoes of a given perimeter or area that are left fixed by the symmetries of the subgroup. The columns *rotation*, *congruence* and *asym* give respectively the number of rotation-type, congruence-type and asymmetric convex polyominoes of the given size.

per	$\{1\}$	$\langle r \rangle, \langle r^3 \rangle$	$\langle r^2 \rangle$	rotation	$\langle h \rangle, \langle v \rangle$	$\langle d_1 \rangle, \langle d_2 \rangle$	congruence	$\langle h, v \rangle$	$\langle d_1, d_2 \rangle$	asym
4	1	1	1	1	1	1	1	1	1	0
6	2	0	2	1	2	0	1	2	0	0
8	7	1	3	3	3	3	3	3	1	0
10	28	0	8	9	6	0	6	4	0	16
12	120	2	16	35	12	14	24	6	4	72
14	528	0	40	142	24	0	77	8	0	456
16	2344	4	84	609	48	70	334	12	8	2064
18	10416	0	200	2654	96	0	1351	16	0	10056
20	46160	8	424	11650	192	348	5960	24	24	44736

Table 1: Convex polyominoes enumerated by their symmetries and perimeter.

area	$\{1\}$	$\langle r \rangle, \langle r^3 \rangle$	$\langle r^2 \rangle$	rotation	$\langle h \rangle, \langle v \rangle$	$\langle d_1 \rangle, \langle d_2 \rangle$	congruence	$\langle h, v \rangle$	$\langle d_1, d_2 \rangle$	asym
1	1	1	1	1	1	1	1	1	1	0
2	2	0	2	1	2	0	1	2	0	0
3	6	0	2	2	2	2	2	2	0	0
4	19	1	7	7	5	1	5	3	1	8
5	59	1	7	17	5	5	11	3	1	40
6	176	0	24	50	12	4	29	4	0	128
7	502	0	22	131	12	14	72	4	2	440
8	1374	2	74	363	26	12	191	6	2	1240
9	3630	2	62	924	26	38	478	6	2	3452
10	9312	0	208	2380	52	32	1211	6	2	8952

Table 2: Convex polyominoes enumerated by their symmetries and area.

Appendix B

We provide here the beginning terms of the series $(C/\mathfrak{C}_4)(t, q)$, $(C/\mathfrak{D}_4)(t, q)$ and $\overline{C}(t, q)$ of rotation-type, congruence-type and asymmetric polyominoes respectively. They are given up to area 10 (q^{10}).

$$\begin{aligned}(C/\mathfrak{C}_4)(t, q) = & t^2q + t^3q^2 + 2t^4q^3 + t^4q^4 + 6t^5q^4 + 2t^5q^5 + t^5q^6 + 15t^6q^5 + 11t^6q^6 + 6t^6q^7 \\ & + 38t^7q^6 + 2t^6q^8 + 38t^7q^7 + t^6q^9 + 36t^7q^8 + 87t^8q^7 + 18t^7q^9 + 124t^8q^8 + 9t^7q^{10} \\ & + 139t^8q^9 + 201t^9q^8 + 115t^8q^{10} + 334t^9q^9 + 470t^9q^{10} + 432t^{10}q^9 + 861t^{10}q^{10} \\ & + 925t^{11}q^{10} + \dots\end{aligned}$$

$$\begin{aligned}(C/\mathfrak{D}_4)(t, q) = & t^2q + t^3q^2 + 2t^4q^3 + t^4q^4 + 4t^5q^4 + t^5q^5 + t^5q^6 + 10t^6q^5 + 7t^6q^6 + 4t^6q^7 \\ & + 21t^7q^6 + 2t^6q^8 + 19t^7q^7 + t^6q^9 + 20t^7q^8 + 49t^8q^7 + 9t^7q^9 + 65t^8q^8 + 6t^7q^{10} \\ & + 74t^8q^9 + 104t^9q^8 + 62t^8q^{10} + 167t^9q^9 + 239t^9q^{10} + 227t^{10}q^9 + 436t^{10}q^{10} \\ & + 468t^{11}q^{10} + \dots\end{aligned}$$

$$\begin{aligned}\overline{C}(t, q) = & 8t^5q^4 + 8t^5q^5 + 32t^6q^5 + 24t^6q^6 + 16t^6q^7 + 104t^7q^6 + 152t^7q^7 + 104t^7q^8 \\ & + 272t^8q^7 + 72t^7q^9 + 448t^8q^8 + 16t^7q^{10} + 496t^8q^9 + 688t^9q^8 + 400t^8q^{10} \\ & + 1336t^9q^9 + 1744t^9q^{10} + 1552t^{10}q^9 + 3344t^{10}q^{10} + 3448t^{11}q^{10} + \dots\end{aligned}$$