

# Stable map resolutions of Richardson varieties

Allen Knutson (Cornell)

Allensdottirs seminar, September 2020

## Abstract

To a simple normal crossings divisor (sncd)  $D$ , one associates its “dual simplicial complex”, with a vertex for each component  $D_i$  and face  $F$  for each stratum  $\bigcap_{f \in F} D_f \neq \emptyset$ . For example, Escobar’s brick manifolds (which among other things, provide resolutions of Richardson varieties) come with an sncd whose dual complex is a subword complex. In good cases (which includes brick manifolds) the dual complex is a sphere.

With no such geometrical input, Björner-Wachs showed that the order complex of a Bruhat interval  $(u, v)$  is a sphere. I’ll define a space of equivariant stable maps from  $\mathbb{P}^1$  to the Richardson variety  $X_u^v$ , and prove that this space is a smooth orbifold, which comes with a natural sncd whose dual is the Björner-Wachs complex. There are no choices, e.g. of reduced words. In the Grassmannian case this space is GKM, and I describe its GKM graph in terms of rim-hook tableaux.

# Simple normal crossing divisors and their dual complexes.

Let  $D_1, D_2, \dots, D_m$  be a collection of smooth divisors in a (complex, say) manifold  $M$ . They are **simple normal crossings** if  $\bigcap_{f \in F} D_f$  is smooth connected of codimension  $|F|$  (when nonempty) for each  $F \subseteq [m]$ , i.e. rather like a set of coordinate hyperplanes in  $\mathbb{C}^n$ . Their union  $D = D_1 \cup \dots \cup D_m$  is a **simple normal crossings divisor** or **sncd**.

A good test case is  $M = \text{TV}_P$  the projective toric variety associated to a polytope  $P$ , and  $D$  the complement of the open  $T$ -orbit. Then  $\bigcap_{f \in F} D_f$  is always irreducible (when nonempty), but will only have always the right codimension when  $M$  is orbifold, i.e. when  $P$  is “simple”. Consider a pyramid for counterexamples.

Another nonexample is  $M = \mathbb{CP}^2 = \{[x : y : z]\}$ ,  $D_1 = \{x = 0\}$ ,  $D_2 = \{y^2 = xz\}$ . The intersection  $D_1 \cap D_2$  is smooth and codim 2 but disconnected.

Yet another is the Schubert divisors in the 3-fold  $GL_3/B$ , two smooth surfaces whose intersection  $\mathbb{P}^1 \cup_{pt} \mathbb{P}^1$  is not smooth.

When  $D$  is snc, define its **dual complex**  $\Delta(D) \subseteq 2^{[m]}$  to be the simplicial complex with vertex set  $[m]$ , where  $F \subseteq [m]$  to be a face iff  $\bigcap_{f \in F} D_f \neq \emptyset$ .

[Kollár '14] showed that every simplicial complex arises as the dual of some sncd – but states in [Kollár-Xu '16] a “folklore conjecture”: if  $D$  is anticanonical in  $M$ , then  $\Delta(D)$  is homeomorphic to a sphere mod a finite group.

# Bott-Samelson manifolds and their boring sncds.

Fix a pinning  $(G, B, T, W)$  of a Lie (or Kac-Moody) group. Given a word  $Q$  in the simple reflections of the Weyl group  $W$ , define the **Bott-Samelson manifold**

$$BS^Q := \left\{ (F_0, \dots, F_{\#Q}) \in (G/B)^{1+\#Q} : F_0 = B/B, \forall i (F_i, F_{i+1}) \in \overline{G_\Delta \cdot (B/B, r_{q_i} B/B)} \right\}$$

of tuples of (generalized) flags, starting at the base flag  $B/B$  and only changing a little bit at each step. This is an iterated  $\mathbb{P}^1$  bundle, hence smooth projective irreducible, and possesses a  $B$ -action, with  $(BS^Q)^T$  isolated and  $\cong 2^Q$ .

The **Bott-Samelson map**  $BS^Q \rightarrow G/B$  takes  $(F_i) \mapsto F_{\#Q}$ , with image some  $B$ -orbit closure  $X^w := \overline{BwB}/B$ . This  $w$  is the **Demazure product** of  $Q$ , the (unique) maximum product of any subword of  $Q$ . (In the boring case for us  $w = \prod Q$ , though people like that  $BS^Q \rightarrow X^w$  is then a resolution of singularities.)

Whenever  $F_{i-1} = F_i$ , we might as well skip letter  $i$  in  $Q$ , giving us an injection  $BS^{Q \setminus i} \hookrightarrow BS^Q$ . Intersecting these images we get a stratum  $\cong BS^R$  for each of the  $2^{\#Q}$  many subwords  $R \subseteq Q$ . Every intersection is nonempty!

Hence if  $D = \bigcup_{i=1}^{\#Q} BS^Q \text{ minus letter } i$ , it forms an sncd in  $BS^Q$  whose  $\Delta(D)$  is the entire simplex, rather than some interesting subcomplex of that simplex.

# Brick manifolds and spherical subword complexes.

The **brick manifold**  $\text{Brick}^Q \subseteq \text{BS}^Q$  is the  $F_{\#Q} = wB/B$  fiber of  $\text{BS}^Q \rightarrow X^w$  ( $w$  being the Demazure product). It is smooth (by Sard),  $T$ -invariant, and of dimension  $\#Q - \ell(w)$  (so, boring when  $Q$  reduced).

Let  $D = \bigcup_{q \in Q} (\text{Brick}^Q \cap \text{BS}^{Q \setminus q}) \subseteq \text{Brick}^Q$ ; it is an sncd in  $\text{Brick}^Q$ .

**Theorem [Escobar '16].**  $\Delta(D)$  is the “subword complex”  $\Delta(Q, w)$  whose facets are the complements  $Q \setminus R$  of reduced subwords  $R \subseteq Q$  with product  $w$ . It is therefore homeomorphic to a sphere [K-Miller '05].

Since  $D$  is anticanonical in  $\text{Brick}^Q$ , this is consonant with the folklore conjecture.


A **Richardson variety**  $X_u^v \subset G/B$  is the transverse intersection of a Schubert variety  $X_u := \overline{B_- u B} / B$  and an opposite Schubert variety  $X^v := \overline{B v B} / B$ .

We can resolve  $X_u = w_0 X^{w_0 u}$  using  $\text{BS}_R := w_0 \text{BS}^R$ , where  $R$  is a reduced word for  $w_0 u$ . Brion constructed a resolution of  $X_u^v$  using the fiber product of  $\text{BS}^Q \rightarrow X^v$  and  $\text{BS}_R \rightarrow X_u$ . This fiber product is naturally identified with the brick manifold  $\text{Brick}^{Q \overleftarrow{R}}$ , where  $\overleftarrow{R}$  is  $R$  reversed, and the map to  $G/B$  takes  $(F_0, F_1, \dots, F_{\#Q}, \dots, F_{\#Q + \#R}) \mapsto F_{\#Q}$ .


In the slides to come, we will give *canonical* resolutions of Richardson varieties (and thus of projected Richardsons too), not dependent on choices of  $Q$  and  $R$ .

## Example: Brion's "log resolutions" of the Richardson stratification of $GL_3/B$ .

Let  $Q = R = 121$ , reduced words in  $S_3$ , so  $Q\overleftarrow{R} = 121121$ . Then the dual complex is a 2-sphere:



The vertices are labeled with the complements of letters, the regions with reduced subwords with product  $w_0$ .  $R = 212$  gives an isomorphic complex:



# Moduli spaces of stable maps of rational curves.

Fix a 2-homology class  $\beta \in H_2(M)$  and a number  $n$  of “marked points”. We consider maps  $\gamma : \Sigma \rightarrow M$ , where  $\Sigma$  is a tree of smooth  $\mathbb{P}^1$ s with simple normal (i.e. nodal) crossings and  $n$  points (not at the nodes) marked  $1 \dots n$ . Also we require  $\gamma_*([\Sigma]) = \beta$ . (The 0 in “ $\overline{\mathcal{M}}_{0,n}$ ” below is for the only genus we consider.)

Call the map  $\gamma$  **stable** if  $\Sigma$  has only finitely many automorphisms compatible with  $\gamma$ . Specifically, each component of  $\Sigma$  collapsed by  $\gamma$  to a point should have at least three nodes + marked points.

There is a natural topology on this space  $\overline{\mathcal{M}}_{0,n}(M, \beta)$  of maps, making it compact (in limits,  $\Sigma$  can break). It is more naturally a stack than a scheme, in that one should remember the finite automorphism groups.

**Theorem [Fulton-Pandharipande '95].**  $\overline{\mathcal{M}}_{0,n}(G/P, \beta)$  is a smooth proper stack, or in other language, a compact orbifold.

This space comes with an sncd, consisting of the reducible  $\Sigma$ .

Already the case  $\overline{\mathcal{M}}_{0,n}(\text{pt}, 0)$  is interesting. Here  $D$  has one component for each of the  $2^{n-1} - n - 1$  nontrivial divisions of the marked points. The classical cross-ratio gives an isomorphism  $\overline{\mathcal{M}}_{0,4} \cong \mathbb{P}^1$ , where the sncd is the values  $0, 1, \infty$ . In particular the sncd is not anticanonical.

## (Now the new stuff!) A moduli space of equivariant maps.

We define a locally closed substack  $\overline{\mathcal{M}}'$ . Assume  $\Sigma$ 's components come in a chain  $\bigcirc \cdots \bigcirc$ , not in a knottier tree. Put a  $\mathbb{G}_m$  action on  $\Sigma$ , speed 1 on each component, with opposed weights  $+1, -1$  at the two tangent lines at each node. The two  $\mathbb{G}_m$ -fixed points in  $\Sigma$  at the ends, with respective tangent weights  $+1, -1$ , we **mark** and call  $0, \infty \in \Sigma$  (note in particular that  $n \geq 2$ ).

If a circle acts on  $M$ , together we get a  $T^2$ -action on  $\overline{\mathcal{M}}'_{0,n}(M, \beta)$ . The fixed points  $\overline{\mathcal{M}}'_{0,n}(M, \beta)^{\mathbb{G}_m}$  for the diagonal are the circle-equivariant stable maps.

**Theorem.**  $\overline{\mathcal{M}}'_{0,n \leq 3}(G/P, \beta)^{\mathbb{G}_m}$  is a smooth stack (albeit disconnected).

Fix a regular dominant weight, say  $\check{\rho}$ , acting on  $G/P$ ; by regularity  $(G/P)^{\check{\rho}} \cong W/W_P$  with Białyński-Birula decompositions the Bruhat and opposite Bruhat decompositions.

Let  $\beta = [\overline{\check{\rho} \cdot x}] \in H_2(G/P)$  where  $x \in X_u^v$  is general in the Richardson variety.

**Theorem.** Let  $\tilde{X}_u^v(m) = \left\{ \gamma \in \overline{\mathcal{M}}'_{0,m+2}(G/P, \beta)^{\mathbb{G}_m} : \gamma(0) = uP/P, \gamma(\infty) = vP/P \right\}$ .

Then  $\tilde{X}_u^v(m)$  is smooth, connected, and for  $m \leq 1$  is proper. The map  $\tilde{X}_u^v(1) \rightarrow X_u^v$  taking  $\gamma \mapsto \gamma(\text{the marked point} \neq 0, \infty)$  is a resolution of singularities.

Effectively, we're not just specifying a class in homology  $H_2(G/P)$ , but in equivariant homology  $H_2^{\mathbb{G}_m}(G/P)$ .

## Main theorems: the sncd $D \subset \widetilde{X}_u^v(0)$ .

**Theorem.** 1. Let  $\gamma : \Sigma \rightarrow X_u^v$  lie in our space  $\widetilde{X}_u^v(0)$ , and enumerate  $\Sigma$ 's fixed points  $p_0 = 0, p_1, \dots, p_c = \infty \in \Sigma^{\mathbb{G}^m}$  so that  $p_{i-1}, p_i$  lie in the same component of  $\Sigma$  for  $i = 1 \dots c$ . Then  $\gamma(p_1) < \dots < \gamma(p_{c-1})$  in the open Bruhat interval  $(u, v)$ .

2. The substack of  $\widetilde{X}_u^v(0)$  consisting of stable curves through  $w_1 < \dots < w_{c-1}$  in the open Bruhat interval  $(u, v)$  is isomorphic to  $\prod_{i=1}^c \widetilde{X}_{w_{i-1}}^{w_i}(0)$ , and in particular is smooth of codimension  $c - 1$ . (Here we take  $w_0 = u, w_c = v$ .)

3. Hence the substack  $D$  consisting of reducible stable curves is sncd, and in the  $G/B$  case, is anticanonical.


4.  $\widetilde{X}_u^v(1) \cong \widetilde{X}_u^v \times \mathbb{P}^1(0)$ . (This doesn't quite work for higher  $n$ .)

#3 prompts us to consider  $D$ 's dual complex, which is exactly the order complex of the Bruhat interval  $(u, v)$ . This simplicial complex was proven in [Björner-Wachs '82] to be homeomorphic to a sphere, using "EL-shellability". Another case confirmed of the folklore conjecture!

By #4, the dual of the sncd for  $\widetilde{X}_u^v(1)$  is almost the suspension of the Björner-Wachs sphere – first cross with an interval, triangulate, *then* cone the ends.

Note that one can define  $\widetilde{X}_u^v(n)$  using stable maps into  $X_u^v$  rather than into  $G/B$ ; we only used maps into  $G/B$  to more easily prove smoothness. The singular variety  $X_u^v$  already contains the seeds of its resolution!

Example: the dual complex  $\Delta(D)$  to the sncd  $D$  in  $\tilde{X}_{123}^{321}(1)$ .



In each component of  $D$ ,  $\Sigma$  breaks into  $\infty$ , with the marked point on one of the two components. Each corresponding vertex of  $\Delta(D)$  is labeled by  $\gamma$  (the node).

When the component with the marked point collapses, taking the node with it, we [box] its image. Otherwise the  $*$  specifies the component of the marked point. A few of the bigger faces of  $\Delta(D)$  are also labeled.

The link of the  $[u]$  (or  $[v]$ ) vertex is a copy of the Björner-Wachs sphere. Deleting those (gold) balls gives a (blue) triangulation of their sphere times an interval.

# GKM spaces and the Grassmannian case.

Call a torus action  $d$ -GKM (for Goresky-Kottwitz-MacPherson) if it fixes only finitely many subvarieties of dimension  $\leq d$  (necessarily toric). [GKM '98] only considered  $d = 1$ , which includes flag manifolds  $G/P$ . The fixed points and curves in a 1-GKM space give the vertices and edges of a graph.

It is easy to see that if  $M$  is  $d$ -GKM, then each  $\overline{M}_{0,n}(M, \beta)$  is  $(d - 1)$ -GKM. For example, the isolated fixed points in  $\tilde{X}_\mu^\nu(0)$  consist of chains of covers of  $T$ -fixed curves, each connecting some  $w_i$  to  $w_{i+1} = w_i r_\delta$ .

[Guillemin-Zara '01] observed that Grassmannians are 2-GKM, which they called “3-independence” (of isotropy weights). Hence each  $\tilde{X}_\mu^\nu(0)$  is 1-GKM.

To describe its GKM graph, we need recall the combinatorial notion of **rim-hook tableau** of shape  $\mu/\nu$ . This is a chain  $\mu = \lambda_0 \subset \lambda_1 \subset \dots \subset \lambda_m = \nu$  of partitions, where each  $\lambda_i/\lambda_{i-1}$  is a **rim-hook**, i.e. connected and containing no  $2 \times 2$  square.

**Theorem.** The  $T$ -fixed points on  $\tilde{X}_\mu^\nu(0)$  correspond to rim-hook tableaux  $\{\tau\}$ . Most of the edges out of  $\tau$  involve breaking a rim-hook into two or gluing two together, making  $\tau'$ . If rim-hooks  $i$  and  $i + 1$  of  $\tau$  together contain a  $2 \times 2$  square (so can't be glued), or share no boundary (ditto), the resulting union has a canonical alternate breaking,  $\tau'$ . These pairs  $(\tau, \tau')$  are the graph edges.



## Bonus: computing the isotropy weights on $\tilde{X}_\mu^\vee(0)$ , up to scale.

Let  $T$  act on the 1-GKM space  $M$ , and  $\rho : \mathbb{G}_m \rightarrow T$  a regular coweight ( $M^\rho = M^T$ ).

A  $T$ -fixed curve  $\delta$  in  $\overline{\mathcal{M}}_{0,n}(M, \beta)^{\mathbb{G}_m}$  is a family  $(\gamma_t)_{t \in \mathbb{P}^1}$  of  $\mathbb{G}_m$ -equivariant stable maps  $\gamma_t : \Sigma_t \rightarrow M$ , the union of whose images forms a toric  $T$ -invariant surface  $S \subseteq M$ . The images  $\gamma_t(0)$  and  $\gamma_t(\infty)$  are constant in  $t$ , and are the sink and source of the  $\mathbb{G}_m$ -action on  $S$ .

Let  $\lambda, \mu$  be the isotropy weights on  $T_{\gamma_t(0)}S$ . Then the coweight lattice of  $\text{Stab}_T(\delta)$  is  $(\lambda^\perp \cap \mu^\perp) + \mathbb{Z}\rho$ , whose perp is  $(\mathbb{Z}\lambda + \mathbb{Z}\mu) \cap \rho^\perp$ .

The isotropy weights of  $T$  on  $\gamma_0, \gamma_\infty \in \delta$  lie in  $+\mathbb{N}\lambda - \mathbb{N}\mu$  and  $-\mathbb{N}\lambda + \mathbb{N}\mu$  respectively, whose intersections with  $\rho^\perp$  are  $\cong \mathbb{N}$ . We have thus determined those isotropy weights up to scale.

In the case  $M = \text{Gr}(k, n)$ , the possible  $S$  boil down to (here  $a < b < c < d$ )

- $\text{Gr}(1, \mathbb{C}^{abc})$ , gluing two rim-hooks along a horizontal edge
- $\text{Gr}(2, \mathbb{C}^{abc})$ , gluing two rim-hooks along a vertical edge
- $\text{Gr}(1, \mathbb{C}^{ab}) \times \text{Gr}(1, \mathbb{C}^{cd})$ , swapping nonoverlapping rim-hooks
- $\text{Gr}(1, \mathbb{C}^{ac}) \times \text{Gr}(1, \mathbb{C}^{bd})$  or  $\text{Gr}(1, \mathbb{C}^{ad}) \times \text{Gr}(1, \mathbb{C}^{bc})$ , gluing then rebreaking.

I computed each isotropy weight with the recipe above, then invented  $\Phi$ , which I set up so the isotropy weight would be a multiple of  $\Phi(\tau) - \Phi(\tau')$ .

Q:  $\exists$  an equivariant ample line bundle  $\mathcal{L}$  on  $\tilde{X}_\mu^\vee(0)$  with  $\Phi(\tau) = T\text{-wt}(\mathcal{L}|_\tau)$ ?