

# The cyclic Bruhat decomposition of flag manifolds

Allen Knutson (Cornell)

Joint Meetings, Atlanta, January 2017

## Abstract

If we intersect the  $n$  cyclic translates of the Bruhat decomposition of the Grassmannian  $Gr(k, n)$ , we get the celebrated **positroid stratification** studied by Lusztig, Postnikov, Williams, Rietsch, Knutson-Lam-Speyer... It is also the stratification by projected Richardson varieties [KLS], and its most natural flag manifold version is just the Richardson varieties  $\{X_\sigma^\pi\}$ .

Nonetheless, I'll look at the intersection of the  $n$  cyclic translates of the Bruhat decomposition, and index the strata with "cyclic flag pipe dreams". Alas: unlike on  $Gr(k, n)$ , these strata can be empty, or of bad dimension.

They are determined by  $(\dim(F_k \cap \mathbb{C}^{[i,j]}))$  where  $[i, j]$  varies over cyclic intervals; unfortunately their closures are *not* given by inequalities on those dimensions, and (relatedly) this decomposition is not a stratification. Taking  $[i, j]$  only from (non-cyclic) intervals, as was useful for Schubert calculus [K], I *do* get  $\neq \emptyset$ ness, smoothness, irreducibility, and dimension.


# Arrays of dimension jumps, with pipes.

Identify  $\text{Fl}(n) := B_- \setminus \text{GL}(n)$ , using  $F_k := \text{span of top } k \text{ rows of } M \in \text{GL}(n)$ .  
 For each  $i \leq j \leq i + n$ , consider columns  $[i, j] \bmod n$  of  $M$ , and record

$J_{ij} := \{k \in [n] : \text{rank}(\text{top } k \text{ rows in cols } [i, j]) > \text{rank}(\text{top } k - 1 \text{ rows in cols } [i, j])\}$ .

e.g.  $M = \begin{pmatrix} 0 & a & b & 0 \\ 0 & a & b & * \\ * & * & * & * \\ * & * & * & * \end{pmatrix}$


$\mapsto$



**Theorem.**  $J_{ij}$  increases by one element as you go North or East,  
 i.e.  $\{(i, j) : J_{ij} \ni k\}$  is an order ideal above some  $k$ -pipe.

## Tiles for CF pipe dreams (CF = “cyclic flag”).

We can therefore think of these CF pipe dreams as being built out of tiles:



It's easy to show the elbows tile (the third type) doesn't occur with  $a > b$ .

The pipe labels on the vertical edges are weakly increasing in each row of  $\mathcal{J}$ :

$$a \leq a, \quad b \leq b, \quad a \leq b \quad \text{in the three tiles.}$$

For  $\mathcal{J}$  an  $n$ -periodic assemblage of these tiles into a **CF pipe dream**, let  $X(\mathcal{J})^\circ \subseteq \text{Fl}(n)$  denote the corresponding locally closed subset of  $\text{Fl}(n)$ .


Based on the example of  $\text{Gr}(k, n)$ , I was moved to

**Conjecture.**  $X(\mathcal{J})^\circ$  is smooth and irreducible, with codimension given by the number of **horizontal tiles**, of the left type.

(Which equals the number of vertical tiles, by the Jordan curve theorem.)

But this turns out to be false, much like most conjectures about matroid strata!

# Counterexamples: an empty stratum, and stratification failure.



The 2, 1, 2, 1 down the  $i = j$  diagonal tell us  $a = c = 0, b, d \neq 0$ .

The 124 at  $[1, 3]$  says that  $\det \begin{bmatrix} a & b & c \\ e & * & f \\ g & * & h \end{bmatrix} = \det \begin{bmatrix} 0 & b & 0 \\ e & * & f \\ g & * & h \end{bmatrix} = b(eh - fg) = 0$ .

But the 123 at  $[3, 5]$  says that  $\det \begin{bmatrix} c & d & a \\ f & * & e \\ h & * & g \end{bmatrix} = d(fg - eh) \neq 0$ . So there are none.



The same phenomenon,  $b \neq 0 \implies eh - fg = 0$ , leads to a stratum whose closure is not a union of strata.

*Moral: When an inequality helps prove an equality, watch out!*

# A flag of positroids.

The data defining  $\mathcal{J}$  tells us which positroid stratum each  $k$ -plane  $F_k$  is in, i.e. we get an  $n$ -tuple of bounded affine permutations.

In this interpretation, the  $k$ -pipe says which dots move when going from the  $k$ -ball affine permutation to the  $(k + 1)$ -ball affine permutation.



**Question.** For which  $\mathcal{J}$  is  $\overline{X(\mathcal{J})}$  determined as a set by intersecting the flag manifold with the preimages of those positroid varieties?

# Partial flag manifolds, and the loop amplituhedron.

The same recipes work on  $\text{Fl}(n_1 < n_2 < \dots < n_m = n)$ , except now we have  $n_i - n_{i-1}$  many  $n_i$ -pipes, and they mustn't cross each other.

In the case  $\text{Gr}(k, n)$ , in row  $i \in \mathbb{Z}$  the pipe labels on the  $n$  vertical edges go  $k, k, \dots, k, n, n, \dots, n$ . If we let  $\pi(i) := i + \#ks$  in that row, we get the corresponding bounded affine permutation defining the positroid stratum.

The  $\ell$ -loop amplituhedron  $\mathcal{A}_{k,\ell}$  is the  $\text{Gr}(2, 4)^\ell$ -bundle over  $\text{Gr}(k, k+4)$ .

So  $\mathcal{A}_{k,1} \cong \text{Fl}(k, k+2, k+4)$ , and  $\mathcal{A}_{k,\ell} \hookrightarrow \text{Fl}(k, k+2, k+4)^\ell$ , from which it inherits a cyclic Bruhat decomposition. Now there are  $k$  pipes labeled  $V$ , and 2 pipes labeled each of  $1, \dots, \ell$ , that can lie along one another (but not along  $V$ -pipes).


If we don't bother drawing the remaining pipes, we get pipe dreams like this:

$$M_V = \begin{bmatrix} 0 & * & 1 & 2 & 0 \end{bmatrix}$$

$$L_1 = \begin{bmatrix} * & * & 1 & 2 & 0 \\ * & * & 1 & 2 & 0 \end{bmatrix}$$

$$L_2 = \begin{bmatrix} * & * & * & 0 & * \\ * & * & * & 0 & * \end{bmatrix}$$

$\mapsto$



## Interval rank flag strata and IF pipe dreams.

If we only study intervals  $[i, j] \subseteq [n]$  of columns, rather than *cyclic* intervals, we get a coarser decomposition into **IF strata**, indexed by triangular (rather than periodic) **IF pipe dreams**. It still is finer than the Richardson stratification.

Associated to an IF pipe dream  $\mathcal{J}$  are two permutations  $\pi$  and  $\sigma$ , from the lists of pipes crossed across the North side and then down the East side.

**Theorems.** Let  $\mathcal{J}$  be an IF pipe dream, and  $\pi$  and  $\sigma$  as above.

- $X(\mathcal{J})_\circ$  is nonempty, smooth, and irreducible.
- $X(\mathcal{J})_\circ \subseteq X_{\pi^{-1}}^{\sigma^{-1}}$
- $\text{codim}(X(\mathcal{J})_\circ \subseteq X^{\sigma^{-1}}) = \#\text{vertical tiles}$ .
- $\text{codim}(X(\mathcal{J})_\circ \subseteq X_{\pi^{-1}}) = \#\text{horizontal tiles (equivalent to the previous)}$ .

Not everything is great: the same counterexample still works to show that this coarser decomposition is not a stratification.

David Speyer and I are trying to relate this decomposition to Deodhar's.

On the Grassmannian, this was the stratification I used in arXiv:1408.1261 to extend Vakil's "geometric Littlewood-Richardson rule" to equivariant K-theory. On there, though, it was *coarser* than the projected Richardson stratification.


# Towards a geometric L-R rule for IF pipe dreams.

I defined the **geometric shift**  $\mathbb{I}\mathbb{I}_\alpha$  of  $X \subseteq P \setminus G$  as  $\lim_{t \rightarrow \infty} \exp(te_\alpha) \cdot X$ , connecting a construction from Vakil with Erdős-Ko-Rado combinatorial shifting.

Here,  $\exp(te_\alpha) \cdot$  means adding  $t$  times column  $i$  to column  $j$ , and taking the limit; the rank conditions on column  $j$  thereby move backwards to column  $i$ .

Vakil gave a list of shifts to apply to (initially Richardson, eventually Schubert) varieties in  $Gr(k, n)$ . His list rasters the rows of the pipe dream bottom to top, and right to left within rows; we indicate his  $\{(i, j)\}$  below at yellow tiles.

If the corners of a yellow tile have  $NW < SE$ , the shift switches those sets (and otherwise does nothing). The resulting array of subsets may be combinatorially illegal, reflecting the geometry that  $\mathbb{I}\mathbb{I}_{i \rightarrow j} \overline{X(\mathcal{J})}$  has become reducible.



This is the calculation  $[X_{213}][X^{312}] = [X_{231}] + [X_{312}]$  (don't forget the inverting!).

The main holdup: **what does  $\mathbb{I}\mathbb{I}_{i \rightarrow j}$  do to the equations-from-inequations?**