

MAT344 Problem Set 7

(due Thursday, November 14, noon)

Notes:

- For all the questions, always *explain your reasoning* and refer to the results you are using. Just a number (even if it is the correct final answer) will **not** get you full credit.
- When submitting to Crowdmark, please ensure that your uploads are *legible, correctly rotated, and properly matched with the correct problems*. Any improperly uploaded problem scans will not be graded.
- Any assignments submitted after the deadline will not be accepted.

Part A

Three randomly chosen questions from this part will be marked.

Problem 1. How many surjections are there from $[7]$ to $[4]$? (for this question, give the answer as a number, not a formula)

Problem 2. How many derangements are there of the set $[6]$? (for this question, give the answer as a number, not a formula)

Problem 3. How many integer solutions are there to the equation

$$x_1 + x_2 + x_3 + x_4 = 32$$

with $0 \leq x_i \leq 10$ for $i = 1, 2, 3, 4$?

Problem 4. How many numbers from 1 to 5000 are divisible by either 3 or 17?

Part B

Two randomly chosen questions from this part will be marked.

Problem 5. The **Euler totient function** is a function $\phi(n)$ that is defined as follows: for a positive integer $n \geq 2$, let

$$\phi(n) = |\{m \in \mathbb{Z} | 1 \leq m \leq n, \gcd(m, n) = 1\}|$$

(that is, $\phi(n)$ is the number of positive integers less than or equal to n relatively prime to n).

- (a) Find $\phi(p)$ for p a prime number.
- (b) Find $\phi(p^k q^l)$ for p and q distinct primes and k, l positive integers.
- (c) Use inclusion-exclusion to find $\phi(n)$ for $n = p_1^{k_1} p_2^{k_2} \cdots p_m^{k_m}$ with the p_i distinct primes and k_i positive integers.

Problem 6. The principle of inclusion-exclusion is not the only approach available for counting derangements. We know that $d_1 = 0$ and $d_2 = 1$. Using this initial information, it is possible to give a recursive form for d_n . Give a combinatorial argument to prove that the number of derangements satisfies the recursive formula

$$d_n = (n - 1)(d_{n-1} + d_{n-2})$$

for $n \geq 2$.

Problem 7. A small merry-go-round has 8 seats occupied by 8 children. In how many ways can the children change places in a way that no child sits behind the same child as on the first ride? The seats do not matter, only the relative positions of the children.

Part C

This question will be marked for completion only.

Problem 8. Give an example of a problem that is difficult to enumerate directly but can be counted using inclusion-exclusion.