

REVIEW

- (1) Total mass of regions
- $\mathcal{D} \subset \mathbb{R}^2$
- and
- $\mathcal{W} \subset \mathbb{R}^3$
- :

$$M = \iint_{\mathcal{D}} \delta(x, y) dA, \quad M = \iiint_{\mathcal{W}} \delta(x, y, z) dV,$$

where δ is the mass density. The same formula can be used to compute the total charge of a region if δ is the charge density.

- (2) Moments:

$$\underline{\text{In } \mathbb{R}^2}: \quad M_x = \iint_{\mathcal{D}} y \delta(x, y) dA, \quad M_y = \iint_{\mathcal{D}} x \delta(x, y) dA$$

$$\underline{\text{In } \mathbb{R}^3}: \quad M_{yz} = \iiint_{\mathcal{W}} x \delta(x, y, z) dV, \quad M_{xz} = \iiint_{\mathcal{W}} y \delta(x, y, z) dV, \quad M_{xy} = \iiint_{\mathcal{W}} z \delta(x, y, z) dV.$$

- (3) Coordinates of the
- center of mass*
- :

$$\underline{\text{In } \mathbb{R}^2}: \quad x_{CM} = \frac{M_y}{M}, \quad y_{CM} = \frac{M_x}{M}.$$

$$\underline{\text{In } \mathbb{R}^3}: \quad x_{CM} = \frac{M_{yz}}{M}, \quad y_{CM} = \frac{M_{xz}}{M}, \quad z_{CM} = \frac{M_{xy}}{M}.$$

- (4) Random variables
- X
- and
- Y
- have joint probability density function
- $p(x, y)$
- if

$$P(a \leq X \leq b; c \leq Y \leq d) = \int_{x=a}^b \int_{y=c}^d p(x, y) dy dx.$$

- (5) A joint probability density function must satisfy
- $p(x, y) \geq 0$
- and

$$\int_{x=-\infty}^{\infty} \int_{y=-\infty}^{\infty} p(x, y) dy dx = 1.$$

- (6) Let
- $G(u, v) = (x(u, v), y(u, v))$
- be a mapping. The
- Jacobian*
- of
- G
- is the determinant

$$\text{Jac}(G) = \frac{\partial(u, v)}{\partial(x, y)} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix}.$$

- (7)
- $\text{Jac}(G) = [\text{Jac}(F)]^{-1}$
- , where
- $F = G^{-1}$
- .

- (8) Change of Variables Formula: If
- $G: \mathcal{D}_0 \rightarrow \mathcal{D}$
- has component functions with continuous partial derivatives and one-to-one on the interior of
- \mathcal{D}_0
- , and if
- f
- is continuous, then

$$\iint_{\mathcal{D}} f(x, y) dx dy = \iint_{\mathcal{D}_0} f(x(u, v), y(u, v)) \frac{\partial(u, v)}{\partial(x, y)} du dv.$$

PROBLEMS

- (1) Compute the total mass of the plate in Figure 1 assuming a mass density of $f(x, y) = \frac{x^2}{(x^2+y^2)}$ g/cm².
- (2) Find the z_{CM} (z coordinate of the center of mass) of a cylinder of radius 2 and height 4 and mass density e^{-z} , where z is the height above the base (an expression in terms of e is fine). Can you find the x_{CM} and y_{CM} without explicitly computing any integral?
- (3) Find a constant C such that

$$p(x, y) = \begin{cases} Cxy & \text{if } 0 \leq x \text{ and } 0 \leq y \leq 1 - x \\ 0 & \text{otherwise} \end{cases}$$

is a joint probability density function, then calculate $P(X \geq Y)$.

- (4) Compute $\iint_{\mathcal{D}} (x + 3y) dx dy$, where \mathcal{D} is the shaded region in Figure 3. *Hint:* Use the map $\Phi(u, v) = (u - 2v, v)$.
- (5) Find a mapping Φ that maps the disk $u^2 + v^2 \leq 1$ onto the interior of the ellipse $(\frac{x}{a})^2 + (\frac{y}{b})^2 \leq 1$. Then use the Change of Variables Formula to prove that the area of the ellipse is πab .




Figure 1: Problem 1




Figure 2: Problem 2




Figure 3: Problem 4