

On the optimal error exponents for classical and quantum antidistinguishability

Hemant K. Mishra,^{1,*} Michael Nussbaum,^{2,†} and Mark M. Wilde^{1,‡}

¹*School of Electrical and Computer Engineering,
Cornell University, Ithaca, New York 14850, USA*

²*Department of Mathematics, Cornell University, Ithaca, New York 14850, USA*

The concept of antidistinguishability of quantum states has been studied to investigate foundational questions in quantum mechanics. It is also called quantum state elimination, because the goal of such a protocol is to guess which state, among finitely many chosen at random, the system is not prepared in (that is, it can be thought of as the first step in a process of elimination). Antidistinguishability has been used to investigate the reality of quantum states, ruling out ψ -epistemic ontological models of quantum mechanics [Pusey *et al.*, *Nat. Phys.*, 8(6):475-478, 2012]. Thus, due to the established importance of antidistinguishability in quantum mechanics, exploring it further is warranted.

In this paper, we provide a comprehensive study of the optimal error exponent—the rate at which the optimal error probability vanishes to zero asymptotically—for classical and quantum antidistinguishability. We derive an exact expression for the optimal error exponent in the classical case and show that it is given by the *multivariate classical Chernoff divergence*. Our work thus provides this divergence with a meaningful operational interpretation as the optimal error exponent for antidistinguishing a set of probability measures. For the quantum case, we provide several bounds on the optimal error exponent: a lower bound given by the best pairwise Chernoff divergence of the states, an upper bound in terms of *max-relative entropy*, and lower and upper bounds in terms of minimal and maximal *multivariate quantum Chernoff divergences*. It remains an open problem to obtain an explicit expression for the optimal error exponent for quantum antidistinguishability.

Dedicated to the memory of Mary Beth Ruskai. She was an important foundational figure in the field of quantum information, and her numerous seminal research contributions and reviews, including [LR73, Rus02, HSR03], have inspired many quantum information scientists.

CONTENTS

| | |
|--|----|
| I. Introduction | 2 |
| A. Contributions | 3 |
| B. Literature review | 4 |
| C. Paper organization | 4 |
| II. Mathematical background | 5 |
| A. Antidistinguishability of probability measures | 5 |
| B. Multivariate classical Chernoff divergence | 7 |
| C. Quantum states, channels, and measurements | 8 |
| D. Antidistinguishability of quantum states | 9 |
| E. Quantum Chernoff divergence | 10 |
| III. Multivariate Chernoff divergence as the optimal error exponent for classical antidistinguishability | 10 |
| A. Multivariate Chernoff divergence versus pairwise Chernoff divergences | 16 |

* hemant.mishra@cornell.edu

† nussbaum@math.cornell.edu

‡ wilde@cornell.edu

| | |
|---|----|
| IV. Achievable error exponent for quantum antidistinguishability | 18 |
| A. One-shot case | 18 |
| B. Asymptotic case | 21 |
| V. Bounds on the optimal error exponent for quantum antidistinguishability from multivariate quantum Chernoff divergences | 21 |
| A. Multivariate quantum Chernoff divergences | 21 |
| B. Bounds on the optimal error exponent for quantum antidistinguishability | 24 |
| VI. Max-relative entropy bounds on the error probability and optimal error exponent | 25 |
| VII. Conclusion | 28 |
| Acknowledgments | 28 |
| Declaration of interests | 28 |
| Data availability statement | 29 |
| References | 29 |
| A. Expectation values at non-corner points | 31 |
| B. Proof of Theorem 7 | 34 |
| C. Proof of Equation (139) | 37 |
| D. Proof of Proposition 15 | 38 |
| E. Proof of Proposition 16 | 39 |
| F. Proof of Proposition 17 | 40 |
| G. Additivity of the optimal error exponent | 41 |
| H. Limit of the regularized maximal multivariate quantum Chernoff divergence | 42 |
| I. Proof of Equation (208) | 42 |

I. INTRODUCTION

Quantum state discrimination is a fundamental component of quantum information science, which plays a key role in quantum computing [BCDvD06], quantum communication [Hel69, BC09], and quantum key distribution [BK15]. The state discrimination or distinguishability task is to infer the actual state of a quantum system by applying a quantum measurement to the system. More formally, consider a quantum system prepared in one of the quantum states ρ_1, \dots, ρ_r . A quantum measurement is specified by a positive operator-valued measure $\{M_1, \dots, M_r\}$ with output i indicating ρ_i as the true state of the system with success probability $\text{Tr}[M_i \rho_i]$, as given by the Born rule [Bor26].

The task that we consider here is in a sense opposite to the aforementioned task of distinguishability, and it is thus called *antidistinguishability* of quantum states or *quantum state elimination*

[CFS02, PBR12, BCLM14, Lei14, BJOP14, HB20, LD20, RS23]. In particular, for the task of antidistinguishability, we are interested in designing a measurement whose outcome corresponds to a state that is not the actual state of the quantum system. In the classical version of the antidistinguishability problem, quantum states are replaced by probability measures on a measurable space, and the task is to rule out one of the probability measures upon observing i.i.d. (independent and identically distributed) data that is not produced by the probability measure.

As an illustrative example in the classical case, suppose that one of three possible dice is tossed, a red one with probability distribution p_R , a green one with probability distribution p_G , or a blue one with probability distribution p_B . The task is then, after observing a sample, to output “not red” if the green or blue die is tossed, “not green” if the red or blue die is tossed, and “not blue” if the red or green die is tossed. It is also of interest to consider the antidistinguishability task when the same colored die is tossed multiple times, leading to several samples that one can use to arrive at a conclusion.

To the best of our knowledge, an analysis of the asymptotics of the error probability of antidistinguishability is missing in the literature for both cases, classical as well as quantum, and it is this scenario that we consider in our paper.

A. Contributions

In this paper, we provide a comprehensive study of the optimal error exponent—the rate at which the optimal error probability vanishes to zero asymptotically—for classical and quantum antidistinguishability.

- We derive an exact expression for the optimal error exponent in the classical case and show that it is given by the *multivariate classical Chernoff divergence* (Theorem 7). Our work thus provides this multivariate divergence with a meaningful operational interpretation as the optimal error exponent for antidistinguishing a set of probability measures.
- We provide several bounds on the optimal error exponent in the quantum case:
 - lower bound given by the best pairwise Chernoff divergence of the states (Theorem 12),
 - upper bound in terms of *max-relative entropy* (Theorem 20), and
 - lower and upper bounds in terms of minimal and maximal *multivariate quantum Chernoff divergences* (Theorem 18).
- Some other interesting consequences of our analysis are as follows:
 - We provide an upper bound on the optimal error probability of antidistinguishing an ensemble of quantum states in terms of the pairwise optimal error probabilities of the states, and consequently, we deduce that the given quantum states are perfectly antidistinguishable if at least two of them are orthogonal to each other (Theorem 9);
 - We derive a lower bound on the optimal quantum antidistinguishability error probability in terms of an exponentiated and optimized max-relative entropy (Proposition 19).

It remains an open problem to determine an explicit expression for the optimal error exponent in the quantum case.

B. Literature review

Let us briefly review some prior contributions to the topic of antidistinguishability. We note here that quantum state discrimination is equivalent to finding a size- $(r - 1)$ subset of $\{\rho_1, \dots, \rho_r\}$ such that none of the states in the subset is the true state of the system; thus, the task is equivalent to what is called *quantum $(r - 1)$ -state exclusion*. A generalization of this task is *quantum m -state exclusion* for $1 \leq m \leq r - 1$, which aims at detecting a size- m subset of $\{\rho_1, \dots, \rho_r\}$ such that none of the states in the subset is the true state of the system [RS23]. Quantum 1-state exclusion is therefore the same as antidistinguishability of quantum states.

The concept of antidistinguishability has been studied to investigate foundational questions in quantum mechanics [CFS02, PBR12, BCLM14, Lei14]. For example, it was used in [PBR12] to investigate the reality of quantum states, ruling out ψ -epistemic ontological models of quantum mechanics. It was also used in studying quantum communication complexity [HB20] and in deriving noncontextuality inequalities [LD20]. Thus, due to the established importance of antidistinguishability in quantum mechanics, exploring it further is warranted. There have been a number of works that determine algebraic conditions on a set of quantum states such that perfect antidistinguishability is possible. A sufficient condition for perfect antidistinguishability of pure states [HK18] is that if some positive linear combination of the pure states is a projection with a “special” kernel, then the states are *antidistinguishable*. In the same paper, a necessary and sufficient condition for antidistinguishability of pure states was given, which demands the existence of projections satisfying three non-trivial conditions. Very recently, a necessary and sufficient condition for non-antidistinguishability of general quantum states was given in [RS23], which also demands the existence of a Hermitian matrix with positive trace satisfying a set of non-trivial inequalities. Even though the conditions given in the aforementioned works are interesting and insightful, verifying them is not straightforward. One of the consequences of our work is that we provide a simple sufficient condition for perfect antidistinguishability of quantum states (Theorem 9).

C. Paper organization

The organization of our paper is as follows. In Section II, we state some definitions and provide a brief mathematical background of relevant topics covered in our paper. We start Section III by building a theory of classical antidistinguishability, where we introduce the notions of optimal error probability and optimal error exponent. We then derive an explicit expression for the optimal error exponent in the classical case, and we show that it is given by the *multivariate classical Chernoff divergence* (Theorem 7). The following sections deal with the optimal error exponent in the quantum case. We begin Section IV by providing an upper bound on the optimal error probability of antidistinguishing an ensemble of quantum states in terms of the pairwise optimal error probabilities of the states (Theorem 9), and we then use this result to derive a lower bound on the optimal error exponent (Theorem 12). Next, we provide both lower and upper bounds on the optimal error exponent in terms of minimal and maximal *multivariate quantum Chernoff divergences* in Section V B (Theorem 18). Lastly, in Section VI, we derive a lower bound on the optimal error probability in terms of an exponentiated and optimized *max-relative entropy* of the states (Proposition 19), and we then use this result to provide an upper bound on the optimal error exponent in terms of the max-relative entropy (Theorem 20). Appendices B through I contain mathematical proofs of various claims made throughout the paper.

II. MATHEMATICAL BACKGROUND

A. Antidistinguishability of probability measures

Let P_1, \dots, P_r be probability measures on a measurable space (Ω, \mathcal{A}) , where \mathcal{A} is a σ -algebra on the set Ω . Let η_1, \dots, η_r be positive real numbers such that $\sum_{i \in [r]} \eta_i = 1$. Set $[r] := \{1, \dots, r\}$. Throughout the paper, we call

$$\mathcal{E}_{\text{cl}} := \{(\eta_i, P_i) : i \in [r]\} \quad (1)$$

an ensemble of probability measures on the measurable space (Ω, \mathcal{A}) . Let μ be the dominating measure

$$\mu := \sum_{i \in [r]} \eta_i P_i, \quad (2)$$

and p_1, \dots, p_r the induced densities

$$p_i := \frac{dP_i}{d\mu}, \quad i \in [r], \quad (3)$$

which are given by the Radon–Nikodým theorem [Bil95].

The problem of distinguishability, i.e., identifying the correct probability density p_i based on i.i.d. (independent and identically distributed) data, has been well studied. This problem is as follows: Suppose that i is sampled with probability η_i , and then n i.i.d. samples are selected according to the product measure $P_i^{\otimes n}$. The task is to identify the correct value of i based on the n i.i.d. samples observed. It is known that the maximum likelihood method is optimal, and the optimal asymptotic error exponent is equal to the Chernoff divergence for the least favorable pair (p_i, p_j) , for $i \neq j$ [Sal73, Tor81, LJ97, Sal99, Sal03]. The total success probability for the identification task, in the case that $n = 1$, is given by

$$\int d\mu (\eta_1 p_1 \vee \dots \vee \eta_r p_r) := \int d\mu(\omega) \max\{\eta_1 p_1(\omega), \dots, \eta_r p_r(\omega)\}. \quad (4)$$

For the antidistinguishability problem in the classical case, the task is to guess a probability density that is not represented by the observed data. For this problem, no literature is available to the best of our knowledge. A reasonable first idea for selecting a density that is unlikely to be the true one is to choose the one such that $\eta_i p_i(\omega)$ is minimum if ω is observed. This corresponds to a *minimum likelihood principle*. In what follows, we discuss this idea more formally.

A deterministic decision rule for the antidistinguishability problem is a function

$$\delta : \Omega \rightarrow \{\mathbf{e}_i : i \in [r]\}, \quad (5)$$

where \mathbf{e}_i is the i th standard unit vector in \mathbb{R}^r , such that $\delta(\omega) = \mathbf{e}_i$ means that we indicate p_i to be our guess for the density that is not the true one. More generally, we can admit a randomized decision rule, along the following lines:

$$\delta : \Omega \rightarrow [0, 1]^r, \quad \sum_{i \in [r]} \delta_i(\omega) = 1. \quad (6)$$

If p_i is the true density, then the antidistinguishability error probability is given by

$$\int d\mu(\omega) \delta_i(\omega) p_i(\omega), \quad (7)$$

and the total error probability is

$$\text{Err}_{\text{cl}}(\delta; \mathcal{E}_{\text{cl}}) := \sum_{i \in [r]} \eta_i \int \mathbf{d}\mu(\omega) \delta_i(\omega) p_i(\omega) = \int \mathbf{d}\mu(\omega) \sum_{i \in [r]} \delta_i(\omega) \eta_i p_i(\omega). \quad (8)$$

To minimize the above expression, we can minimize the integrand for every ω . Since $\delta_i(\omega)$ is a weight, we should place maximum weight on the smallest of $\eta_i p_i(\omega)$. So, the optimal decision for given ω corresponds to the *minimum likelihood rule*: $\delta^*(\omega) = \mathbf{e}_i$, if $i \in [r]$ is the minimum index such that $\eta_i p_i(\omega) = \min\{\eta_1 p_1(\omega), \dots, \eta_r p_r(\omega)\}$. The total error probability when using the decision rule δ^* is the optimal error probability, given by

$$\text{Err}_{\text{cl}}(\mathcal{E}_{\text{cl}}) := \text{Err}_{\text{cl}}(\delta^*; \mathcal{E}_{\text{cl}}) = \int \mathbf{d}\mu(\omega) \min\{\eta_1 p_1(\omega), \dots, \eta_r p_r(\omega)\} = \int \mathbf{d}\mu (\eta_1 p_1 \wedge \dots \wedge \eta_r p_r). \quad (9)$$

In the asymptotic treatment of the problem, we consider the n -fold ensemble $\mathcal{E}_{\text{cl}}^n := \{(\eta_i, P_i^{\otimes n}) : i \in [r]\}$ on the n -fold measurable space $(\Omega^n, \mathcal{A}^{(n)})$, where Ω^n is the n -fold Cartesian product of Ω and $\mathcal{A}^{(n)}$ is the σ -algebra on Ω^n generated by the n -fold Cartesian product of \mathcal{A} . It then follows that the optimal antidistinguishability error probability, in this case, is

$$\text{Err}_{\text{cl}}(\mathcal{E}_{\text{cl}}^n) = \int \mathbf{d}\mu^{\otimes n} (\eta_1 p_1^{\otimes n} \wedge \dots \wedge \eta_r p_r^{\otimes n}). \quad (10)$$

Set $\eta_{\min} := \min\{\eta_1, \dots, \eta_r\}$ and $\eta_{\max} := \max\{\eta_1, \dots, \eta_r\}$. It is easy to see that for all $n \in \mathbb{N}$, we have

$$\eta_{\min} \int \mathbf{d}\mu^{\otimes n} (p_1^{\otimes n} \wedge \dots \wedge p_r^{\otimes n}) \leq \text{Err}_{\text{cl}}(\mathcal{E}_{\text{cl}}^n) \leq \eta_{\max} \int \mathbf{d}\mu^{\otimes n} (p_1^{\otimes n} \wedge \dots \wedge p_r^{\otimes n}). \quad (11)$$

This implies

$$\liminf_{n \rightarrow \infty} -\frac{1}{n} \ln \text{Err}_{\text{cl}}(\mathcal{E}_{\text{cl}}^n) = \liminf_{n \rightarrow \infty} -\frac{1}{n} \ln \int \mathbf{d}\mu^{\otimes n} (p_1^{\otimes n} \wedge \dots \wedge p_r^{\otimes n}), \quad (12)$$

which is independent of η_1, \dots, η_r .

Definition 1. The optimal error exponent for antidistinguishing the probability measures of a given ensemble $\mathcal{E}_{\text{cl}} = \{(\eta_i, P_i) : i \in [r]\}$ is defined by

$$E_{\text{cl}}(P_1, \dots, P_r) := \liminf_{n \rightarrow \infty} -\frac{1}{n} \ln \text{Err}_{\text{cl}}(\mathcal{E}_{\text{cl}}^n). \quad (13)$$

Remark 1. We note that the above definition of the optimal error exponent is independent of the choice of the dominating measure μ . This is because the development in (4)–(12) is independent of the choice of the probability measure μ dominating P_1, \dots, P_r . Indeed, if μ' is an arbitrary probability measure dominating P_1, \dots, P_r , then μ' also dominates μ . Let $\nu := \frac{\mathbf{d}\mu}{\mathbf{d}\mu'}$. We then have

$$p'_i := \frac{\mathbf{d}P_i}{\mathbf{d}\mu'} = \frac{\mathbf{d}P_i}{\mathbf{d}\mu} \cdot \frac{\mathbf{d}\mu}{\mathbf{d}\mu'} = p_i \nu, \quad \text{for all } i \in [r]. \quad (14)$$

Consequently, the quantity in (4) is given by

$$\int \mathbf{d}\mu (\eta_1 p_1 \vee \dots \vee \eta_r p_r) = \int \mathbf{d}\mu' (\eta_1 p_1 \vee \dots \vee \eta_r p_r) \nu \quad (15)$$

$$= \int \mathbf{d}\mu' (\eta_1 p_{1\nu} \vee \cdots \vee \eta_r p_{r\nu}) \quad (16)$$

$$= \int \mathbf{d}\mu' (\eta_1 p'_1 \vee \cdots \vee \eta_r p'_r). \quad (17)$$

Similarly, the remaining quantities in (4)–(12) can be shown to be independent of the choice of μ . See [Shi16, p. 233].

B. Multivariate classical Chernoff divergence

In 1909, Hellinger introduced a function of multiple probability distributions [Hel09]. Known as the *Hellinger transform* in the literature on probability and statistics, this quantity plays an important role in our work. We recall its definition below. Let $\mathcal{E}_{\text{cl}} = \{(\eta_i, P_i) : i \in [r]\}$ be an ensemble of probability measures on a measurable space (Ω, \mathcal{A}) . Let μ be the dominating measure defined by (2), and let p_1, \dots, p_r be the induced probability densities given by (3). Denote by \mathbb{S}_r the unit simplex in \mathbb{R}^r :

$$\mathbb{S}_r := \left\{ \mathbf{s} \in [0, 1]^r : \mathbf{s} = (s_1, \dots, s_r), \sum_{i \in [r]} s_i = 1 \right\}. \quad (18)$$

Definition 2. *The Hellinger transform of the probability measures P_1, \dots, P_r is a function on the unit simplex, defined as*

$$\mathbb{H}_{\mathbf{s}}(P_1, \dots, P_r) := \int \mathbf{d}\mu p_1^{s_1} \cdots p_r^{s_r}, \quad \text{for } \mathbf{s} := (s_1, \dots, s_r) \in \mathbb{S}_r. \quad (19)$$

Here we use the convention $0^0 = 1$ of [Str11, Definition 5.10].

The term ‘‘Hellinger transform’’ was perhaps first used in [LeC70], followed by several works in the area of probability and statistics. See [Tor81, Jac89, Tor91, Gri93, FL96, LM10, Str11] and references therein.

Remark 2. We emphasize that the Hellinger transform given in Definition 2, and hence our further analysis of the classical antidistinguishability error probability, is independent of the choice of the dominating measure μ . This easily follows by similar arguments as in Remark 1. See [Shi16, Chapter 3, Section 9, Lemma 3] for a proof in the case of $r = 2$.

If P_1, \dots, P_r are mutually absolutely continuous, i.e., for all $A \subseteq \Omega$, $P_1(A) = 0$ if and only if $P_i(A) = 0$ for all $i \in [r]$, then the Hellinger transform is continuous on \mathbb{S}_r . Indeed, we have $P_i \leq \eta_i^{-1} \mu$, implying that $p_i \leq \eta_i^{-1}$ for all $i \in [r]$. This gives $\prod_{i \in [r]} (p_i + 1)$ as an integrable upper bound on $\prod_{i \in [r]} p_i^{s_i}$ for all $(s_1, \dots, s_r) \in \mathbb{S}_r$. If the probability measures are mutually absolutely continuous, then

$$\mu(\{\omega \in \Omega : p_i(\omega) > 0 \forall i \in [r]\}) = 1. \quad (20)$$

Without loss of generality, we can then assume that $p_i(\omega) > 0$ for all $\omega \in \Omega$ and $i \in [r]$. Thus, for every $\mathbf{s} := (s_1, \dots, s_r) \in \mathbb{S}_r$ and for all $n \in \mathbb{N}$ and every sequence $\mathbf{s}^{(n)} := (s_1^{(n)}, \dots, s_r^{(n)})$ in the interior of \mathbb{S}_r such that $\lim_{n \rightarrow \infty} \mathbf{s}^{(n)} = \mathbf{s}$, we have

$$\lim_{n \rightarrow \infty} \prod_{i \in [r]} (p_i(\omega))^{s_i^{(n)}} = \prod_{i \in [r]} (p_i(\omega))^{s_i}, \quad \text{for } \omega \in \Omega. \quad (21)$$

By the Lebesgue dominated convergence theorem, we then have $\lim_{n \rightarrow \infty} \mathbb{H}_s^{(n)}(P_1, \dots, P_r) = \mathbb{H}_s(P_1, \dots, P_r)$, thereby proving continuity of the Hellinger transform on \mathbb{S}_r .

In general, the Hellinger transform is a measure of “closeness” among several probability distributions. It is easy to see that

$$0 \leq \mathbb{H}_s(P_1, \dots, P_r) \leq 1, \quad (22)$$

which follows from Hölder’s inequality [Str11, Lemma 53.3]. As the value of $\mathbb{H}_s(P_1, \dots, P_r)$ gets close to 0, the distance among the measures increases in some sense [FL96].

The following quantity plays an important role in our paper.

Definition 3. We define the multivariate Chernoff divergence of the probability measures P_1, \dots, P_r by

$$\xi_{\text{cl}}(P_1, \dots, P_r) := -\ln \inf_{s \in \mathbb{S}_r} \mathbb{H}_s(P_1, \dots, P_r), \quad (23)$$

where \mathbb{H}_s is defined in (19).

The divergence can be viewed as a generalization of the classical Chernoff divergence, the latter being a special case of the former for $r = 2$ [Che52]. One of the main results of our paper is that the optimal error exponent for antidistinguishing an ensemble of probability measures is equal to their multivariate Chernoff divergence (Theorem 7).

C. Quantum states, channels, and measurements

A quantum system is associated with a complex Hilbert space \mathcal{H} . We focus exclusively on systems with finite-dimensional Hilbert spaces in this paper. Let $\dim(\mathcal{H})$ denote the dimension of \mathcal{H} . We denote every element of \mathcal{H} using the *ket* notation as $|\psi\rangle, |\phi\rangle$, etc., and every element of its dual using the *bra* notation as $\langle\psi|, \langle\phi|$, etc. The notations go well with the natural action of a dual element $\langle\psi|$ on a vector $|\phi\rangle$ in terms of the inner product of the two vectors: $\langle\psi|(|\phi\rangle) = \langle\psi|\phi\rangle$.

A quantum state of a system is identified by a density operator ρ , which is a self-adjoint, positive semidefinite operator of unit trace acting on \mathcal{H} . A pure state is given by a state vector $|\psi\rangle \in \mathcal{H}$ whose corresponding density operator is $|\psi\rangle\langle\psi|$. The set of density operators forms a convex set with pure states as the extreme points. Let $\mathcal{D}(\mathcal{H})$ denote the set of density operators and $\mathcal{L}(\mathcal{H})$ the space of linear operators acting on \mathcal{H} . We shall use the notation \mathcal{D} for the set of density operators whenever the underlying Hilbert space is clear from the context. A quantum channel \mathcal{N} , between two quantum systems represented by Hilbert spaces \mathcal{H} and \mathcal{K} , is a completely positive, trace-preserving linear map from $\mathcal{L}(\mathcal{H})$ to $\mathcal{L}(\mathcal{K})$. In particular, for all $\rho \in \mathcal{D}(\mathcal{H})$, we have that $\mathcal{N}(\rho) \in \mathcal{D}(\mathcal{K})$.

A quantum measurement is described by a positive operator-valued measure (POVM) $\mathcal{M} = \{M_1, \dots, M_r\}$, which is a finite set of positive semi-definite operators whose sum is the identity operator, i.e.,

$$M_i \geq 0 \text{ for all } i \in [r], \quad \sum_{i \in [r]} M_i = \mathbb{I}, \quad (24)$$

where \mathbb{I} is the identity operator acting on \mathcal{H} .

The projection onto the support of an operator A is denoted by $\text{Supp}(A)$, its absolute value is denoted by $|A| := \sqrt{A^\dagger A}$, and its positive part by $A_+ := \frac{1}{2}(A + |A|)$. For two Hermitian operators A

and B , we use the notation

$$A \wedge B := \frac{1}{2}(A + B - |A - B|), \quad (25)$$

in analogy with $\min(a, b) = \frac{1}{2}(a + b - |a - b|) \equiv a \wedge b$ for $a, b \in \mathbb{R}$.

D. Antidistinguishability of quantum states

Suppose that a quantum system is prepared in one of the quantum states ρ_1, \dots, ρ_r with *a priori* probability distribution η_1, \dots, η_r such that $\eta_i > 0$ for all $i \in [r]$. Throughout the paper, we call $\{(\eta_i, \rho_i) : i \in [r]\}$ an ensemble of quantum states over a Hilbert space \mathcal{H} and denote it by \mathcal{E} . Antidistinguishability of the states performed by a POVM $\mathcal{M} = \{M_1, \dots, M_r\}$ can be described as follows: “the measurement outcome i occurring corresponds to a guess that the true state of the system is not ρ_i .” Thus, if ρ_i is the true state of the system, then $\text{Tr}[M_i \rho_i]$ is the error probability for making an incorrect guess. The average error probability of antidistinguishability, for a fixed POVM \mathcal{M} , is then given by

$$\text{Err}(\mathcal{M}; \mathcal{E}) := \sum_{i \in [r]} \eta_i \text{Tr}[M_i \rho_i]. \quad (26)$$

We are interested in determining the optimal antidistinguishability error probability, which is optimized over all possible measurements:

$$\text{Err}(\mathcal{E}) := \inf_{\mathcal{M}} \text{Err}(\mathcal{M}; \mathcal{E}), \quad (27)$$

where the infimum is taken over all POVMs of the form $\mathcal{M} = \{M_1, \dots, M_r\}$ acting on \mathcal{H} .

The quantum states are said to be perfectly antidistinguishable if there exists a quantum measurement whose outputs always correspond to a false state of the system; i.e., there exists a POVM \mathcal{M} such that $\text{Err}(\mathcal{M}; \mathcal{E}) = 0$. In general, an ensemble of quantum states may not be antidistinguishable, which means, for such an ensemble \mathcal{E} , that $\text{Err}(\mathcal{M}; \mathcal{E}) > 0$ for every POVM \mathcal{M} . For instance, two non-orthogonal quantum states are not perfectly antidistinguishable [Lei14].

In the asymptotic treatment of the antidistinguishability problem for a given ensemble $\mathcal{E} = \{(\eta_i, \rho_i) : i \in [r]\}$, we consider the n -fold ensemble $\mathcal{E}^n := \{(\eta_i, \rho_i^{\otimes n}) : i \in [r]\}$. The optimal error probability of antidistinguishability for \mathcal{E}^n is by definition given as

$$\text{Err}(\mathcal{E}^n) = \inf_{\mathcal{M}^{(n)}} \sum_{i \in [r]} \text{Tr} \left[\eta_i M_i^{(n)} \rho_i^{\otimes n} \right], \quad (28)$$

where the infimum is taken over the set of POVMs $\mathcal{M}^{(n)} = \{M_1^{(n)}, \dots, M_r^{(n)}\}$ acting on the n -fold tensor product Hilbert space $\mathcal{H}^{\otimes n}$. This gives for all $n \in \mathbb{N}$,

$$\eta_{\min} \inf_{\mathcal{M}^{(n)}} \sum_{i \in [r]} \text{Tr} \left[M_i^{(n)} \rho_i^{\otimes n} \right] \leq \text{Err}(\mathcal{E}^n) \leq \eta_{\max} \inf_{\mathcal{M}^{(n)}} \sum_{i \in [r]} \text{Tr} \left[M_i^{(n)} \rho_i^{\otimes n} \right], \quad (29)$$

which implies that

$$\liminf_{n \rightarrow \infty} -\frac{1}{n} \ln \text{Err}(\mathcal{E}^n) = \liminf_{n \rightarrow \infty} -\frac{1}{n} \ln \inf_{\mathcal{M}^{(n)}} \sum_{i \in [r]} \text{Tr} \left[M_i^{(n)} \rho_i^{\otimes n} \right], \quad (30)$$

the latter being independent of η_1, \dots, η_r .

Definition 4. The optimal error exponent for antidistinguishing the states of a quantum ensemble $\mathcal{E} = \{(\eta_i, \rho_i) : i \in [r]\}$ is defined by

$$E(\rho_1, \dots, \rho_r) := \liminf_{n \rightarrow \infty} -\frac{1}{n} \ln \text{Err}(\mathcal{E}^n). \quad (31)$$

E. Quantum Chernoff divergence

Here we briefly recall known results for distinguishability of two or more states; a key quantity for this purpose is as follows:

Definition 5. The quantum Chernoff divergence between two states ρ_1 and ρ_2 is defined as

$$\xi(\rho_1, \rho_2) := -\ln \inf_{s \in [0,1]} \text{Tr}[\rho_1^s \rho_2^{1-s}]. \quad (32)$$

If $\rho_1 = |\psi\rangle\langle\psi|$ and $\rho_2 = |\phi\rangle\langle\phi|$ are pure states, then

$$\xi(\rho_1, \rho_2) = -\ln |\langle\psi|\phi\rangle|^2. \quad (33)$$

The quantum Chernoff divergence between two states is known to be the *optimal error exponent* in distinguishing them [ACMT⁺07, NS09]; i.e.,

$$\lim_{n \rightarrow \infty} -\frac{1}{n} \ln (\text{Tr}[\eta_1 \rho_1^{\otimes n} \wedge \eta_2 \rho_2^{\otimes n}]) = \lim_{n \rightarrow \infty} -\frac{1}{n} \ln \left(\frac{1}{2} [1 - \|\eta_1 \rho_1^{\otimes n} - \eta_2 \rho_2^{\otimes n}\|_1] \right) = \xi(\rho_1, \rho_2), \quad (34)$$

where we have used the well known fact that the optimal error probability in distinguishing $\rho_1^{\otimes n}$ from $\rho_2^{\otimes n}$ is equal to

$$\text{Tr}[\eta_1 \rho_1^{\otimes n} \wedge \eta_2 \rho_2^{\otimes n}] = \frac{1}{2} [1 - \|\eta_1 \rho_1^{\otimes n} - \eta_2 \rho_2^{\otimes n}\|_1], \quad (35)$$

with $\rho_1^{\otimes n}$ prepared with probability η_1 and $\rho_2^{\otimes n}$ with probability η_2 [Hel69, Hol72].

It is known more generally that the optimal error exponent in distinguishing the ensemble $\{(\eta_i, \rho_i^{\otimes n}) : i \in [r]\}$ is equal to the minimum pairwise Chernoff divergence [Li16].

III. MULTIVARIATE CHERNOFF DIVERGENCE AS THE OPTIMAL ERROR EXPONENT FOR CLASSICAL ANTIDISTINGUISHABILITY

In this section, we prove that the optimal error exponent for antidistinguishing an ensemble of probability measures (13) is equal to the multivariate Chernoff divergence of the probability measures. We first prove this result for mutually absolutely continuous probability measures because the proof in this case is more readable. The proof in the general case is similar, albeit more technical, and so we present it separately in Appendix B. Some aspects of the proof presented here follow the development in the appendix of [NS09].

Theorem 6. Consider an ensemble $\mathcal{E}_{\text{cl}} = \{(\eta_i, P_i) : i \in [r]\}$ of mutually absolutely continuous probability measures on a measurable space (Ω, \mathcal{A}) . The optimal error exponent for antidistinguishing-

ing the probability measures is given by their multivariate Chernoff divergence, i.e.,

$$E_{\text{cl}}(P_1, \dots, P_r) = \lim_{n \rightarrow \infty} -\frac{1}{n} \ln \text{Err}_{\text{cl}}(\mathcal{E}_{\text{cl}}^n) = \xi_{\text{cl}}(P_1, \dots, P_r), \quad (36)$$

where, recalling (18), (19), and (23), the multivariate classical Chernoff divergence ξ_{cl} is defined as

$$\xi_{\text{cl}}(P_1, \dots, P_r) := -\ln \inf_{\mathbf{s} \in \mathbb{S}_r} \int \mathbf{d}\mu p_1^{s_1} \cdots p_r^{s_r}. \quad (37)$$

Proof. Let μ be the dominating measure given by (2) and p_1, \dots, p_r the induced densities defined in (3). The assumption that the probability measures are mutually absolutely continuous implies that

$$\mu(\{\omega \in \Omega : p_i(\omega) > 0 \forall i \in [r]\}) = 1. \quad (38)$$

So, without loss of generality, we assume that $p_i(\omega) > 0$ for all $\omega \in \Omega$ and $i \in [r]$.

We have from (10) that

$$\text{Err}_{\text{cl}}(\mathcal{E}_{\text{cl}}^n) = \int \mathbf{d}\mu^{\otimes n} (\eta_1 p_1^{\otimes n} \wedge \cdots \wedge \eta_r p_r^{\otimes n}) \leq \int \mathbf{d}\mu^{\otimes n} (p_1^{\otimes n} \wedge \cdots \wedge p_r^{\otimes n}). \quad (39)$$

Let $\mathbf{s} \in \mathbb{S}_r$ be arbitrary. We can write the right-hand side of (39) as

$$\int \mathbf{d}\mu^{\otimes n} (p_1^{\otimes n} \wedge \cdots \wedge p_r^{\otimes n}) = \int \mathbf{d}\mu^{\otimes n} (p_1^{\otimes n} \wedge \cdots \wedge p_r^{\otimes n})^{s_1} \cdots (p_1^{\otimes n} \wedge \cdots \wedge p_r^{\otimes n})^{s_r} \quad (40)$$

$$\leq \int \mathbf{d}\mu^{\otimes n} (p_1^{\otimes n})^{s_1} \cdots (p_r^{\otimes n})^{s_r}. \quad (41)$$

The expression on the right-hand side of (41) has a product structure. Indeed, for every generic $\omega^n := (\omega_1, \dots, \omega_n) \in \Omega^n$, we have that

$$\int \mathbf{d}\mu^{\otimes n}(\omega^n) (p_1^{\otimes n}(\omega^n))^{s_1} \cdots (p_r^{\otimes n}(\omega^n))^{s_r} = \int \prod_{k \in [n]} \mathbf{d}\mu(\omega_k) \left(\prod_{k \in [n]} p_1(\omega_k) \right)^{s_1} \cdots \left(\prod_{k \in [n]} p_r(\omega_k) \right)^{s_r} \quad (42)$$

$$= \int \prod_{k \in [n]} \mathbf{d}\mu(\omega_k) \prod_{k \in [n]} p_1^{s_1}(\omega_k) \cdots p_r^{s_r}(\omega_k) \quad (43)$$

$$= \int \prod_{k \in [n]} (\mathbf{d}\mu(\omega_k) p_1^{s_1}(\omega_k) \cdots p_r^{s_r}(\omega_k)) \quad (44)$$

$$= \prod_{k \in [n]} \int \mathbf{d}\mu(\omega_k) (p_1^{s_1} \cdots p_r^{s_r})(\omega_k) \quad (45)$$

$$= \left(\int \mathbf{d}\mu p_1^{s_1} \cdots p_r^{s_r} \right)^n \quad (46)$$

$$= \mathbb{H}_{\mathbf{s}}(P_1, \dots, P_r)^n. \quad (47)$$

From (39), (41), and (47), we thus get

$$\text{Err}_{\text{cl}}(\mathcal{E}_{\text{cl}}^n) \leq \mathbb{H}_{\mathbf{s}}(P_1, \dots, P_r)^n, \quad \text{for all } \mathbf{s} \in \mathbb{S}_r. \quad (48)$$

This implies, for all $n \in \mathbb{N}$, that

$$-\frac{1}{n} \ln \text{Err}_{\text{cl}}(\mathcal{E}_{\text{cl}}^n) \geq -\ln \inf_{\mathbf{s} \in \mathbb{S}_r} \mathbb{H}_{\mathbf{s}}(P_1, \dots, P_r) = \xi_{\text{cl}}(P_1, \dots, P_r). \quad (49)$$

Therefore, we get

$$\liminf_{n \rightarrow \infty} -\frac{1}{n} \ln \text{Err}_{\text{cl}}(\mathcal{E}_{\text{cl}}^n) \geq \xi_{\text{cl}}(P_1, \dots, P_r). \quad (50)$$

This proves the achievability part of the optimal error exponent.

To prove the optimality part, we apply multivariable calculus and the law of large numbers. For this purpose, let us parameterize the unit simplex of \mathbb{R}^r by the corner of the standard unit cube of \mathbb{R}^{r-1} , defined as

$$\mathbb{T}_r := \left\{ \mathbf{t} \in [0, 1]^{r-1} : \mathbf{t} := (t_1, \dots, t_{r-1}), \sum_{i \in [r-1]} t_i \leq 1 \right\}. \quad (51)$$

The unit simplex (18) can be expressed as

$$\mathbb{S}_r = \left\{ (t_1, \dots, t_{r-1}, 1 - \sum_{i \in [r-1]} t_i) : (t_1, \dots, t_{r-1}) \in \mathbb{T}_r \right\}. \quad (52)$$

Using the new parameterization, let us denote the elements of \mathbb{S}_r by $\mathbf{s}_{\mathbf{t}} := (t_1, \dots, t_{r-1}, 1 - \sum_{i \in [r-1]} t_i)$ for $\mathbf{t} := (t_1, \dots, t_{r-1}) \in \mathbb{T}_r$. The Hellinger transform of P_1, \dots, P_r can then be expressed as the following function on \mathbb{T}_r :

$$\mathbb{H}(\mathbf{t}) := \mathbb{H}_{\mathbf{s}_{\mathbf{t}}}(P_1, \dots, P_r), \quad \text{for } \mathbf{t} \in \mathbb{T}_r. \quad (53)$$

Thus, the multivariate Chernoff divergence of P_1, \dots, P_r has the form

$$\xi_{\text{cl}}(P_1, \dots, P_r) = \sup_{\mathbf{t} \in \mathbb{T}_r} -\ln \mathbb{H}(\mathbf{t}). \quad (54)$$

In what follows, using the reparametrized Hellinger transform (53), we define an *exponential family* of densities $p_{\mathbf{t}}$, as given in (59) and with $\mathbf{t} \in \mathbb{T}_r$, which enables us to express each $p_i^{\otimes n}$ in terms of $p_{\mathbf{t}}^{\otimes n}$ for all $n \in \mathbb{N}$. This then allows for the use of the law of large numbers to deduce a family of upper bounds on the asymptotic error exponent, given by $-\min_{1 \leq i \leq r} \gamma_i(\mathbf{t})$ for the *non-corner* points in \mathbb{T}_r , as defined later on in (76). Lastly, we use multivariable calculus to prove that there exists a non-corner point \mathbf{t}^* such that $\ln \mathbb{H}(\mathbf{t}^*) = \min_{1 \leq i \leq r} \gamma_i(\mathbf{t}^*)$. This implies that the multivariate Chernoff divergence is the optimal bound for the asymptotic error rate.

For every $\mathbf{t} \in \mathbb{T}_r$, let us express $\mathbb{H}(\mathbf{t})$ in an *exponential-integral* form as follows:

$$\mathbb{H}(\mathbf{t}) = \int \mathbf{d}\mu p_1^{t_1} \cdots p_{r-1}^{t_{r-1}} p_r^{1 - \sum_{i \in [r-1]} t_i} \quad (55)$$

$$= \int \mathbf{d}\mu (p_1/p_r)^{t_1} \cdots (p_{r-1}/p_r)^{t_{r-1}} p_r \quad (56)$$

$$= \int \mathbf{d}\mu p_r \exp\left(\sum_{i \in [r-1]} t_i \ln(p_i/p_r) \right) \quad (57)$$

$$= \int \mathbf{d}\mu p_r \exp\left(\sum_{i \in [r-1]} t_i q_i \right), \quad (58)$$

where $q_i := \ln(p_i/p_r)$. Consider the following *exponential family* of densities with respect to μ given by

$$p_{\mathbf{t}} := \frac{1}{H(\mathbf{t})} p_r \exp\left(\sum_{j \in [r-1]} t_j q_j\right), \quad \mathbf{t} \in \mathbb{T}_r. \quad (59)$$

Also, define a function $K : \mathbb{T}_r \rightarrow \mathbb{R}$ by

$$K(\mathbf{t}) := \ln H(\mathbf{t}). \quad (60)$$

Let $\mathbf{e}_1, \dots, \mathbf{e}_{r-1}$ denote the standard unit vectors in \mathbb{R}^{r-1} , and let \mathbb{T}_r° denote the interior of \mathbb{T}_r which is given by

$$\mathbb{T}_r^\circ := \left\{ (t_1, \dots, t_{r-1}) \in (0, 1)^{r-1} : \sum_{i \in [r-1]} t_i < 1 \right\}. \quad (61)$$

By Theorem 2.64 of [Sch12] we know that H is a smooth function on \mathbb{T}_r° ; also its partial derivatives are given for $\mathbf{t} \in \mathbb{T}_r^\circ$ and $i \in [r-1]$ by

$$\partial_i H(\mathbf{t}) := \lim_{h \rightarrow 0} \frac{H(\mathbf{t} + h\mathbf{e}_i) - H(\mathbf{t})}{h} \quad (62)$$

$$= \int \mathbf{d}\mu q_i p_r \exp\left(\sum_{j \in [r-1]} t_j q_j\right) \quad (63)$$

$$= H(\mathbf{t}) \mathbb{E}_{\mathbf{t}}[q_i], \quad (64)$$

where $\mathbb{E}_{\mathbf{t}}$ is the expectation under the density $p_{\mathbf{t}}$. We know that the Hellinger transform is a continuous function taking only positive values on \mathbb{S}_r . This implies that K is a real-valued continuous function on \mathbb{T}_r . Additionally, the smoothness of H on \mathbb{T}_r° implies the smoothness of K on \mathbb{T}_r° . From (64) we have that

$$\partial_i K(\mathbf{t}) = \frac{1}{H(\mathbf{t})} \partial_i H(\mathbf{t}) = \mathbb{E}_{\mathbf{t}}[q_i], \quad i \in [r-1], \mathbf{t} \in \mathbb{T}_r^\circ. \quad (65)$$

By Theorem 18.7 of [Das11], the Hessian matrix of K at $\mathbf{t} \in \mathbb{T}_r^\circ$ is the covariance matrix of the random vector $X := (q_1, \dots, q_{r-1})$ with the probability density $p_{\mathbf{t}}$, whence the Hessian matrix of K is positive semidefinite at each point in \mathbb{T}_r° . This implies that K is a convex function on \mathbb{T}_r° . By continuity, K is convex on \mathbb{T}_r . Let \mathbb{T}_r^1 denote the set

$$\mathbb{T}_r^1 := \left\{ (t_1, \dots, t_{r-1}) \in \mathbb{T}_r : \sum_{i \in [r-1]} t_i < 1 \right\}. \quad (66)$$

We call \mathbb{T}_r^1 the set of *non-corner* points of \mathbb{T}_r . It is easy to see that $\mathbb{T}_r^\circ \subset \mathbb{T}_r^1$. For any $\mathbf{t} \in \mathbb{T}_r^1$ and $i \in [r-1]$, the limit

$$\partial_i^+ K(\mathbf{t}) := \lim_{h \searrow 0} \frac{K(\mathbf{t} + h\mathbf{e}_i) - K(\mathbf{t})}{h} \quad (67)$$

exists in $\mathbb{R} \cup \{-\infty\}$ (Lemma 21). Observe that for $\mathbf{t} \in \mathbb{T}_r^\circ$, we have $\partial_i K(\mathbf{t}) = \partial_i^+ K(\mathbf{t})$ for all $i \in [r-1]$. It is shown in Appendix A that for $\mathbf{t} \in \mathbb{T}_r^1$ and $i \in [r-1]$, the expectation value $\mathbb{E}_{\mathbf{t}}[q_i]$ exists in $\mathbb{R} \cup \{-\infty\}$ and satisfies

$$\partial_i^+ K(\mathbf{t}) = \mathbb{E}_{\mathbf{t}}[q_i]. \quad (68)$$

Define a set

$$\mathbb{T}_{r,f}^1 := \{\mathbf{t} \in \mathbb{T}_r^1 : \partial_i^+ K(\mathbf{t}) \neq -\infty, \forall i \in [r-1]\}. \quad (69)$$

Note that $\mathbb{T}_r^\circ \subset \mathbb{T}_{r,f}^1$.

Using the definition (59) of the density $p_{\mathbf{t}}$ for $\mathbf{t} := (t_1, \dots, t_{r-1}) \in \mathbb{T}_{r,f}^1$ and $i \in [r]$, we have

$$\ln \frac{p_i}{p_{\mathbf{t}}} = \ln p_i - \ln p_{\mathbf{t}} \quad (70)$$

$$= \ln p_i - \sum_{j \in [r-1]} t_j q_j - \ln p_r + \ln H(\mathbf{t}) \quad (71)$$

$$= \ln \frac{p_i}{p_r} - \sum_{j \in [r-1]} t_j q_j + K(\mathbf{t}) \quad (72)$$

$$= q_i - \sum_{j \in [r-1]} t_j q_j + K(\mathbf{t}), \quad (73)$$

where q_r is the zero function on Ω . We write (73) in a more compact form as

$$\ln \frac{p_i}{p_{\mathbf{t}}} = \sum_{j \in [r-1]} (\delta_{ij} - t_j) q_j + K(\mathbf{t}), \quad \text{for } i \in [r], \quad \mathbf{t} \in \mathbb{T}_{r,f}^1. \quad (74)$$

Here δ_{ij} is the Kronecker delta (taking the value 1 if $i = j$, and 0 otherwise). By taking the expectation on both sides of (74) under the density $p_{\mathbf{t}}$, and then using (68), we get

$$\gamma_i(\mathbf{t}) := \mathbb{E}_{\mathbf{t}} \left[\ln \frac{p_i}{p_{\mathbf{t}}} \right] = \sum_{j \in [r-1]} (\delta_{ij} - t_j) \mathbb{E}_{\mathbf{t}}[q_j] + K(\mathbf{t}) \quad (75)$$

$$= \sum_{j \in [r-1]} (\delta_{ij} - t_j) \partial_i^+ K(\mathbf{t}) + K(\mathbf{t}) \quad (76)$$

for all $i \in [r]$ and $\mathbf{t} \in \mathbb{T}_{r,f}^1$. We can write (76) in a more compact form as

$$\gamma_i(\mathbf{t}) = \begin{cases} \partial_i^+ K(\mathbf{t}) - \mathbf{t}^T \nabla^+ K(\mathbf{t}) + K(\mathbf{t}), & i \in [r-1], \\ -\mathbf{t}^T \nabla^+ K(\mathbf{t}) + K(\mathbf{t}), & i = r, \end{cases} \quad \text{for } \mathbf{t} \in \mathbb{T}_{r,f}^1, \quad (77)$$

where $\nabla^+ K(\mathbf{t}) := (\partial_1^+ K(\mathbf{t}), \dots, \partial_{r-1}^+ K(\mathbf{t}))^T$.

Let $\omega^n := (\omega_1, \dots, \omega_n) \in \Omega^n$ and $\mathbf{t} \in \mathbb{T}_{r,f}^1$ be arbitrary. We have that

$$p_i^{\otimes n}(\omega^n) = \left(\prod_{j \in [n]} \frac{p_i}{p_{\mathbf{t}}}(\omega_j) \right) p_{\mathbf{t}}^{\otimes n}(\omega^n) = \exp(n G_{\mathbf{t},n}^{(i)}(\omega^n)) p_{\mathbf{t}}^{\otimes n}(\omega^n), \quad (78)$$

where

$$G_{\mathbf{t},n}^{(i)}(\omega^n) := \frac{1}{n} \sum_{j \in [n]} \ln \frac{p_i}{p_{\mathbf{t}}}(\omega_j), \quad \text{for } i \in [r]. \quad (79)$$

Let $P_{\mathbf{t}}^{\otimes n}$ be the product measure corresponding to the density $p_{\mathbf{t}}^{\otimes n}$, and let $\mathbb{E}_{\mathbf{t}}^n$ be the pertaining expectation. By the definition in (75), we then have that

$$\mathbb{E}_{\mathbf{t}}^n \left[G_{\mathbf{t},n}^{(i)} \right] = \gamma_i(\mathbf{t}), \quad \text{for } i \in [r]. \quad (80)$$

Since $G_{\mathbf{t},n}^{(i)}$ is an i.i.d. average, the law of large numbers [Bil95] implies that for arbitrary $\delta > 0$, there exists $n_\delta \in \mathbb{N}$ such that the probability of the event

$$U_{n,\delta} := \{\omega^n \in \Omega^n : \forall i \in [r], G_{\mathbf{t},n}^{(i)}(\omega^n) \geq \gamma_i(\mathbf{t}) - \delta\} \quad (81)$$

satisfies

$$P_{\mathbf{t}}^{\otimes n}(U_{n,\delta}) \geq 1 - \delta, \quad \text{for } n \geq n_\delta. \quad (82)$$

The development in (78)–(82) implies that, for all $n \geq n_\delta$,

$$\text{Err}_{\text{cl}}(\mathcal{E}_{\text{cl}}^n) = \int \mathbf{d}\mu^{\otimes n} (\eta_1 p_1^{\otimes n} \wedge \cdots \wedge \eta_r p_r^{\otimes n}) \quad (83)$$

$$\geq \eta_{\min} \int \mathbf{d}\mu^{\otimes n} (p_1^{\otimes n} \wedge \cdots \wedge p_r^{\otimes n}) \quad (84)$$

$$= \eta_{\min} \int \mathbf{d}\mu^{\otimes n} \left(\exp(n G_{\mathbf{t},n}^{(1)}) \wedge \cdots \wedge \exp(n G_{\mathbf{t},n}^{(r)}) \right) p_{\mathbf{t}}^{\otimes n} \quad (85)$$

$$= \eta_{\min} \mathbb{E}_{\mathbf{t}}^n \left[\exp(n G_{\mathbf{t},n}^{(1)}) \wedge \cdots \wedge \exp(n G_{\mathbf{t},n}^{(r)}) \right] \quad (86)$$

$$\geq \eta_{\min} \mathbb{E}_{\mathbf{t}}^n \left[\mathbf{1}_{U_{n,\delta}} \left(\exp(n G_{\mathbf{t},n}^{(1)}) \wedge \cdots \wedge \exp(n G_{\mathbf{t},n}^{(r)}) \right) \right] \quad (87)$$

$$\geq \eta_{\min} P_{\mathbf{t}}^{\otimes n}(U_{n,\delta}) \exp\left(n \min_{1 \leq i \leq r} (\gamma_i(\mathbf{t}) - \delta)\right) \quad (88)$$

$$\geq \eta_{\min}(1 - \delta) \exp\left(n \min_{1 \leq i \leq r} \gamma_i(\mathbf{t}) - n\delta\right). \quad (89)$$

Here $\mathbf{1}_{U_{n,\delta}}$ denotes the indicator function of the set $U_{n,\delta}$. Therefore, we have that

$$-\frac{1}{n} \ln \text{Err}_{\text{cl}}(\mathcal{E}_{\text{cl}}^n) \leq -\frac{\ln(\eta_{\min}(1 - \delta))}{n} - \left(\min_{1 \leq i \leq r} \gamma_i(\mathbf{t}) - \delta \right), \quad \text{for } n \geq n_\delta. \quad (90)$$

By taking the limit superior as $n \rightarrow \infty$ on both sides of (90) and then the limit $\delta \rightarrow 0$, we thus get

$$\limsup_{n \rightarrow \infty} -\frac{1}{n} \ln \text{Err}_{\text{cl}}(\mathcal{E}_{\text{cl}}^n) \leq -\min_{1 \leq i \leq r} \gamma_i(\mathbf{t}), \quad \text{for } \mathbf{t} \in \mathbb{T}_{r,f}^1. \quad (91)$$

Recall from (54) and the fact $K(\mathbf{t}) = \ln H(\mathbf{t})$, our goal is to show that

$$\limsup_{n \rightarrow \infty} -\frac{1}{n} \ln \text{Err}_{\text{cl}}(\mathcal{E}_{\text{cl}}^n) \leq \sup_{\mathbf{t} \in \mathbb{T}_r} -K(\mathbf{t}). \quad (92)$$

In view of (91), it suffices to show that for some $\mathbf{t}^* \in \mathbb{T}_{r,f}^1$, the following holds

$$\min_{1 \leq i \leq r} \gamma_i(\mathbf{t}^*) \geq K(\mathbf{t}^*). \quad (93)$$

We now argue that such a \mathbf{t}^* exists. Since K is a continuous function on the compact set \mathbb{T}_r , there exists $\mathbf{t}^* := (t_1^*, \dots, t_{r-1}^*) \in \mathbb{T}_r$ that minimizes K over \mathbb{T}_r , i.e.,

$$K(\mathbf{t}^*) = \min_{\mathbf{t} \in \mathbb{T}_r} K(\mathbf{t}). \quad (94)$$

Consider the following two cases.

Case A: Suppose $\mathbf{t}^* \in \mathbb{T}_r^1$. Choose arbitrary $i \in [r-1]$. If $t_i^* = 0$, then by convexity, continuity of K , and the fact that \mathbf{t}^* is a minimizer, we have $\partial_i^+ K(\mathbf{t}^*) \geq 0$. (see Lemma 21). Else we have $0 < t_i^* < 1$ and the first order necessary condition for a minimizer implies $\partial_i^+ K(\mathbf{t}^*) = 0$. Combining these, we get $\partial_i^+ K(\mathbf{t}^*) \geq 0$ for all $i \in [r-1]$ and hence $\mathbf{t}^* \in \mathbb{T}_{r,f}^1$, and $\mathbf{t}^{*T} \nabla^+ K(\mathbf{t}^*) = 0$. From (77), we thus get

$$\gamma_i(\mathbf{t}^*) = \begin{cases} \partial_i^+ K(\mathbf{t}^*) + K(\mathbf{t}^*), & i \in [r-1], \\ K(\mathbf{t}^*), & i = r. \end{cases} \quad (95)$$

This implies that the inequality (93) holds for the minimizer \mathbf{t}^* .

Case B: Suppose $\mathbf{t}^* \in \mathbb{T}_r \setminus \mathbb{T}_r^1$, i.e., $t_1^* + \dots + t_{r-1}^* = 1$. For some $i \in [r-1]$, we have $t_i^* > 0$. According to the current parameterization of the unit simplex given in (52), \mathbf{t}^* corresponds to the vector $(t_1^*, \dots, t_{r-1}^*, 0)$ in \mathbb{S}_r . We reparameterize the unit simplex \mathbb{S}_r as

$$\mathbf{s}_{\mathbf{u}} = \left(u_1, \dots, u_i, 1 - \sum_{j \in [r-1]} u_j, u_{i+1}, \dots, u_{r-1} \right), \quad \mathbf{u} \in \mathbb{T}_r. \quad (96)$$

In the reparameterized problem, the corresponding minimizer \mathbf{u}^* of K satisfies $\mathbf{s}_{\mathbf{u}^*} = (t_1^*, \dots, t_{r-1}^*, 0)$, which implies

$$1 - \sum_{j \in [r-1]} u_j^* = t_i^* > 0. \quad (97)$$

This reduces the problem to **Case A**, which implies that (93) holds.

Combining the above two cases, we conclude that (93) holds for the minimizer \mathbf{t}^* and this completes the proof. \blacksquare

Theorem 7. Consider an ensemble $\mathcal{E}_{\text{cl}} = \{(\eta_i, P_i) : i \in [r]\}$ of probability measures, with possibly unequal supports, on a measurable space (Ω, \mathcal{A}) . The optimal error exponent for antidistinguishing the probability measures is given by their multivariate Chernoff divergence, i.e.,

$$E_{\text{cl}}(P_1, \dots, P_r) = \lim_{n \rightarrow \infty} -\frac{1}{n} \ln \text{Err}_{\text{cl}}(\mathcal{E}_{\text{cl}}^n) = \xi_{\text{cl}}(P_1, \dots, P_r), \quad (98)$$

where, recalling (18), (19), and (23), the multivariate classical Chernoff divergence ξ_{cl} is defined as

$$\xi_{\text{cl}}(P_1, \dots, P_r) := -\ln \inf_{\mathbf{s} \in \mathbb{S}_r} \int d\mu p_1^{s_1} \dots p_r^{s_r}. \quad (99)$$

Proof. See Appendix B for a detailed proof. \blacksquare

A. Multivariate Chernoff divergence versus pairwise Chernoff divergences

Identifying the true probability measure out of the given r probability measures is the same as eliminating all the remaining $r-1$ false probability measures. As such, general intuition says that, upon observing i.i.d. data, it is easier to eliminate a false probability measure than to identify the true probability measure. This also means that the optimal error exponent of classical antidistinguishability should be greater than that of multiple classical hypothesis testing, the

former being the multivariate classical Chernoff divergence and the latter being the minimum of the pairwise Chernoff divergences of the probability measures [Sal73] (see also [Tor81, Theorem 4.2] and [LJ97, Sal99, Sal03]). Indeed, for any two indices $i, j \in [r]$ define a subset of \mathbb{S}_r :

$$\mathbb{S}_r^{(i,j)} = \{\mathbf{s} \in \mathbb{S}_r : \mathbf{s} := (s_1, \dots, s_r), s_i + s_j = 1\}. \quad (100)$$

By definition, we have

$$\xi_{\text{cl}}(P_1, \dots, P_r) \geq -\ln \inf_{\mathbf{s} \in \mathbb{S}_r^{(i,j)}} \mathbb{H}_{\mathbf{s}}(P_1, \dots, P_r) \quad (101)$$

$$= -\ln \inf_{s \in [0,1]} \int \mathbf{d}\mu p_i^s p_j^{(1-s)} \quad (102)$$

$$= \xi_{\text{cl}}(P_i, P_j), \quad (103)$$

where $\xi_{\text{cl}}(P_i, P_j)$ is the Chernoff divergence of the probability measures P_i and P_j . This gives

$$\xi_{\text{cl}}(P_1, \dots, P_r) \geq \max_{i < j} \xi_{\text{cl}}(P_i, P_j) \geq \min_{i < j} \xi_{\text{cl}}(P_i, P_j). \quad (104)$$

The following example illustrates an instance for which the first inequality in (104) is strict.

Example 8. Consider a uniform ensemble $\mathcal{E}_{\text{cl}} = \{(1/3, P_1), (1/3, P_2), (1/3, P_3)\}$ of probability measures on a discrete space $\Omega = \{x, y, z\}$ whose densities with respect to the counting measure μ are given by

$$p_1 = \frac{1}{2} \mathbf{1}_{\{x,y\}}, \quad p_2 = \frac{1}{2} \mathbf{1}_{\{x,z\}}, \quad p_3 = \frac{1}{3} \mathbf{1}_{\Omega}. \quad (105)$$

We have for $\omega^n \in \Omega^n$,

$$(p_1^{\otimes n} \wedge p_2^{\otimes n} \wedge p_3^{\otimes n})(\omega^n) = \begin{cases} \frac{1}{3^n}, & \text{if } \omega^n = \underbrace{(x, \dots, x)}_{n \text{ times}}, \\ 0, & \text{otherwise.} \end{cases} \quad (106)$$

By the minimum likelihood principle, we thus get

$$\text{Err}_{\text{cl}}(\mathcal{E}_{\text{cl}}^n) = \frac{1}{3} \int \mathbf{d}\mu^{\otimes n} (p_1^{\otimes n} \wedge p_2^{\otimes n} \wedge p_3^{\otimes n}) \quad (107)$$

$$= \frac{1}{3} \cdot \mu^{\otimes n}(\underbrace{\{(x, \dots, x)\}}_{n \text{ times}}) \cdot \frac{1}{3^n} \quad (108)$$

$$= \frac{1}{3} \cdot \mu(\{x\})^n \cdot \frac{1}{3^n} \quad (109)$$

$$= \frac{1}{3} \cdot 1 \cdot \frac{1}{3^n} \quad (110)$$

$$= \frac{1}{3^{n+1}}. \quad (111)$$

This gives the optimal error exponent

$$E_{\text{cl}}(P_1, P_2, P_3) = \liminf_{n \rightarrow \infty} -\frac{1}{n} \ln \text{Err}_{\text{cl}}(\mathcal{E}_{\text{cl}}^n) = \ln 3. \quad (112)$$

We now compute the pairwise Chernoff divergences of the probability measures as follows.

$$\xi_{\text{cl}}(P_1, P_2) = -\ln \inf_{s \in [0,1]} \int \mathbf{d}\mu p_1^s p_2^{(1-s)} \quad (113)$$

$$= -\ln \inf_{s \in [0,1]} \int_{\{x\}} \mathbf{d}\mu \frac{1}{2^s} \frac{1}{2^{(1-s)}} \quad (114)$$

$$= -\ln \inf_{s \in [0,1]} \frac{1}{2^s} \frac{1}{2^{(1-s)}} \quad (115)$$

$$= -\ln \left(\frac{1}{2} \right) \quad (116)$$

$$= \ln 2. \quad (117)$$

Also,

$$\xi_{\text{cl}}(P_1, P_3) = -\ln \inf_{s \in [0,1]} \int \mathbf{d}\mu p_1^s p_3^{(1-s)} \quad (118)$$

$$= -\ln \inf_{s \in [0,1]} \int_{\{x,y\}} \mathbf{d}\mu \frac{1}{2^s} \frac{1}{3^{(1-s)}} \quad (119)$$

$$= -\ln \left[\mu(\{x, y\}) \cdot \frac{1}{3} \cdot \inf_{s \in [0,1]} \left(\frac{3}{2} \right)^s \right] \quad (120)$$

$$= -\ln \left[2 \cdot \frac{1}{3} \cdot 1 \right] \quad (121)$$

$$= \ln(3/2). \quad (122)$$

By similar arguments, we get $\xi_{\text{cl}}(P_2, P_3) = \ln(3/2)$. This implies

$$\max\{\xi_{\text{cl}}(P_1, P_2), \xi_{\text{cl}}(P_1, P_3), \xi_{\text{cl}}(P_2, P_3)\} = \ln 2. \quad (123)$$

$$\min\{\xi_{\text{cl}}(P_1, P_2), \xi_{\text{cl}}(P_1, P_3), \xi_{\text{cl}}(P_2, P_3)\} = \ln(3/2). \quad (124)$$

From (112), (123), and (124), we have

$$E_{\text{cl}}(P_1, P_2, P_3) > \max\{\xi_{\text{cl}}(P_1, P_2), \xi_{\text{cl}}(P_1, P_3), \xi_{\text{cl}}(P_2, P_3)\} \quad (125)$$

$$> \min\{\xi_{\text{cl}}(P_1, P_2), \xi_{\text{cl}}(P_1, P_3), \xi_{\text{cl}}(P_2, P_3)\}. \quad (126)$$

IV. ACHIEVABLE ERROR EXPONENT FOR QUANTUM ANTIDISTINGUISHABILITY

A. One-shot case

Observe that the “antidistinguishability problem” between any two states ρ_1 and ρ_2 is the same as the state discrimination problem. Indeed, if we say that “ ρ_1 is not the true state,” then we are saying “ ρ_2 is the true state.” Using this observation, we obtain an upper bound on the optimal error probability of antidistinguishing the states of a given quantum ensemble by considering “special” POVMs that focus on pairs of states, as expounded upon in the proof of the following theorem:

Theorem 9. Consider a quantum ensemble $\mathcal{E} = \{(\eta_i, \rho_i) : i \in [r]\}$. An upper bound on the optimal error probability of antidistinguishing the states of the ensemble is given by

$$\text{Err}(\mathcal{E}) \leq \min_{1 \leq i < j \leq r} \text{Tr}[\eta_i \rho_i \wedge \eta_j \rho_j] \quad (127)$$

$$= \min_{1 \leq i < j \leq r} \frac{1}{2} \left(\eta_i + \eta_j - \|\eta_i \rho_i - \eta_j \rho_j\|_1 \right). \quad (128)$$

In particular, if at least two states in ρ_1, \dots, ρ_r are mutually orthogonal then $\text{Err}(\mathcal{E}) = 0$.

Proof. Given two fixed indices $i, j \in [r]$, let $\Xi_r^{(i,j)}$ denote the set of POVMs $\mathcal{M} = \{M_1, \dots, M_r\}$ such that $M_k = 0$ if $k \notin \{i, j\}$. For such POVMs, we have $M_i + M_j = \mathbb{I}$ and

$$\text{Err}(\mathcal{M}; \mathcal{E}) = \eta_i \text{Tr}[M_i \rho_i] + \eta_j \text{Tr}[(\mathbb{I} - M_i) \rho_j] \quad (129)$$

$$= \eta_i \text{Tr}[M_i \rho_i] + \eta_j - \eta_j \text{Tr}[M_i \rho_j] \quad (130)$$

$$= \eta_j - \text{Tr}[M_i (\eta_j \rho_j - \eta_i \rho_i)]. \quad (131)$$

By taking the infimum over $\Xi_r^{(i,j)}$ on both sides of (131), we get

$$\inf_{\mathcal{M} \in \Xi_r^{(i,j)}} \text{Err}(\mathcal{M}; \mathcal{E}) = \eta_j - \sup_{0 \leq M_i \leq \mathbb{I}} \text{Tr}[M_i (\eta_j \rho_j - \eta_i \rho_i)], \quad (132)$$

where the supremum on the right-hand side of (132) is taken over every positive semi-definite operator M_i such that $0 \leq M_i \leq \mathbb{I}$. The supremum is attained by the *Helstrom–Holevo measurement* [Hel69, Hol72] given by $M_i = \text{Supp}(\eta_j \rho_j - \eta_i \rho_i)_+$ (note the order of i and j). We thus get

$$\inf_{\mathcal{M} \in \Xi_r^{(i,j)}} \text{Err}(\mathcal{M}; \mathcal{E}) = \eta_j - \text{Tr}[(\eta_j \rho_j - \eta_i \rho_i)_+] \quad (133)$$

$$= \text{Tr}[\eta_j \rho_j] - \text{Tr}[(\eta_j \rho_j - \eta_i \rho_i + |\eta_j \rho_j - \eta_i \rho_i|) / 2] \quad (134)$$

$$= \text{Tr}[(\eta_i \rho_i + \eta_j \rho_j - |\eta_i \rho_i - \eta_j \rho_j|) / 2] \quad (135)$$

$$= \text{Tr}[\eta_i \rho_i \wedge \eta_j \rho_j] \quad (136)$$

$$= \frac{1}{2} \left(\eta_i + \eta_j - \|\eta_i \rho_i - \eta_j \rho_j\|_1 \right). \quad (137)$$

It is clear that the optimal antidistinguishability error probability satisfies

$$\text{Err}(\mathcal{E}) \leq \inf_{\mathcal{M} \in \Xi_r^{(i,j)}} \text{Err}(\mathcal{M}; \mathcal{E}) = \text{Tr}[\eta_i \rho_i \wedge \eta_j \rho_j], \quad \text{for all } 1 \leq i < j \leq r. \quad (138)$$

By combining (136)–(138), we thus get the upper bound on the optimal antidistinguishability error probability stated in the theorem. ■

The expression on the right-hand side of (127) can be further simplified for pure states. This is a consequence of the following identity (see Proposition 22 in Appendix C):

$$\|\varphi \chi \varphi - |\zeta \chi \zeta|\|_1^2 = (\langle \varphi | \varphi \rangle + \langle \zeta | \zeta \rangle)^2 - 4 |\langle \zeta | \varphi \rangle|^2, \quad (139)$$

which holds for vectors $|\varphi\rangle$ and $|\zeta\rangle$, as well as Theorem 1 of [ACMT⁺07] which states that for all positive semi-definite operators A, B and all $0 \leq s \leq 1$, we have

$$\text{Tr}[A \wedge B] \leq \text{Tr} A^s B^{1-s}. \quad (140)$$

Corollary 10. *If the quantum states in Theorem 9 are pure, i.e., given by $\rho_i = |\psi_i\rangle\langle\psi_i|$, then we have*

$$\text{Err}(\mathcal{E}) \leq \min_{1 \leq i < j \leq r} \frac{\eta_i + \eta_j}{2} \left(1 - \sqrt{1 - \frac{4\eta_i\eta_j|\langle\psi_i|\psi_j\rangle|^2}{(\eta_i + \eta_j)^2}} \right) \quad (141)$$

$$\leq \frac{1}{2} \min_{1 \leq i < j \leq r} |\langle\psi_i|\psi_j\rangle|^2. \quad (142)$$

Proof. Applying (128) and (139), we find that for all $1 \leq i < j \leq r$,

$$\text{Err}(\mathcal{E}) \leq \frac{1}{2} \left(\eta_i + \eta_j - \|\eta_i|\psi_i\rangle\langle\psi_i| - \eta_j|\psi_j\rangle\langle\psi_j|\|_1 \right) \quad (143)$$

$$= \frac{1}{2} \left(\eta_i + \eta_j - \sqrt{(\eta_i + \eta_j)^2 - 4\eta_i\eta_j|\langle\psi_i|\psi_j\rangle|^2} \right) \quad (144)$$

$$= \frac{\eta_i + \eta_j}{2} \left(1 - \sqrt{1 - \frac{4\eta_i\eta_j|\langle\psi_i|\psi_j\rangle|^2}{(\eta_i + \eta_j)^2}} \right). \quad (145)$$

This proves the inequality (141). By (140), we get that for all $1 \leq i < j \leq r$ and $0 \leq s \leq 1$,

$$\text{Tr}[\eta_i|\psi_i\rangle\langle\psi_i| \wedge \eta_j|\psi_j\rangle\langle\psi_j|] \leq \eta_i^s \eta_j^{1-s} \text{Tr}[|\psi_i\rangle\langle\psi_i| |\psi_j\rangle\langle\psi_j|] = \eta_i^s \eta_j^{1-s} |\langle\psi_i|\psi_j\rangle|^2. \quad (146)$$

Since (146) holds for all $s \in [0, 1]$, we get

$$\text{Tr}[\eta_i|\psi_i\rangle\langle\psi_i| \wedge \eta_j|\psi_j\rangle\langle\psi_j|] \leq (\eta_i \wedge \eta_j) |\langle\psi_i|\psi_j\rangle|^2 \leq \frac{1}{2} |\langle\psi_i|\psi_j\rangle|^2. \quad (147)$$

The desired inequality (142) thus follows by using the inequality (147) in (127). \blacksquare

The sufficient condition for perfect antidistinguishability given in Theorem 9 is not a necessary condition, even in the simple case of commuting states. This is illustrated in the following example.

Example 11. Consider states ρ_1 , ρ_2 , and ρ_3 diagonalizable in a common eigenbasis $\{|1\rangle\langle 1|, |2\rangle\langle 2|, |3\rangle\langle 3|\}$, given by

$$\rho_1 = \frac{1}{2} (|1\rangle\langle 1| + |2\rangle\langle 2|), \quad (148)$$

$$\rho_2 = \frac{1}{2} (|1\rangle\langle 1| + |3\rangle\langle 3|), \quad (149)$$

$$\rho_3 = \frac{1}{2} (|2\rangle\langle 2| + |3\rangle\langle 3|). \quad (150)$$

Consider a POVM $\mathcal{M} = \{M_1, M_2, M_3\}$ given by

$$M_1 = |3\rangle\langle 3|, \quad (151)$$

$$M_2 = |2\rangle\langle 2|, \quad (152)$$

$$M_3 = |1\rangle\langle 1|. \quad (153)$$

The POVM \mathcal{M} antidistinguishes the states perfectly because $\text{Tr}[M_i \rho_i] = 0$ for $i \in [3]$. However, no pair of states are mutually orthogonal to each other.

B. Asymptotic case

As a consequence of Theorem 9, we arrive at a lower bound on the optimal error exponent, as stated in the following theorem.

Theorem 12. *Consider a quantum ensemble $\mathcal{E} = \{(\eta_i, \rho_i) : i \in [r]\}$. A lower bound on the optimal error exponent for antidistinguishing the states of the ensemble is given by the maximum of the pairwise Chernoff divergence of the states; i.e., we have*

$$E(\rho_1, \dots, \rho_r) \geq \max_{1 \leq i < j \leq r} \xi(\rho_i, \rho_j). \quad (154)$$

Proof. By Theorem 9, we have

$$\text{Err}(\mathcal{E}^n) \leq \min_{1 \leq i < j \leq r} \text{Tr}[\eta_i \rho_i^{\otimes n} \wedge \eta_j \rho_j^{\otimes n}]. \quad (155)$$

By combining (155) with (34), we get the desired inequality in (154). ■

Let us recall from Example 8 that the inequality in (154) can be strict in some cases.

Corollary 13. *If the quantum states in Theorem 12 are pure, given by $\rho_i = |\psi_i\rangle\langle\psi_i|$, then we have*

$$E(|\psi_1\rangle\langle\psi_1|, \dots, |\psi_r\rangle\langle\psi_r|) \geq \max_{1 \leq i < j \leq r} -\ln |\langle\psi_i|\psi_j\rangle|^2. \quad (156)$$

Proof. It follows directly from (154) and the representation (33) of the quantum Chernoff divergence for pure states. ■

V. BOUNDS ON THE OPTIMAL ERROR EXPONENT FOR QUANTUM ANTIDISTINGUISHABILITY FROM MULTIVARIATE QUANTUM CHERNOFF DIVERGENCES

In this section, we begin by introducing the general concept of multivariate quantum Chernoff divergences, and after that, we employ this concept in order to obtain bounds on the optimal error exponent for quantum antidistinguishability. The reasoning used here is inspired by similar reasoning used for distinguishability problems between two states [Mat13, Mat14, MO15, HT16, HM17, Mat18].

A. Multivariate quantum Chernoff divergences

Definition 14. *Let $r \geq 2$ be an integer. We call a function $\xi : \mathcal{D}^r \rightarrow [0, \infty]$ a multivariate quantum Chernoff divergence if it satisfies the following properties:*

1. *Data processing: for states ρ_1, \dots, ρ_r and a channel \mathcal{N} ,*

$$\xi(\rho_1, \dots, \rho_r) \geq \xi(\mathcal{N}(\rho_1), \dots, \mathcal{N}(\rho_r)), \quad (157)$$

2. *Reduction to the multivariate classical Chernoff divergence for commuting states: if the states*

ρ_1, \dots, ρ_r commute, then

$$\xi(\rho_1, \dots, \rho_r) = \xi_{\text{cl}}(P_1, \dots, P_r), \quad (158)$$

where ξ_{cl} is defined in (23), P_1, \dots, P_r are probability measures on $[\dim(\mathcal{H})]$,

$$P_\ell(X) := \sum_{i \in X} \lambda_{\ell,i}, \quad \text{for } X \subseteq [\dim(\mathcal{H})], \quad (159)$$

given by a spectral decomposition of the states in a common eigenbasis

$$\rho_\ell = \sum_{i \in [\dim(\mathcal{H})]} \lambda_{\ell,i} |i\rangle\langle i|, \quad \text{for } \ell \in [r]. \quad (160)$$

As stated above, all multivariate quantum Chernoff divergences agree on commuting states and are equal to the multivariate classical Chernoff divergence of the corresponding probability measures induced by the states in a common eigenbasis. If ρ_1, \dots, ρ_r are commuting states, then we denote their divergence by $\xi_{\text{cl}}(\rho_1, \dots, \rho_r)$. In this case, it is easy to verify that

$$\xi_{\text{cl}}(\rho_1, \dots, \rho_r) = -\ln \inf_{s \in \mathbb{S}_r} \sum_{i \in [\dim(\mathcal{H})]} \left(\prod_{\ell \in [r]} \lambda_{\ell,i}^{s_\ell} \right). \quad (161)$$

As a first starting point, let us explicitly note that the optimal error exponent in (31) is itself a multivariate quantum Chernoff divergence.

Proposition 15. *The optimal error exponent $E : \mathcal{D}^r \rightarrow [0, \infty]$ defined by (31) is a multivariate quantum Chernoff divergence.*

Proof. See Appendix D. ■

Let us note that other multivariate quantum Chernoff divergences can be constructed from the multivariate log-Euclidean divergence, as discussed in Remark 4 below, as well as by means of the multivariate quantum Rényi divergences proposed in [FLO23, MBV22]. In what follows, we discuss some other constructions of multivariate quantum Chernoff divergences.

We say that a multivariate quantum Chernoff divergence ξ_{\min} is minimal if it is a lower bound to any other multivariate quantum Chernoff divergence; i.e., for any multivariate quantum Chernoff divergence ξ , we have

$$\xi_{\min}(\rho_1, \dots, \rho_r) \leq \xi(\rho_1, \dots, \rho_r), \quad \text{for } (\rho_1, \dots, \rho_r) \in \mathcal{D}^r. \quad (162)$$

A minimal multivariate quantum Chernoff divergence is unique by definition, and it can be obtained as an optimization over *quantum-to-classical* or *measurement* channels as presented in Proposition 16 below.

Let \mathcal{K} be a complex Hilbert space of dimension t with an orthonormal basis $\{|1\rangle, \dots, |t\rangle\}$. Associated with a POVM $\{M_1, \dots, M_t\}$ acting on the Hilbert space \mathcal{H} is a channel \mathcal{M} , called measurement channel, which has the following action on an input state $\rho \in \mathcal{D}(\mathcal{H})$:

$$\mathcal{M}(\rho) = \sum_{\omega \in [t]} \text{Tr}[M_\omega \rho] |\omega\rangle\langle \omega|. \quad (163)$$

The action of the measurement channel on any given states ρ_1, \dots, ρ_r produces commuting states $\mathcal{M}(\rho_1), \dots, \mathcal{M}(\rho_r)$. This induces probability measures $P_1^{\mathcal{M}}, \dots, P_r^{\mathcal{M}}$ on the discrete space $\Omega = [t]$, defined by

$$P_i^{\mathcal{M}}(X) := \sum_{x \in X} \text{Tr}[M_x \rho_i], \quad \text{for } X \subseteq \Omega. \quad (164)$$

It can be easily verified that the optimal error probability of antidistinguishing the commuting states $\mathcal{M}(\rho_1), \dots, \mathcal{M}(\rho_r)$ is equal to that of antidistinguishing the corresponding probability measures $P_1^{\mathcal{M}}, \dots, P_r^{\mathcal{M}}$. See (D7)–(D14) in Appendix D.

Proposition 16. *The minimal multivariate quantum Chernoff divergence is given by*

$$\xi_{\min}(\rho_1, \dots, \rho_r) = \sup_{\mathcal{M}} \xi_{\text{cl}}(P_1^{\mathcal{M}}, \dots, P_r^{\mathcal{M}}), \quad (165)$$

where the supremum is taken over all measurement channels \mathcal{M} with a t -dimensional classical output space for all $t \in \mathbb{N}$ and each probability measure $P_i^{\mathcal{M}}$ is defined in (164).

Proof. See Appendix E. ■

Similar to the definition of minimal multivariate quantum Chernoff divergence, we can define the maximal multivariate quantum Chernoff divergence. We say that a multivariate quantum Chernoff divergence ξ_{\max} is maximal if it is an upper bound to any other multivariate quantum Chernoff divergence; i.e., for any multivariate quantum Chernoff divergence ξ , we have

$$\xi_{\max}(\rho_1, \dots, \rho_r) \geq \xi(\rho_1, \dots, \rho_r), \quad \text{for } (\rho_1, \dots, \rho_r) \in \mathcal{D}^r. \quad (166)$$

A maximal multivariate quantum Chernoff divergence is unique by definition, and it can be obtained as an optimization over *classical-to-quantum* or *preparation* channels as given in Proposition 17 below.

We can view any probability measure P on the discrete space $\Omega = [t]$ as a quantum state in \mathcal{K} with the fixed eigenbasis $\{|1\rangle\langle 1|, \dots, |t\rangle\langle t|\}$, i.e.,

$$P \equiv \sum_{\omega \in \Omega} P(\{\omega\}) |\omega\rangle\langle \omega|. \quad (167)$$

A quantum channel $\mathcal{P} : \mathcal{L}(\mathcal{K}) \rightarrow \mathcal{L}(\mathcal{H})$ is said to prepare a state $\rho \in \mathcal{D}(\mathcal{H})$ from a probability measure P if it satisfies $\mathcal{P}(P) = \rho$ and is called a preparation channel or classical-to-quantum channel (see [Wil17, Section 4.6.5] for a review of classical-to-quantum channels).

Proposition 17. *The maximal multivariate quantum Chernoff divergence is given by*

$$\xi_{\max}(\rho_1, \dots, \rho_r) = \inf_{(\mathcal{P}, \{P_i\}_{i \in [r]})} \{ \xi_{\text{cl}}(P_1, \dots, P_r) : \mathcal{P}(P_i) = \rho_i \text{ for all } i \in [r] \}, \quad (168)$$

where the infimum involves preparation channels \mathcal{P} with t -dimensional classical input system, for all $t \in \mathbb{N}$, as well as probability measures $\{P_1, \dots, P_r\}$ of the form in (167).

Proof. See Appendix F. ■

B. Bounds on the optimal error exponent for quantum antidistinguishability

The optimal error exponent for quantum antidistinguishability can be bounded from above and below by the minimal and the maximal multivariate quantum Chernoff divergences, respectively, as stated in the following theorem.

Theorem 18. *Let $\mathcal{E} = \{(\eta_i, \rho_i) : i \in [r]\}$ be a quantum ensemble. We have*

$$\xi_{\min}(\rho_1, \dots, \rho_r) \leq \mathbb{E}(\rho_1, \dots, \rho_r) \leq \xi_{\max}(\rho_1, \dots, \rho_r), \quad (169)$$

where ξ_{\min} and ξ_{\max} are given by (165) and (168), respectively. Additionally, the bounds in (169) can be strengthened through regularization as

$$\sup_{\ell \in \mathbb{N}} \frac{1}{\ell} \xi_{\min}(\rho_1^{\otimes \ell}, \dots, \rho_r^{\otimes \ell}) \leq \mathbb{E}(\rho_1, \dots, \rho_r) \leq \inf_{\ell \in \mathbb{N}} \frac{1}{\ell} \xi_{\max}(\rho_1^{\otimes \ell}, \dots, \rho_r^{\otimes \ell}). \quad (170)$$

Proof. We know from Proposition 15 that the optimal error exponent is a multivariate quantum Chernoff divergence, which, along with (162) and (166), justifies the inequalities in (169).

We know from Lemma 23 in Appendix G that

$$\mathbb{E}(\rho_1, \dots, \rho_r) = \frac{1}{\ell} \mathbb{E}(\rho_1^{\otimes \ell}, \dots, \rho_r^{\otimes \ell}) \quad \text{for all } \ell \in \mathbb{N}. \quad (171)$$

Substituting the above equality into (169) gives

$$\frac{1}{\ell} \xi_{\min}(\rho_1^{\otimes \ell}, \dots, \rho_r^{\otimes \ell}) \leq \mathbb{E}(\rho_1, \dots, \rho_r) \leq \frac{1}{\ell} \xi_{\max}(\rho_1^{\otimes \ell}, \dots, \rho_r^{\otimes \ell}) \quad \text{for all } \ell \in \mathbb{N}, \quad (172)$$

which implies the inequalities (170). ■

We note that in the upper bound in (170), the infimum over $\ell \in \mathbb{N}$ can be replaced with the limit $\ell \rightarrow \infty$:

$$\inf_{\ell \in \mathbb{N}} \frac{1}{\ell} \xi_{\max}(\rho_1^{\otimes \ell}, \dots, \rho_r^{\otimes \ell}) = \lim_{\ell \rightarrow \infty} \frac{1}{\ell} \xi_{\max}(\rho_1^{\otimes \ell}, \dots, \rho_r^{\otimes \ell}). \quad (173)$$

See Appendix H. It is open to determine whether the supremum over $\ell \in \mathbb{N}$ in the lower bound in (170) can be replaced with the limit $\ell \rightarrow \infty$, if the limit exists.

It is known from [MO15, Corollary III.8] and [HT16, Corollary 4] (see also [Mat14, Section 9.3]) that when $r = 2$, the following equality holds

$$\sup_{\ell \in \mathbb{N}} \frac{1}{\ell} \xi_{\min}(\rho_1^{\otimes \ell}, \rho_2^{\otimes \ell}) = \tilde{\xi}(\rho_1, \rho_2) := \sup_{s \in (0,1)} \left[-\ln \tilde{Q}_s(\rho_1, \rho_2) \right], \quad (174)$$

where

$$\tilde{Q}_s(\rho_1, \rho_2) := \begin{cases} \text{Tr} \left[\left(\rho_2^{(1-s)/2s} \rho_1 \rho_2^{(1-s)/2s} \right)^s \right] & : s \in [1/2, 1) \\ \text{Tr} \left[\left(\rho_1^{s/2(1-s)} \rho_2 \rho_1^{s/2(1-s)} \right)^{1-s} \right] & : s \in (0, 1/2) \end{cases}. \quad (175)$$

Since the optimal error exponent is known in this case to be $\xi(\rho_1, \rho_2)$, which is defined in (32), and it is also known from [DL14, Lemma 3] that

$$\xi(\rho_1, \rho_2) \geq \tilde{\xi}(\rho_1, \rho_2), \quad (176)$$

where the inequality is strict if ρ_1 and ρ_2 are invertible and do not commute (see [Hia94, Theorem 2.1]), it follows that the lower bound in (170) cannot be optimal in general.

It is also known from [Mat13, Mat18], that when $r = 2$, we have

$$\inf_{\ell \in \mathbb{N}} \frac{1}{\ell} \xi_{\max}(\rho_1^{\otimes \ell}, \rho_2^{\otimes \ell}) \geq \widehat{\xi}(\rho_1, \rho_2) := \sup_{s \in (0,1)} -\ln \widehat{Q}_s(\rho_1, \rho_2), \quad (177)$$

where

$$\widehat{Q}_s(\rho_1, \rho_2) := \text{Tr} \left[\rho_2 \left(\rho_2^{-1/2} \tilde{\rho}_1 \rho_2^{-1/2} \right)^s \right]. \quad (178)$$

Here $\tilde{\rho}_1$ is the absolutely continuous part of ρ_1 with respect to ρ_2 [And76], and the negative power of ρ_2 is taken in on its support. Since the optimal error exponent is known in this case to be $\xi(\rho_1, \rho_2)$ given in (32), and it is also known from [Mat13, Mat18] that

$$\xi(\rho_1, \rho_2) \leq \widehat{\xi}(\rho_1, \rho_2), \quad (179)$$

where the inequality is strict if ρ_1 and ρ_2 are invertible and do not commute (see [HM17, Theorem 4.3]), it follows that the upper bound in (170) cannot be the tightest possible upper bound in general.

VI. MAX-RELATIVE ENTROPY BOUNDS ON THE ERROR PROBABILITY AND OPTIMAL ERROR EXPONENT

In this section, we provide a lower bound on the minimum error probability of antidistinguishability (Proposition 19), as well as a corresponding upper bound on the optimal error exponent (Theorem 20). To begin with, let us recall the definition of the max-relative entropy of two positive semi-definite operators A and B with $\text{Tr} A \leq 1$ [Dat09]:

$$D_{\max}(A\|B) := \ln \inf_{\lambda \geq 0} \{ \lambda : A \leq \lambda B \}. \quad (180)$$

We have that $D_{\max}(A\|B) = +\infty$ if the support of A is not contained in the support of B . Also, whenever the support of A is contained in the support of B , we have $D_{\max}(A\|B) < +\infty$ and

$$D_{\max}(A\|B) = \ln \left\| B^{-\frac{1}{2}} A B^{-\frac{1}{2}} \right\|_{\infty}, \quad (181)$$

where the inverse is understood to be taken on the support of B . Let us also recall that the minimum error probability of antidistinguishability of an ensemble $\mathcal{E} := \{(\eta_i, \rho_i) : i \in [r]\}$ can also be expressed in terms of the following primal and dual semi-definite programs [BJOP14, Section II] (see, also [YKL75, (III.15)]):

$$\text{Err}(\mathcal{E}) = \inf_{\{M_i\}_{i \in [r]}} \left\{ \sum_{i \in [r]} \eta_i \text{Tr}[M_i \rho_i] : M_i \geq 0 \text{ for all } i \in [r], \sum_{i \in [r]} M_i = \mathbb{I} \right\} \quad (182)$$

$$= \sup_{Y \in \text{Herm}} \{ \text{Tr}[Y] : Y \leq \eta_i \rho_i \text{ for all } i \in [r] \}, \quad (183)$$

where Herm denotes the set of Hermitian operators. The equality holds as a consequence of Slater's condition; indeed we see this by noting that $M_i = I/r$ is strictly feasible for the primal, and $Y = 0$ is feasible for the dual.

Proposition 19. For an ensemble $\mathcal{E} := \{(\eta_i, \rho_i) : i \in [r]\}$, the following lower bound holds:

$$\text{Err}(\mathcal{E}) \geq \exp\left(-\inf_{\omega \in \mathcal{D}} \max_{i \in [r]} D_{\max}(\omega \| \eta_i \rho_i)\right). \quad (184)$$

Proof. By applying (182)–(183) and restricting the optimization over Hermitian operators to be over positive semi-definite operators, we conclude that

$$\text{Err}(\mathcal{E}) \geq \sup_{Y \geq 0} \{\text{Tr}[Y] : Y \leq \eta_i \rho_i, \quad \forall i \in [r]\} \quad (185)$$

$$= \sup_{\lambda \geq 0, \omega \in \mathcal{D}} \{\text{Tr}[\lambda \omega] : \lambda \omega \leq \eta_i \rho_i, \quad \forall i \in [r]\} \quad (186)$$

$$= \sup_{\lambda \geq 0, \omega \in \mathcal{D}} \{\lambda : \lambda \omega \leq \eta_i \rho_i, \quad \forall i \in [r]\} \quad (187)$$

$$\geq \sup_{\lambda > 0, \omega \in \mathcal{D}} \{\lambda : \lambda \omega \leq \eta_i \rho_i, \quad \forall i \in [r]\} \quad (188)$$

$$= \sup_{\lambda > 0, \omega \in \mathcal{D}} \left\{ \lambda : \omega \leq \frac{1}{\lambda} \eta_i \rho_i, \quad \forall i \in [r] \right\} \quad (189)$$

$$= \sup_{\lambda' > 0, \omega \in \mathcal{D}} \left\{ \frac{1}{\lambda'} : \omega \leq \lambda' \eta_i \rho_i, \quad \forall i \in [r] \right\} \quad (190)$$

$$= \left[\inf_{\lambda' > 0, \omega \in \mathcal{D}} \{\lambda' : \omega \leq \lambda' \eta_i \rho_i, \quad \forall i \in [r]\} \right]^{-1} \quad (191)$$

$$= \left[\inf_{\omega \in \mathcal{D}} \exp\left(\max_{i \in [r]} D_{\max}(\omega \| \eta_i \rho_i)\right) \right]^{-1} \quad (192)$$

$$= \left[\exp\left(\inf_{\omega \in \mathcal{D}} \max_{i \in [r]} D_{\max}(\omega \| \eta_i \rho_i)\right) \right]^{-1} \quad (193)$$

$$= \exp\left(-\inf_{\omega \in \mathcal{D}} \max_{i \in [r]} D_{\max}(\omega \| \eta_i \rho_i)\right). \quad (194)$$

The equality (186) follows from the substitution $Y = \lambda \omega$, which is possible because every positive semi-definite operator can be written as a scaled density operator. The equality (190) follows from the substitution $\lambda = \frac{1}{\lambda'}$. ■

Remark 3. Observe that

$$\inf_{\omega \in \mathcal{D}} \max_{i \in [r]} D_{\max}(\omega \| \eta_i \rho_i) = \inf_{\omega \in \mathcal{D}} \max_{\{s_i\}_{i \in [r]}} \sum_{i \in [r]} s_i D_{\max}(\omega \| \eta_i \rho_i) \quad (195)$$

$$= \max_{\{s_i\}_{i \in [r]}} \inf_{\omega \in \mathcal{D}} \sum_{i \in [r]} s_i D_{\max}(\omega \| \eta_i \rho_i), \quad (196)$$

where $\{s_i\}_{i \in [r]}$ is a probability distribution. The first equality follows because the maximum over a finite set can be replaced with a maximum of the expected value of the elements of the set, with the maximum taken over all possible distributions. The second equality follows from an application of the Sion minimax theorem [Sio58]: indeed, the objective function $\sum_{i \in [r]} s_i D_{\max}(\omega \| \eta_i \rho_i)$ is linear and continuous in the probability distribution $\{s_i\}_{i \in [r]}$ and it is lower semi-continuous and quasi-convex in $\omega \in \mathcal{D}$.

Theorem 20. *The following upper bound holds for the optimal error exponent:*

$$E(\rho_1, \dots, \rho_r) \leq \inf_{\omega \in \mathcal{D}} \max_{i \in [r]} D_{\max}(\omega \| \rho_i) \quad (197)$$

$$= \max_{\{s_i\}_{i \in [r]}} \inf_{\omega \in \mathcal{D}} \sum_{i \in [r]} s_i D_{\max}(\omega \| \rho_i). \quad (198)$$

Proof. Consider that

$$E(\rho_1, \dots, \rho_r) = \liminf_{n \rightarrow \infty} -\frac{1}{n} \ln \text{Err}(\mathcal{E}^n) \quad (199)$$

$$\leq \liminf_{n \rightarrow \infty} \frac{1}{n} \inf_{\omega^{(n)} \in \mathcal{D}^{\otimes n}} \max_{i \in [r]} D_{\max}(\omega^{(n)} \| \eta_i \rho_i^{\otimes n}) \quad (200)$$

$$= \liminf_{n \rightarrow \infty} \frac{1}{n} \inf_{\omega^{(n)} \in \mathcal{D}^{\otimes n}} \max_{i \in [r]} D_{\max}(\omega^{(n)} \| \rho_i^{\otimes n}) \quad (201)$$

$$\leq \liminf_{n \rightarrow \infty} \frac{1}{n} \inf_{\omega \in \mathcal{D}} \max_{i \in [r]} D_{\max}(\omega^{\otimes n} \| \rho_i^{\otimes n}) \quad (202)$$

$$= \inf_{\omega \in \mathcal{D}} \max_{i \in [r]} D_{\max}(\omega \| \rho_i). \quad (203)$$

The first inequality follows by applying Proposition 19. The second equality is justified at the end of the proof. The second inequality follows by choosing $\omega^{(n)} = \omega^{\otimes n}$. The final equality follows because the max-relative entropy is additive for tensor-power states. This establishes (197), and (198) follows by the same reasoning as in Remark 3.

We now justify the second equality. Using the fact that, for $c > 0$,

$$D_{\max}(A \| cB) = D_{\max}(A \| B) - \ln c, \quad (204)$$

it follows that

$$\begin{aligned} & \liminf_{n \rightarrow \infty} \frac{1}{n} \inf_{\omega^{(n)} \in \mathcal{D}^{\otimes n}} \max_{i \in [r]} D_{\max}(\omega^{(n)} \| \eta_i \rho_i^{\otimes n}) \\ &= \liminf_{n \rightarrow \infty} \frac{1}{n} \inf_{\omega^{(n)} \in \mathcal{D}^{\otimes n}} \max_{i \in [r]} \left[D_{\max}(\omega^{(n)} \| \rho_i^{\otimes n}) - \ln \eta_i \right] \end{aligned} \quad (205)$$

$$= \liminf_{n \rightarrow \infty} \frac{1}{n} \inf_{\omega^{(n)} \in \mathcal{D}^{\otimes n}} \max_{i \in [r]} D_{\max}(\omega^{(n)} \| \rho_i^{\otimes n}) \quad (206)$$

because

$$D_{\max}(\omega^{(n)} \| \rho_i^{\otimes n}) - \ln \eta_{\min} \geq D_{\max}(\omega^{(n)} \| \rho_i^{\otimes n}) - \ln \eta_i \geq D_{\max}(\omega^{(n)} \| \rho_i^{\otimes n}), \quad (207)$$

where $\eta_{\min} := \min_{i \in [r]} \eta_i$. ■

Remark 4. The upper bound in Theorem 20 bears a resemblance to the following divergence:

$$\max_{\{s_i\}_{i \in [r]}} \inf_{\omega \in \mathcal{D}} \sum_{i \in [r]} s_i D(\omega \| \rho_i) = \max_{\{s_i\}_{i \in [r]}} \left(-\ln \text{Tr} \left[\exp \left(\sum_{i \in [r]} s_i \ln \rho_i \right) \right] \right), \quad (208)$$

where the equality follows whenever each ρ_i is positive definite. Indeed, the only difference between (198) and (208) is the substitution $D_{\max}(\rho \| \sigma) \rightarrow D(\rho \| \sigma) := \text{Tr}[\rho(\ln \rho - \ln \sigma)]$, where

the latter denotes the standard quantum relative entropy [Ume62]. The equality in (208) was established in Eq. (V.121) and Example V.25 of [MBV22]. See Appendix I for a review of the proof of (208). Finally, note that (208) reduces to the multivariate classical Chernoff divergence when the states in the set $\{\rho_i\}_{i \in [r]}$ commute (have a common eigenbasis). As such, this quantity is a multivariate quantum Chernoff divergence according to Definition 14.

VII. CONCLUSION

Summary. We have solved the classical antidistinguishability problem of finding the optimal error exponent, which we proved to be equal to the multivariate classical Chernoff divergence of the given probability measures. To the best of our knowledge, this result constitutes the first operational interpretation of the divergence involving three or more states. We have also given various upper and lower bounds on the optimal error exponent in the quantum case, while it still remains an open problem to compute its exact expression. In analogy with the classical case, we believe that the quantity that gives the exact error exponent in the quantum case should be called *the multivariate quantum Chernoff divergence*.

Future directions. Recall from [BJOP14] that quantum m -state exclusion can be thought of as antidistinguishability of a set of states related to the original set. We leave it as an intriguing open question to determine the optimal asymptotic error exponent for quantum m -state exclusion.

Analogous to the task of antidistinguishing quantum states, one may consider the problem of antidistinguishing an ensemble of quantum channels. In this problem, a quantum channel is chosen randomly from a finite set of quantum channels, with known *a priori* probability distribution. The antidistinguisher is allowed to pass one share of a bipartite quantum state through the channel, after which both the reference system and the channel output system are measured. Based on the measurement outcome, the antidistinguisher's goal is to rule out a quantum channel other than the selected one. It would be an interesting future work to study the asymptotics of the error rates for antidistinguishing an ensemble of quantum channels.

ACKNOWLEDGMENTS

We are especially grateful to Milán Mosonyi for several clarifying discussions about quantum hypothesis testing, as well as to Kaiyuan Ji, Felix Leditzky, Vishal Singh, and Aaron Wagner for insightful discussions. We also thank the anonymous referee and the editor for many helpful comments that improved the manuscript. HKM and MMW acknowledge support from the National Science Foundation under Grant No. 2304816.

DECLARATION OF INTERESTS

On behalf of all authors, the corresponding author states that there is no conflict of interest.

DATA AVAILABILITY STATEMENT

Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

-
- [ACMT⁺07] Koenraad M. R. Audenaert, John Calsamiglia, Ramón Muñoz-Tapia, Emilio Bagan, Ll. Masanes, Antonio Acín, and Frank Verstraete. Discriminating states: The quantum Chernoff bound. *Physical Review Letters*, 98(16):160501, April 2007. [arXiv:quant-ph/0610027](#).
- [And76] T. Ando. Lebesgue-type decomposition of positive operators. *Acta Sci. Math. (Szeged)*, 38(3-4):253–260, January 1976.
- [BC09] Stephen M. Barnett and Sarah Croke. Quantum state discrimination. *Advances in Optics and Photonics*, 1(2):238–278, April 2009. [arXiv:0810.1970](#).
- [BCDvD06] Dave Bacon, Andrew M. Childs, and Wim Dam van Dam. Optimal measurements for the dihedral hidden subgroup problem. *Chicago Journal of Theoretical Computer Science*, 2006:2, October 2006. [arXiv:quant-ph/0501044](#).
- [BCLM14] Jonathan Barrett, Eric G. Cavalcanti, Raymond Lal, and Owen J. E. Maroney. No ψ -epistemic model can fully explain the indistinguishability of quantum states. *Physical Review Letters*, 112(25):250403, June 2014. [arXiv:1310.8302](#).
- [Bil95] Patrick Billingsley. *Probability and Measure*. Wiley Series in Probability and Statistics. Wiley, 1995.
- [BJOP14] Somshubhro Bandyopadhyay, Rahul Jain, Jonathan Oppenheim, and Christopher Perry. Conclusive exclusion of quantum states. *Physical Review A*, 89(2):022336, February 2014. [arXiv:1306.4683](#).
- [BK15] Joonwoo Bae and Leong-Chuan Kwek. Quantum state discrimination and its applications. *Journal of Physics A: Mathematical and Theoretical*, 48(8):083001, January 2015. [arXiv:1707.02571](#).
- [BL06] J. Borwein and A. Lewis. *Convex Analysis*. Springer, New York, 2006.
- [Bor26] Max Born. Quantum mechanics of collision processes. *Uspekhi Fizich*, June 1926.
- [CFS02] Carlton M. Caves, Christopher A. Fuchs, and Rüdiger Schack. Conditions for compatibility of quantum-state assignments. *Physical Review A*, 66(6):062111, December 2002. [arXiv:quant-ph/0206110](#).
- [Che52] Herman Chernoff. A measure of asymptotic efficiency for tests of a hypothesis based on the sum of observations. *The Annals of Mathematical Statistics*, 23(4):493–507, December 1952.
- [Das11] Anirban DasGupta. *Probability for Statistics and Machine Learning: Fundamentals and Advanced Topics*. Springer, 2011.
- [Dat09] Nilanjana Datta. Min- and max-relative entropies and a new entanglement monotone. *IEEE Transactions on Information Theory*, 55:2816–2826, 2009. [arXiv:0803.2770](#).
- [DL14] Nilanjana Datta and Felix Leditzky. A limit of the quantum Rényi divergence. *Journal of Physics A: Mathematical and Theoretical*, 47(4):045304, 2014.
- [Fek23] Michael Fekete. über die verteilung der wurzeln bei gewissen algebraischen gleichungen mit ganzzahligen koeffizienten. *Mathematische Zeitschrift*, 17(1):228–249, December 1923.
- [FL96] I. Fazekas and F. Liese. Some properties of the Hellinger transform and its application in classification problems. *Computers & Mathematics with Applications*, 31(8):107–116, April 1996.
- [FLO23] Keiichiro Furuya, Nima Lashkari, and Shoy Ouseph. Monotonic multi-state quantum f-divergences. *Journal of Mathematical Physics*, 64(4):042203, April 2023.
- [Gri93] B. Grigelionis. *On Hellinger transforms for solutions of martingale problems*, pages 107–116. Springer New York, 1993.
- [HB20] Vojtěch Havlíček and Jonathan Barrett. Simple communication complexity separation from quantum state antidistinguishability. *Physical Review Research*, 2(1):013326, March 2020. [arXiv:1911.01927](#).
- [Hel09] Ernst Hellinger. Neue begründung der theorie quadratischer formen von unendlichvielen veränderlichen. *Journal für die reine und angewandte Mathematik*, 1909(136):210–271, July 1909.
- [Hel69] Carl W. Helstrom. Quantum detection and estimation theory. *Journal of Statistical Physics*, 1:231–252, June 1969.
- [Hia94] Fumio Hiai. Equality cases in matrix norm inequalities of Golden-Thompson type. *Linear and Multilinear Algebra*, 36(4):239–249, April 1994.

- [HK18] Teiko Heinosaari and Oskari Kerppo. Antidistinguishability of pure quantum states. *Journal of Physics A: Mathematical and Theoretical*, 51(36):365303, July 2018. [arXiv:1804.10457](https://arxiv.org/abs/1804.10457).
- [HM17] Fumio Hiai and Milán Mosonyi. Different quantum f -divergences and the reversibility of quantum operations. *Reviews in Mathematical Physics*, 29(07):1750023, August 2017.
- [Hol72] Alexander S. Holevo. An analogue of statistical decision theory and noncommutative probability theory. *Trudy Moskovskogo Matematicheskogo Obshchestva*, 26:133–149, 1972.
- [HSR03] Michael Horodecki, Peter W. Shor, and Mary Beth Ruskai. Entanglement breaking channels. *Reviews in Mathematical Physics*, 15(06):629–641, 2003.
- [HT16] Masahito Hayashi and Marco Tomamichel. Correlation detection and an operational interpretation of the Rényi mutual information. *Journal of Mathematical Physics*, 57(10):102201, October 2016.
- [Jac89] Jean Jacod. Filtered statistical models and Hellinger processes. *Stochastic Processes and their Applications*, 32(1):3–45, June 1989.
- [LD20] Matthew Leifer and Cristhiano Duarte. Noncontextuality inequalities from antidistinguishability. *Physical Review A*, 101(6):062113, June 2020. [arXiv:2001.11485](https://arxiv.org/abs/2001.11485).
- [LeC70] L. LeCam. On the assumptions used to prove asymptotic normality of maximum likelihood estimates. *The Annals of Mathematical Statistics*, 41(3):802–828, June 1970.
- [Lei14] Matthew Saul Leifer. Is the quantum state real? An extended review of ψ -ontology theorems. *Quanta*, 3:67–155, November 2014. [arXiv:1409.1570](https://arxiv.org/abs/1409.1570).
- [Li16] Ke Li. Discriminating quantum states: The multiple Chernoff distance. *The Annals of Statistics*, 44(4):1661–1679, August 2016. [arXiv:1508.06624](https://arxiv.org/abs/1508.06624).
- [LJ97] C. C. Leang and D. H. Johnson. On the asymptotics of m -hypothesis Bayesian detection. *IEEE Transactions on Information Theory*, 43(1):280–282, January 1997.
- [LM10] F. Liese and Klaus-J. Miescke. *Statistical Decision Theory*. Springer New York, NY, 2010.
- [LR73] Elliott H. Lieb and Mary Beth Ruskai. Proof of the strong subadditivity of quantum-mechanical entropy. *Journal of Mathematical Physics*, 14(12):1938–1941, December 1973.
- [Mat13] Keiji Matsumoto. A new quantum version of f -divergence, 2013. [arXiv:1311.4722](https://arxiv.org/abs/1311.4722).
- [Mat14] Keiji Matsumoto. On maximization of measured f -divergence between a given pair of quantum states, 2014. [arXiv:1412.3676](https://arxiv.org/abs/1412.3676).
- [Mat18] Keiji Matsumoto. A new quantum version of f -divergence. In Masanao Ozawa, Jeremy Butterfield, Hans Halvorson, Miklós Rédei, Yuichiro Kitajima, and Francesco Buscemi, editors, *Reality and Measurement in Algebraic Quantum Theory*, volume 261, pages 229–273, Singapore, 2018. Springer Singapore. Series Title: Springer Proceedings in Mathematics & Statistics.
- [MBV22] Milán Mosonyi, Gergely Bunnth, and Péter Vrana. Geometric relative entropies and barycentric Rényi divergences. [arXiv:2207.14282v2](https://arxiv.org/abs/2207.14282v2), July 2022.
- [MO15] Milán Mosonyi and Tomohiro Ogawa. Quantum hypothesis testing and the operational interpretation of the quantum Rényi relative entropies. *Communications in Mathematical Physics*, 334:1617–1648, 2015.
- [NS09] Michael Nussbaum and Arleta Szkoła. The Chernoff lower bound for symmetric quantum hypothesis testing. *The Annals of Statistics*, 37(2):1040–1057, April 2009. [arXiv:quant-ph/0607216](https://arxiv.org/abs/quant-ph/0607216).
- [PBR12] Matthew F. Pusey, Jonathan Barrett, and Terry Rudolph. On the reality of the quantum state. *Nature Physics*, 8(6):475–478, June 2012. [arXiv:1111.3328](https://arxiv.org/abs/1111.3328).
- [RS23] Vincent Russo and Jamie Sikora. Inner products of pure states and their antidistinguishability. *Physical Review A*, 107(3):L030202, March 2023. [arXiv:2206.08313](https://arxiv.org/abs/2206.08313).
- [Rus02] Mary Beth Ruskai. Inequalities for quantum entropy: A review with conditions for equality. *Journal of Mathematical Physics*, 43(9):4358–4375, 2002.
- [Sal73] N. P. Salikhov. Asymptotic properties of error probabilities of tests for distinguishing between several multinomial testing schemes. *Doklady Akademii Nauk*, 209(1):54–57, 1973.
- [Sal99] N. P. Salikhov. On one generalization of Chernov’s distance. *Theory of Probability & Its Applications*, 43(2):239–255, 1999.
- [Sal03] N. P. Salikhov. Optimal sequences of tests for several polynomial schemes of trials. *Theory of Probability & Its Applications*, 47(2):286–298, 2003.
- [Sch12] Mark J. Schervish. *Theory of Statistics*. Springer Science & Business Media, 2012.
- [Shi16] Albert N. Shiryaev. *Probability-1*, volume 95. Springer, 2016.
- [Sio58] Maurice Sion. On general minimax theorems. *Pacific Journal of Mathematics*, 8:171–176, 1958.

- [Str11] Helmut Strasser. *Mathematical Theory of Statistics: Statistical Experiments and Asymptotic Decision Theory*, volume 7. Walter de Gruyter, 2011.
- [Tor81] Erik N. Torgersen. Measures of information based on comparison with total information and with total ignorance. *The Annals of Statistics*, 9(3):638–657, 1981.
- [Tor91] Erik N. Torgersen. *Comparison of Statistical Experiments*, volume 36. Cambridge University Press, 1991.
- [Ume62] Hisaharu Umegaki. Conditional expectation in an operator algebra, IV (Entropy and information). *Kodai Mathematical Seminar Reports*, 14:59–85, 1962.
- [Wil17] Mark M. Wilde. *Quantum Information Theory*. Cambridge University Press, second edition, 2017.
- [YKL75] H. Yuen, R. Kennedy, and M. Lax. Optimum testing of multiple hypotheses in quantum detection theory. *IEEE Transactions on Information Theory*, 21(2):125–134, March 1975.

Appendix A: Expectation values at non-corner points

We begin by stating a known property of convex functions in the lemma below. We include a proof of the statement for the sake of completeness.

Lemma 21. *Let $a > 0$ be arbitrary. Let $f : [0, a] \rightarrow \mathbb{R}$ be a convex and continuous function on $[0, a]$, and suppose f is differentiable on $(0, a)$. Then the one-sided derivative*

$$f'_+(0) := \lim_{t \searrow 0} \frac{f(t) - f(0)}{t} \quad (\text{A1})$$

exists and fulfills

$$f'_+(0) = \lim_{t \searrow 0} f'(t). \quad (\text{A2})$$

Here $f'_+(0)$ is either finite or takes the value $-\infty$; if f takes its minimum value at 0, then $f'_+(0)$ is finite and $f'_+(0) \geq 0$.

Proof. The map $t \mapsto (f(t) - f(0))/t$ defined on $(0, a)$ is non-decreasing. See [BL06, Section 2.1, Exercise 7]). Also, the limit in (A1) exists in $\mathbb{R} \cup \{-\infty\}$ [BL06, Proposition 3.1.2]. By the Lagrange mean-value theorem, for any $t \in (0, a)$ there exists $u_t \in (0, t)$ such that

$$\frac{f(t) - f(0)}{t} = f'(u_t). \quad (\text{A3})$$

This implies

$$f'_+(0) = \lim_{t \searrow 0} \frac{f(t) - f(0)}{t} = \lim_{t \searrow 0} f'(u_t). \quad (\text{A4})$$

The derivative of f is a non-decreasing function on $(0, a)$. Indeed, let $t_1, t_2 \in (0, a)$ such that $t_1 < t_2$. Let $h \in (t_1, t_2)$ be arbitrary. We have

$$h = \left(\frac{h - t_1}{t_2 - t_1} \right) t_2 + \left(1 - \frac{h - t_1}{t_2 - t_1} \right) t_1. \quad (\text{A5})$$

By convexity of f , we get

$$f(h) \leq \left(\frac{h - t_1}{t_2 - t_1} \right) f(t_2) + \left(1 - \frac{h - t_1}{t_2 - t_1} \right) f(t_1), \quad (\text{A6})$$

which implies

$$\frac{f(h) - f(t_1)}{h - t_1} \leq \frac{f(t_2) - f(t_1)}{t_2 - t_1}. \quad (\text{A7})$$

By taking the limit $h \searrow t_1$ in the above inequality, we get

$$f'(t_1) \leq \frac{f(t_2) - f(t_1)}{t_2 - t_1}. \quad (\text{A8})$$

A similar argument shows that

$$\frac{f(t_2) - f(t_1)}{t_2 - t_1} \leq f'(t_2). \quad (\text{A9})$$

By (A8) and (A9), we thus get $f'(t_1) \leq f'(t_2)$ and hence f' is non-decreasing on $(0, a)$. From (A4) and the fact that f' is non-decreasing, we thus get

$$f'_+(0) = \lim_{t \searrow 0} f'(t), \quad (\text{A10})$$

with a possible value $-\infty$. If f is minimized at 0, then we have $f(t) - f(0) \geq 0$ for all $t \in (0, a)$. It then directly follows from the definition (A1) that $f'_+(0) \geq 0$. ■

Recall that \mathbb{T}_r^1 is the set of non-corner points of \mathbb{T}_r given by (66). Let $\mathbf{t} \in \mathbb{T}_r^1$. Define a set

$$B_{\mathbf{t}} := \{i \in [r-1] : t_i > 0\}, \quad (\text{A11})$$

and let $B_{\mathbf{t}}^c := [r-1] \setminus B_{\mathbf{t}}$. Let β denote the cardinality of the set $B_{\mathbf{t}}$ so that $1 \leq \beta \leq r-1$. We emphasize that \mathbf{t} corresponds to an interior point of $\mathbb{T}_{\beta+1}$, which is the β -vector obtained by discarding the zero entries of \mathbf{t} . This allows us to use the properties of exponential family of densities given in (59). So, by the similar arguments as given for (65), it follows that for any $i \in B_{\mathbf{t}}$, the expectation value $\mathbb{E}_{\mathbf{t}}[q_i]$ exists, and it satisfies $\partial_i K(\mathbf{t}) = \mathbb{E}_{\mathbf{t}}[q_i]$. It remains to show for $i \in B_{\mathbf{t}}^c$ that $\mathbb{E}_{\mathbf{t}}[q_i]$ exists, and it is equal to $\partial_i^+ K(\mathbf{t})$. Let us fix an arbitrary index $i \in B_{\mathbf{t}}^c$. Choose a small number $\varepsilon > 0$ such that $\mathbf{t} + h\mathbf{e}_i \in \mathbb{T}_r^1$ for all $h \in [0, \varepsilon]$. The function $h \mapsto K(\mathbf{t} + h\mathbf{e}_i)$ is continuous, convex on $[0, \varepsilon]$, and it is differentiable on $(0, \varepsilon)$. Lemma 21 thus implies that

$$\partial_i^+ K(\mathbf{t}) = \lim_{h \searrow 0} \partial_i K(\mathbf{t} + h\mathbf{e}_i) = \lim_{h \searrow 0} \mathbb{E}_{\mathbf{t} + h\mathbf{e}_i}[q_i]. \quad (\text{A12})$$

Here we used the relation $\partial_i K(\mathbf{t} + h\mathbf{e}_i) = \mathbb{E}_{\mathbf{t} + h\mathbf{e}_i}[q_i]$ proved earlier. We now claim that $\mathbb{E}_{\mathbf{t}}[q_i]$ exists and satisfies

$$\lim_{h \searrow 0} \mathbb{E}_{\mathbf{t} + h\mathbf{e}_i}[q_i] = \mathbb{E}_{\mathbf{t}}[q_i] \quad (\text{A13})$$

with a possible value $-\infty$. Indeed, we have

$$\mathbb{E}_{\mathbf{t} + h\mathbf{e}_i}[q_i] = \frac{1}{H(\mathbf{t} + h\mathbf{e}_i)} \int \mathbf{d}\mu \, q_i p_r \exp\left(\sum_{j \in [r-1]} t_j q_j + h q_i\right). \quad (\text{A14})$$

By continuity of H , we have $H(\mathbf{t} + h\mathbf{e}_i) \rightarrow H(\mathbf{t})$ as $h \searrow 0$. Thus, for (A13) to hold, it suffices to prove

$$\lim_{h \searrow 0} \int \mathbf{d}\mu \, q_i p_r \exp\left(\sum_{j \in [r-1]} t_j q_j + h q_i\right) = \int \mathbf{d}\mu \, q_i p_r \exp\left(\sum_{j \in [r-1]} t_j q_j\right). \quad (\text{A15})$$

Let $q_i = q_i^+ - q_i^-$, where q_i^+ and q_i^- are non-negative functions with mutually disjoint supports. This gives

$$\begin{aligned} & \int \mathbf{d}\mu q_i p_r \exp\left(\sum_{j \in [r-1]} t_j q_j + h q_i\right) \\ &= \int \mathbf{d}\mu q_i^+ p_r \exp\left(\sum_{j \in [r-1]} t_j q_j + h q_i^+\right) - \int \mathbf{d}\mu q_i^- p_r \exp\left(\sum_{j \in [r-1]} t_j q_j - h q_i^-\right). \end{aligned} \quad (\text{A16})$$

The first integral term in the right-hand side of (A16) is finite. Indeed, since $h \in (0, \varepsilon)$, we have

$$q_i^+ p_r \exp\left(\sum_{j \in [r-1]} t_j q_j + h q_i^+\right) \leq q_i^+ p_r \exp\left(\sum_{j \in [r-1]} t_j q_j + \varepsilon q_i^+\right). \quad (\text{A17})$$

The function in the right-hand side of (A17) is μ -integrable, because $\mathbf{t} + \varepsilon \mathbf{e}_i$ corresponds to an interior point of $\mathbb{T}_{r-\beta+1}$ so that the properties of an exponential family of densities apply. Moreover, we have pointwise convergence

$$q_i^+ p_r \exp\left(\sum_{j \in [r-1]} t_j q_j + h q_i^+\right) \rightarrow q_i^+ p_r \exp\left(\sum_{j \in [r-1]} t_j q_j\right) \quad \text{as } h \searrow 0. \quad (\text{A18})$$

By the Lebesgue dominated convergence theorem, we have

$$\lim_{h \searrow 0} \int \mathbf{d}\mu q_i^+ p_r \exp\left(\sum_{j \in [r-1]} t_j q_j + h q_i^+\right) = \int \mathbf{d}\mu q_i^+ p_r \exp\left(\sum_{j \in [r-1]} t_j q_j\right) < \infty. \quad (\text{A19})$$

We now consider the second integral term in the right-hand side of (A16). We have the pointwise monotone convergence

$$q_i^- p_r \exp\left(\sum_{j \in [r-1]} t_j q_j - h q_i^-\right) \nearrow q_i^- p_r \exp\left(\sum_{j \in [r-1]} t_j q_j\right), \quad \text{as } h \searrow 0. \quad (\text{A20})$$

By the monotone convergence theorem, we get

$$\lim_{h \searrow 0} \int \mathbf{d}\mu q_i^- p_r \exp\left(\sum_{j \in [r-1]} t_j q_j - h q_i^-\right) = \int \mathbf{d}\mu q_i^- p_r \exp\left(\sum_{j \in [r-1]} t_j q_j\right) \quad (\text{A21})$$

regardless of whether the right-hand integral in (A21) is finite or infinite. The latter point is explicitly stressed in Theorem 16.2 of [Bil95]. By taking the limit $h \searrow 0$ in (A16), and then using (A12), (A19) and (A21), we get

$$\partial_i^+ K(\mathbf{t}) = \mathbb{E}_{\mathbf{t}}[q_i^+] - \mathbb{E}_{\mathbf{t}}[q_i^-] = \mathbb{E}_{\mathbf{t}}[q_i]. \quad (\text{A22})$$

Since $\mathbb{E}_{\mathbf{t}}[q_i^+]$ is a real number, $\mathbb{E}_{\mathbf{t}}[q_i]$ takes a value in $\mathbb{R} \cup \{-\infty\}$. If \mathbf{t} is a minimizer of K then by Lemma 21 we have $\partial_i^+ K(\mathbf{t}) \geq 0$, and hence $\mathbb{E}_{\mathbf{t}}[q_i]$ is finite. We have thus accomplished that if $\mathbf{t} \in \mathbb{T}_r^1$ is a minimizer of K and $i \in [r-1]$, then the expectation value $\mathbb{E}_{\mathbf{t}}[q_i]$ exists, is finite, and satisfies $\partial_i^+ K(\mathbf{t}) = \mathbb{E}_{\mathbf{t}}[q_i]$.

Appendix B: Proof of Theorem 7

One of the major differences between the special case of mutually absolutely continuous probability measures and the general case of probability measures with unequal supports is that, the Hellinger transform is continuous on the entire unit simplex in the special case, but only in the interior of the unit simplex in the general case. As it turns out, in the definition (23), restricting the infimum to be over the interior of the unit simplex does not change the value of the multivariate classical Chernoff divergence.

Define

$$\xi_{\text{cl}}^{\circ}(P_1, \dots, P_r) := -\ln \inf_{\mathbf{s} \in \mathbb{S}_r^{\circ}} \mathbb{H}_{\mathbf{s}}(P_1, \dots, P_r), \quad (\text{B1})$$

where \mathbb{S}_r° denotes the interior of the unit simplex \mathbb{S}_r . We will show that $\xi_{\text{cl}}^{\circ}(P_1, \dots, P_r)$ coincides with the multivariate classical Chernoff divergence $\xi_{\text{cl}}(P_1, \dots, P_r)$ defined in (23), and also that it is the optimal error exponent for antidistinguishing the given probability measures.

As a first step towards showing $\xi_{\text{cl}}^{\circ}(P_1, \dots, P_r) = \xi_{\text{cl}}(P_1, \dots, P_r)$, we discuss a continuous extension of the Hellinger transform defined in the interior of the unit simplex to its boundary. Recall that μ is the dominating measure given by (2) and p_1, \dots, p_r are the induced densities defined in (3). Define

$$D := \{\omega \in \Omega : p_i(\omega) > 0, \forall i \in [r]\}. \quad (\text{B2})$$

For any $\mathbf{s} := (s_1, \dots, s_r) \in \mathbb{S}_r$, define

$$\mathbb{H}_{\mathbf{s}}^{-}(P_1, \dots, P_r) := \int_D \mathbf{d}\mu p_1^{s_1} \cdots p_r^{s_r}. \quad (\text{B3})$$

We note that $\mathbb{H}_{\mathbf{s}}^{-}(P_1, \dots, P_r) = \mathbb{H}_{\mathbf{s}}(P_1, \dots, P_r)$ for all $\mathbf{s} \in \mathbb{S}_r^{\circ}$, which implies that the map $\mathbf{s} \mapsto \mathbb{H}_{\mathbf{s}}^{-}(P_1, \dots, P_r)$ is continuous in the interior of the unit simplex \mathbb{S}_r . Also, its continuity on the boundary of the unit simplex follows by the Lebesgue dominated convergence theorem. It is easy to see that $\mathbb{H}_{\mathbf{s}}^{-}(P_1, \dots, P_r) \leq \mathbb{H}_{\mathbf{s}}(P_1, \dots, P_r)$ for all $\mathbf{s} \in \mathbb{S}_r$. Indeed, let $\mathbf{s} := (s_1, \dots, s_r) \in \mathbb{S}_r$ be arbitrary, set $J := \{j \in [r] : s_j > 0\}$ and $D_J := \{\omega \in \Omega : p_j(\omega) > 0, \forall j \in J\}$. Since $D \subset D_J$, we get

$$\mathbb{H}_{\mathbf{s}}^{-}(P_1, \dots, P_r) = \int_D \mathbf{d}\mu \prod_{j \in J} p_j^{s_j} \quad (\text{B4})$$

$$\leq \int_{D_J} \mathbf{d}\mu \prod_{j \in J} p_j^{s_j} \quad (\text{B5})$$

$$= \mathbb{H}_{\mathbf{s}}(P_1, \dots, P_r). \quad (\text{B6})$$

It should be noted that the last equality follows from the convention $0^0 = 1$ in the definition (19) of the Hellinger transform. We thus get from (B1) that

$$\xi_{\text{cl}}^{\circ}(P_1, \dots, P_r) = -\ln \inf_{\mathbf{s} \in \mathbb{S}_r^{\circ}} \mathbb{H}_{\mathbf{s}}^{-}(P_1, \dots, P_r) \quad (\text{B7})$$

$$= -\ln \inf_{\mathbf{s} \in \mathbb{S}_r} \mathbb{H}_{\mathbf{s}}^{-}(P_1, \dots, P_r) \quad (\text{B8})$$

$$= -\ln \inf_{\mathbf{s} \in \mathbb{S}_r} \mathbb{H}_{\mathbf{s}}(P_1, \dots, P_r) \quad (\text{B9})$$

$$= \xi_{\text{cl}}(P_1, \dots, P_r), \quad (\text{B10})$$

as was claimed earlier.

For an achievable bound on the optimal error exponent, we follow the same arguments as in the development (39)–(48) to get

$$\text{Err}_{\text{cl}}(\mathcal{E}_{\text{cl}}^n) \leq \mathbb{H}_{\mathbf{s}}(P_1, \dots, P_r)^n, \quad \text{for all } \mathbf{s} \in \mathbb{S}_r^\circ. \quad (\text{B11})$$

This gives

$$\liminf_{n \rightarrow \infty} -\frac{1}{n} \ln \text{Err}_{\text{cl}}(\mathcal{E}_{\text{cl}}^n) \geq -\ln \inf_{\mathbf{s} \in \mathbb{S}_r^\circ} \mathbb{H}_{\mathbf{s}}(P_1, \dots, P_r) = \xi_{\text{cl}}^\circ(P_1, \dots, P_r) = \xi_{\text{cl}}(P_1, \dots, P_r). \quad (\text{B12})$$

We now present an optimality bound on the optimal error exponent. The case $\mu(D) = 0$ is trivial, since the antidistinguishability error probability is equal to zero in this case. So, we assume $\mu(D) > 0$. Define

$$\alpha_i := P_i(D) > 0, \quad i \in [r] \quad (\text{B13})$$

and *conditional probability densities* on the set D as

$$\tilde{p}_i := \frac{p_i}{\alpha_i}, \quad i \in [r]. \quad (\text{B14})$$

Observe that the conditional probability densities $\tilde{p}_1, \dots, \tilde{p}_r$ correspond to mutually absolutely continuous probability measures $P_1/\alpha_1, \dots, P_r/\alpha_r$ on the set D . Let us define for $\mathbf{t} := (t_1, \dots, t_{r-1}) \in \mathbb{T}_r$ and $\mathbf{s}_{\mathbf{t}} = (t_1, \dots, t_{r-1}, 1 - \sum_{i \in [r-1]} t_i) \in \mathbb{S}_r$,

$$\text{H}^-(\mathbf{t}) := \mathbb{H}_{\mathbf{s}_{\mathbf{t}}}^-(P_1, \dots, P_r), \quad (\text{B15})$$

$$\text{K}^-(\mathbf{t}) := \ln \text{H}^-(\mathbf{t}). \quad (\text{B16})$$

From (B14) we get,

$$\text{H}^-(\mathbf{t}) = \int_D \mathbf{d}\mu p_1^{t_1} \cdots p_{r-1}^{t_{r-1}} p_r^{1 - \sum_{i \in [r-1]} t_i} \quad (\text{B17})$$

$$= \alpha_1^{t_1} \cdots \alpha_{r-1}^{t_{r-1}} \alpha_r^{1 - \sum_{i \in [r-1]} t_i} \int_D \mathbf{d}\mu \tilde{p}_1^{t_1} \cdots \tilde{p}_{r-1}^{t_{r-1}} \tilde{p}_r^{1 - \sum_{i \in [r-1]} t_i}. \quad (\text{B18})$$

Define

$$\tilde{\text{H}}(\mathbf{t}) := \int_D \mathbf{d}\mu \tilde{p}_1^{t_1} \cdots \tilde{p}_{r-1}^{t_{r-1}} \tilde{p}_r^{1 - \sum_{i \in [r-1]} t_i}, \quad (\text{B19})$$

$$\tilde{\text{K}}(\mathbf{t}) := \ln \tilde{\text{H}}(\mathbf{t}). \quad (\text{B20})$$

From the above development (B15)–(B20), we get

$$\text{K}^-(\mathbf{t}) = \tilde{\text{K}}(\mathbf{t}) + \text{L}(\mathbf{t}), \quad \text{for all } \mathbf{t} \in \mathbb{T}_r, \quad (\text{B21})$$

where L is a linear map defined on \mathbb{T}_r as

$$\text{L}(\mathbf{t}) := \ln \alpha_r + \sum_{i \in [r-1]} t_i \ln \left(\frac{\alpha_i}{\alpha_r} \right). \quad (\text{B22})$$

Analogous to (59), we define an exponential family of densities on D as

$$\tilde{p}_{\mathbf{t}} := \frac{1}{\tilde{\text{H}}(\mathbf{t})} \tilde{p}_1^{t_1} \cdots \tilde{p}_{r-1}^{t_{r-1}} \tilde{p}_r^{1 - \sum_{i \in [r-1]} t_i}. \quad (\text{B23})$$

Recall from the definition of the antidistinguishability error probability (9) that we only need to work with the probability densities p_1, \dots, p_r restricted to the set D ; i.e., we have for all positive integers n ,

$$\text{Err}_{\text{cl}}(\mathcal{E}_{\text{cl}}^n) = \int_D \mathbf{d}\mu^{\otimes n} (\eta_1 p_1^{\otimes n} \wedge \dots \wedge \eta_r p_r^{\otimes n}). \quad (\text{B24})$$

Define

$$\tilde{\mathbb{T}}_{r,f}^1 := \{\mathbf{t} \in \mathbb{T}_r^1 : \partial_i^+ \tilde{\mathbf{K}}(\mathbf{t}) \neq -\infty, \forall i \in [r-1]\} \quad (\text{B25})$$

analogous to (69). Let $\omega^n := (\omega_1, \dots, \omega_n) \in D^n$ and $\mathbf{t} \in \tilde{\mathbb{T}}_{r,f}^1$ be arbitrary. We have that

$$p_i^{\otimes n}(\omega^n) = \exp\left(n \tilde{G}_{\mathbf{t},n}^{(i)}(\omega^n)\right) \tilde{p}_{\mathbf{t}}^{\otimes n}(\omega^n), \quad (\text{B26})$$

where

$$\tilde{G}_{\mathbf{t},n}^{(i)}(\omega^n) := \frac{1}{n} \sum_{j \in [n]} \ln \frac{\tilde{p}_i}{\tilde{p}_{\mathbf{t}}}(\omega_j) + \ln \alpha_i \quad \text{for } i \in [r]. \quad (\text{B27})$$

By following similar arguments as given in the development (78)–(91), we get

$$\limsup_{n \rightarrow \infty} -\frac{1}{n} \ln \text{Err}_{\text{cl}}(\mathcal{E}_{\text{cl}}^n) \leq -\min_{1 \leq i \leq r} (\tilde{\gamma}_i(\mathbf{t}) + \ln \alpha_i), \quad (\text{B28})$$

such that, similar to (77),

$$\tilde{\gamma}_i(\mathbf{t}) = \begin{cases} \partial_i^+ \tilde{\mathbf{K}}(\mathbf{t}) - \mathbf{t}^T \nabla^+ \tilde{\mathbf{K}}(\mathbf{t}) + \tilde{\mathbf{K}}(\mathbf{t}), & i \in [r-1], \\ -\mathbf{t}^T \nabla^+ \tilde{\mathbf{K}}(\mathbf{t}) + \tilde{\mathbf{K}}(\mathbf{t}), & i = r, \end{cases} \quad \text{for } \mathbf{t} \in \tilde{\mathbb{T}}_{r,f}^1. \quad (\text{B29})$$

From (B21), we have for $\mathbf{t} \in \mathbb{T}_r^1$,

$$\partial_i^+ \tilde{\mathbf{K}}(\mathbf{t}) = \partial_i^+ \mathbf{K}^-(\mathbf{t}) - \ln\left(\frac{\alpha_i}{\alpha_r}\right) \quad (\text{B30})$$

and hence

$$\mathbf{t}^T \nabla^+ \tilde{\mathbf{K}}(\mathbf{t}) = \mathbf{t}^T \nabla^+ \mathbf{K}^-(\mathbf{t}) - \mathbf{L}(\mathbf{t}) + \ln \alpha_r. \quad (\text{B31})$$

Substituting the relations (B30) and (B31) in (B29), we obtain

$$\tilde{\gamma}_i(\mathbf{t}) + \ln \alpha_i = \begin{cases} \partial_i^+ \mathbf{K}^-(\mathbf{t}) - \mathbf{t}^T \nabla^+ \mathbf{K}^-(\mathbf{t}) + \mathbf{K}^-(\mathbf{t}), & i \in [r-1], \\ -\mathbf{t}^T \nabla^+ \mathbf{K}^-(\mathbf{t}) + \mathbf{K}^-(\mathbf{t}), & i = r, \end{cases} \quad \text{for } \mathbf{t} \in \tilde{\mathbb{T}}_{r,f}^1. \quad (\text{B32})$$

As the function \mathbf{L} in (B21) is linear and $\tilde{\mathbf{K}}$ is convex on \mathbb{T}_r , we conclude that \mathbf{K}^- is convex as well, and it inherits the smoothness properties of $\tilde{\mathbf{K}}$. One can now essentially follow the steps described in the development (93)–(97) to show that there is no loss in assuming that there exists a minimizer $\mathbf{t}^* \in \tilde{\mathbb{T}}_{r,f}^1$ of \mathbf{K}^- , and that it satisfies

$$-\min_{1 \leq i \leq r} (\tilde{\gamma}_i(\mathbf{t}^*) + \ln \alpha_i) \leq -\mathbf{K}^-(\mathbf{t}^*). \quad (\text{B33})$$

Combining (B28) and (B33), we thus get

$$\limsup_{n \rightarrow \infty} -\frac{1}{n} \ln \text{Err}_{\text{cl}}(\mathcal{E}_{\text{cl}}^n) \leq \sup_{\mathbf{t} \in \mathbb{T}_r} -\mathbf{K}^-(\mathbf{t}) = \xi_{\text{cl}}^\circ(P_1, \dots, P_r) = \xi_{\text{cl}}(P_1, \dots, P_r). \quad (\text{B34})$$

This completes the optimality part. It follows from (B12) and (B34) that the limit $\lim_{n \rightarrow \infty} -\frac{1}{n} \ln \text{Err}_{\text{cl}}(\mathcal{E}_{\text{cl}}^n)$ exists and the desired equality (98) holds.

Appendix C: Proof of Equation (139)

Proposition 22. For arbitrary (not necessarily normalized) vectors $|\varphi\rangle, |\zeta\rangle \in \mathcal{H}$, the following equality holds:

$$\| |\varphi\rangle\langle\varphi| - |\zeta\rangle\langle\zeta| \|_1^2 = (\langle\varphi|\varphi\rangle + \langle\zeta|\zeta\rangle)^2 - 4|\langle\zeta|\varphi\rangle|^2. \quad (\text{C1})$$

Proof. The equality (C1) trivially holds if one of the vectors is zero. So, we assume that both $|\varphi\rangle$ and $|\zeta\rangle$ are non-zero vectors. Define

$$|\varphi'\rangle := \frac{|\varphi\rangle}{\| |\varphi\rangle \|}, \quad |\zeta'\rangle := \frac{|\zeta\rangle}{\| |\zeta\rangle \|}. \quad (\text{C2})$$

Then the desired equality is equivalent to

$$\| c|\varphi'\rangle\langle\varphi'| - d|\zeta'\rangle\langle\zeta'| \|_1^2 = (c + d)^2 - 4cd |\langle\zeta'|\varphi'\rangle|^2, \quad (\text{C3})$$

where

$$c := \| |\varphi\rangle \|^2, \quad d := \| |\zeta\rangle \|^2. \quad (\text{C4})$$

Defining $|\varphi^\perp\rangle$ to be the unit vector orthogonal to $|\varphi'\rangle$ in $\text{span}\{|\varphi'\rangle, |\zeta'\rangle\}$, we find that

$$|\zeta'\rangle = \cos(\theta)|\varphi'\rangle + \sin(\theta)|\varphi^\perp\rangle, \quad (\text{C5})$$

where

$$\cos(\theta) = \langle\varphi'|\zeta'\rangle. \quad (\text{C6})$$

Then it follows that

$$\begin{aligned} & c|\varphi'\rangle\langle\varphi'| - d|\zeta'\rangle\langle\zeta'| \\ &= c|\varphi'\rangle\langle\varphi'| - d(\cos(\theta)|\varphi'\rangle + \sin(\theta)|\varphi^\perp\rangle)(\cos(\theta)\langle\varphi'| + \sin(\theta)\langle\varphi^\perp|) \end{aligned} \quad (\text{C7})$$

$$\begin{aligned} &= [c - d\cos^2(\theta)]|\varphi'\rangle\langle\varphi'| - d\sin(\theta)\cos(\theta)|\varphi^\perp\rangle\langle\varphi'| \\ &\quad - d\sin(\theta)\cos(\theta)|\varphi'\rangle\langle\varphi^\perp| - d\sin^2(\theta)|\varphi^\perp\rangle\langle\varphi^\perp|. \end{aligned} \quad (\text{C8})$$

As a matrix with respect to the basis $\{|\varphi'\rangle, |\varphi^\perp\rangle\}$, the last line has the following form:

$$\begin{bmatrix} c - d\cos^2(\theta) & -d\sin(\theta)\cos(\theta) \\ -d\sin(\theta)\cos(\theta) & -d\sin^2(\theta) \end{bmatrix}, \quad (\text{C9})$$

and this matrix has the following eigenvalues:

$$\lambda_1 = \frac{1}{2} \left(c - d + \sqrt{(c + d)^2 - 4cd\cos^2(\theta)} \right), \quad (\text{C10})$$

$$\lambda_2 = \frac{1}{2} \left(c - d - \sqrt{(c + d)^2 - 4cd\cos^2(\theta)} \right). \quad (\text{C11})$$

Note that $c \geq 0$ and $d \geq 0$. Without loss of generality, suppose that $c \geq d$. Then

$$0 \leq 4cd\sin^2(\theta) \quad (\text{C12})$$

$$= 4cd(1 - \cos^2(\theta)) \quad (\text{C13})$$

$$\Rightarrow -2cd \leq 2cd - 4cd \cos^2(\theta) \quad (\text{C14})$$

$$\Rightarrow c^2 - 2cd + d^2 \leq c^2 + 2cd + d^2 - 4cd \cos^2(\theta) \quad (\text{C15})$$

$$\Rightarrow (c - d)^2 \leq (c + d)^2 - 4cd \cos^2(\theta) \quad (\text{C16})$$

$$\Rightarrow c - d \leq \sqrt{(c + d)^2 - 4cd \cos^2(\theta)}. \quad (\text{C17})$$

Then it follows that the square of the trace norm of $c|\varphi'\rangle\langle\varphi'| - d|\zeta'\rangle\langle\zeta'|$ is given by

$$\begin{aligned} & \|c|\varphi'\rangle\langle\varphi'| - d|\zeta'\rangle\langle\zeta'|\|_1^2 \\ &= (|\lambda_1| + |\lambda_2|)^2 \end{aligned} \quad (\text{C18})$$

$$= \left(\frac{1}{2} \left(c - d + \sqrt{(c + d)^2 - 4cd \cos^2(\theta)} \right) - \frac{1}{2} \left(c - d - \sqrt{(c + d)^2 - 4cd \cos^2(\theta)} \right) \right)^2 \quad (\text{C19})$$

$$= (c + d)^2 - 4cd \cos^2(\theta), \quad (\text{C20})$$

concluding the proof. \blacksquare

Appendix D: Proof of Proposition 15

To prove the data-processing inequality, let \mathcal{N} be any quantum channel. We denote by $\mathcal{N}(\mathcal{E})$ the ensemble $\{(\eta_i, \mathcal{N}(\rho_i)) : i \in [r]\}$, which results from applying the channel \mathcal{N} to each state in \mathcal{E} . The optimal antidistinguishability error probability for the ensemble $\text{Err}(\mathcal{E})$ is not more than that for the ensemble $\mathcal{N}(\mathcal{E})$. To see this, let $\mathcal{M} = \{M_1, \dots, M_r\}$ be any POVM. We have

$$\text{Err}(\mathcal{M}; \mathcal{N}(\mathcal{E})) = \sum_{i \in [r]} \eta_i \text{Tr}[M_i \mathcal{N}(\rho_i)] \quad (\text{D1})$$

$$= \sum_{i \in [r]} \eta_i \text{Tr}[\mathcal{N}^\dagger(M_i) \rho_i] \quad (\text{D2})$$

$$\geq \text{Err}(\mathcal{E}). \quad (\text{D3})$$

The inequality (D3) follows because $\{\mathcal{N}^\dagger(M_1), \dots, \mathcal{N}^\dagger(M_r)\}$ is a POVM. Since (D3) holds for every POVM \mathcal{M} , we have

$$\text{Err}(\mathcal{E}) \leq \text{Err}(\mathcal{N}(\mathcal{E})). \quad (\text{D4})$$

Therefore, for all $n \in \mathbb{N}$, we get

$$-\frac{1}{n} \ln \text{Err}(\mathcal{E}^n) \geq -\frac{1}{n} \ln \text{Err}(\mathcal{N}(\mathcal{E})^n), \quad (\text{D5})$$

which implies

$$\text{E}(\rho_1, \dots, \rho_r) \geq \text{E}(\mathcal{N}(\rho_1), \dots, \mathcal{N}(\rho_r)). \quad (\text{D6})$$

Now, assume that the states in the given ensemble commute with each other. The following arguments show that the optimal error of antidistinguishing the given states is equal to that of the induced probability measures. Let P_1, \dots, P_r be the probability measures on the discrete space $[\dim(\mathcal{H})]$ induced by the states in a common eigenbasis as defined in (159), and let \mathcal{E}_{cl} be the classical ensemble $\{(\eta_i, P_i) : i \in [r]\}$. Suppose p_1, \dots, p_r are the corresponding densities

of the probability measures with respect to the counting measure μ . This gives the following representation of each state

$$\rho_i = \int_{[\dim(\mathcal{H})]} \mathbf{d}\mu(\omega) p_i(\omega) |\omega\rangle\langle\omega|, \quad i \in [r]. \quad (\text{D7})$$

We have

$$\text{Err}(\mathcal{M}; \mathcal{E}) = \sum_{i \in [r]} \eta_i \text{Tr}[M_i \rho_i] \quad (\text{D8})$$

$$= \sum_{i \in [r]} \eta_i \text{Tr} \left[M_i \left(\int_{[\dim(\mathcal{H})]} \mathbf{d}\mu(\omega) p_i(\omega) |\omega\rangle\langle\omega| \right) \right] \quad (\text{D9})$$

$$= \int_{[\dim(\mathcal{H})]} \mathbf{d}\mu(\omega) \sum_{i \in [r]} \langle\omega| M_i |\omega\rangle \eta_i p_i(\omega) \quad (\text{D10})$$

$$= \text{Err}_{\text{cl}}(\delta; \mathcal{E}_{\text{cl}}), \quad (\text{D11})$$

where δ is the decision rule given by $\delta(\omega) := (\langle\omega|M_1|\omega\rangle, \dots, \langle\omega|M_r|\omega\rangle)$. We note here that for any POVM \mathcal{M} , there corresponds a decision rule δ that satisfies (D8)–(D11). Conversely, given any decision rule δ for antidistinguishing the classical ensemble \mathcal{E}_{cl} there corresponds a POVM $\mathcal{M} = \{M_1, \dots, M_r\}$, given by

$$M_i := \int_{[\dim(\mathcal{H})]} \mathbf{d}\mu(\omega) \delta_i(\omega) |\omega\rangle\langle\omega|, \quad (\text{D12})$$

that satisfies (D8)–(D11). This then implies

$$\inf_{\mathcal{M}} \text{Err}(\mathcal{M}; \mathcal{E}) = \inf_{\delta} \text{Err}(\delta; \mathcal{E}_{\text{cl}}), \quad (\text{D13})$$

where the infima are taken over all POVMs \mathcal{M} and decision rules δ corresponding to the given quantum and classical ensembles, respectively. We have thus proved that

$$\text{Err}(\mathcal{E}) = \text{Err}_{\text{cl}}(\mathcal{E}_{\text{cl}}), \quad (\text{D14})$$

which directly implies

$$\text{E}(\rho_1, \dots, \rho_r) = \text{E}_{\text{cl}}(P_1, \dots, P_r). \quad (\text{D15})$$

Appendix E: Proof of Proposition 16

Define a map $\xi' : \mathcal{D}^r \rightarrow [0, \infty]$ by

$$\xi'(\rho_1, \dots, \rho_r) := \sup_{\mathcal{M}} \xi_{\text{cl}}(P_1^{\mathcal{M}}, \dots, P_r^{\mathcal{M}}) \quad (\text{E1})$$

as given in the right-hand side of (165). We first show that ξ' is a lower bound on any multivariate Chernoff divergence. Let $\xi : \mathcal{D}^r \rightarrow [0, \infty]$ be any multivariate quantum Chernoff divergence and ρ_1, \dots, ρ_r be arbitrary quantum states. For any measurement channel \mathcal{M} , we have

$$\xi(\rho_1, \dots, \rho_r) \geq \xi(\mathcal{M}(\rho_1), \dots, \mathcal{M}(\rho_r)) = \xi_{\text{cl}}(P_1^{\mathcal{M}}, \dots, P_r^{\mathcal{M}}). \quad (\text{E2})$$

Here we used the assumptions that ξ satisfies the data-processing inequality and reduces to the multivariate classical Chernoff divergence for commuting states. Since the inequality (E2) holds for an arbitrary measurement channel \mathcal{M} , taking the supremum over \mathcal{M} gives

$$\xi(\rho_1, \dots, \rho_r) \geq \xi'(\rho_1, \dots, \rho_r). \quad (\text{E3})$$

We now show that ξ' is a multivariate quantum Chernoff divergence; i.e., it satisfies the data processing inequality and reduces to the multivariate classical Chernoff divergence for commuting states. Consider a quantum channel \mathcal{N} and any measurement channel \mathcal{M} corresponding to a POVM $\{M_1, \dots, M_t\}$ on the output Hilbert space of the channel \mathcal{N} . Let $\mathcal{M}_{\mathcal{N}}$ be the measurement channel corresponding to the POVM $\{\mathcal{N}^\dagger(M_1), \dots, \mathcal{N}^\dagger(M_t)\}$. Let $P_1^{\mathcal{M}_{\mathcal{N}}}, \dots, P_r^{\mathcal{M}_{\mathcal{N}}}$ denote the probability measures induced by $\mathcal{M}_{\mathcal{N}}$ corresponding to the states ρ_1, \dots, ρ_r as given in the development (163)–(164). Similarly, let $Q_1^{\mathcal{M}}, \dots, Q_r^{\mathcal{M}}$ denote the probability measures induced by \mathcal{M} corresponding to the states $\mathcal{N}(\rho_1), \dots, \mathcal{N}(\rho_r)$. Since $\text{Tr}[M_j \mathcal{N}(\rho_i)] = \text{Tr}[\mathcal{N}^\dagger(M_j)(\rho_i)]$ for all i, j , it follows that $Q_i^{\mathcal{M}} = P_i^{\mathcal{M}_{\mathcal{N}}}$ for $i \in [r]$. This implies

$$\xi'(\mathcal{N}(\rho_1), \dots, \mathcal{N}(\rho_r)) = \sup_{\mathcal{M}} \xi_{\text{cl}}(Q_1^{\mathcal{M}}, \dots, Q_r^{\mathcal{M}}) \quad (\text{E4})$$

$$= \sup_{\mathcal{M}} \xi_{\text{cl}}(P_1^{\mathcal{M}_{\mathcal{N}}}, \dots, P_r^{\mathcal{M}_{\mathcal{N}}}) \quad (\text{E5})$$

$$\leq \xi'(\rho_1, \dots, \rho_r), \quad (\text{E6})$$

which means that ξ' satisfies the data-processing inequality. In the case when the states ρ_1, \dots, ρ_r commute, Theorem 7 and Proposition 15 give the following classical data-processing inequality

$$\xi_{\text{cl}}(\rho_1, \dots, \rho_r) \geq \xi_{\text{cl}}(P_1^{\mathcal{M}}, \dots, P_r^{\mathcal{M}}). \quad (\text{E7})$$

Also, the inequality in (E7) is saturated for the measurement channel corresponding to a common eigenbasis of the commuting states. Therefore, we get

$$\xi'(\rho_1, \dots, \rho_r) = \xi_{\text{cl}}(\rho_1, \dots, \rho_r). \quad (\text{E8})$$

We thus conclude that ξ' is the minimal multivariate quantum Chernoff divergence.

Appendix F: Proof of Proposition 17

Define a map $\xi'' : \mathcal{D}^r \rightarrow [0, \infty]$ by

$$\xi''(\rho_1, \dots, \rho_r) := \inf_{(\mathcal{P}, \{P_i\}_{i \in [r]})} \{\xi_{\text{cl}}(P_1, \dots, P_r) : \mathcal{P}(P_i) = \rho_i \text{ for all } i \in [r]\}, \quad (\text{F1})$$

as given in the right-hand side of (168). We first show that ξ'' is an upper bound on any multivariate Chernoff divergence. Let $\xi : \mathcal{D}^r \rightarrow [0, \infty]$ be any multivariate quantum Chernoff divergence and ρ_1, \dots, ρ_r be arbitrary quantum states. Given a preparation channel \mathcal{P} and probability measures P_1, \dots, P_r satisfying

$$\mathcal{P}(P_i) = \rho_i, \quad \text{for } i \in [r], \quad (\text{F2})$$

we have

$$\xi(\rho_1, \dots, \rho_r) = \xi(\mathcal{P}(P_1), \dots, \mathcal{P}(P_r)) \leq \xi_{\text{cl}}(P_1, \dots, P_r). \quad (\text{F3})$$

In (F3), we used the assumptions that ξ satisfies the data-processing inequality and reduces to the multivariate classical Chernoff divergence for commuting states. By taking the infimum in (F3) over preparation channels and probability measures satisfying (F2), we thus get

$$\xi(\rho_1, \dots, \rho_r) \leq \xi''(\rho_1, \dots, \rho_r). \quad (\text{F4})$$

We now show that ξ'' is a multivariate quantum Chernoff divergence; i.e., it satisfies the data processing inequality and reduces to the multivariate classical Chernoff divergence for commuting states. Let \mathcal{N} be any quantum channel. We have

$$\xi''(\mathcal{N}(\rho_1), \dots, \mathcal{N}(\rho_r)) = \inf_{\substack{(\mathcal{P}, \{P_i\}_{i \in [r]}) \\ \mathcal{P}(P_i) = \mathcal{N}(\rho_i)}} \xi_{\text{cl}}(P_1, \dots, P_r) \quad (\text{F5})$$

$$\leq \inf_{\substack{(\mathcal{P}, \{P_i\}_{i \in [r]}) \\ \mathcal{P}(P_i) = \rho_i}} \xi_{\text{cl}}(P_1, \dots, P_r) \quad (\text{F6})$$

$$= \xi''(\rho_1, \dots, \rho_r), \quad (\text{F7})$$

where the inequality follows because for every preparation channel \mathcal{P} satisfying $\mathcal{P}(P_i) = \rho_i$, its concatenation with \mathcal{N} gives another preparation channel $\mathcal{N} \circ \mathcal{P}$ that satisfies $(\mathcal{N} \circ \mathcal{P})(P_i) = \mathcal{N}(\mathcal{P}(P_i)) = \mathcal{N}(\rho_i)$. If the states ρ_1, \dots, ρ_r commute, then by the classical data-processing inequality, for any preparation channel \mathcal{P} and probability measures P_1, \dots, P_r satisfying (F2), we get

$$\xi_{\text{cl}}(\rho_1, \dots, \rho_r) = \xi_{\text{cl}}(\mathcal{P}(P_1), \dots, \mathcal{P}(P_r)) \leq \xi_{\text{cl}}(P_1, \dots, P_r). \quad (\text{F8})$$

Also, the last inequality is equality for probability distributions prepared from a spectral decomposition of the commuting states in a common orthonormal basis. Therefore, we get

$$\xi''(\rho_1, \dots, \rho_r) = \xi_{\text{cl}}(\rho_1, \dots, \rho_r). \quad (\text{F9})$$

We thus conclude that ξ'' is the maximal multivariate quantum Chernoff divergence.

Appendix G: Additivity of the optimal error exponent

Lemma 23. *Let $\mathcal{E} = \{(\eta_i, \rho_i) : i \in [r]\}$ be an ensemble of states. The following equality holds*

$$E(\rho_1, \dots, \rho_r) = \frac{1}{\ell} E(\rho_1^{\otimes \ell}, \dots, \rho_r^{\otimes \ell}) \quad \text{for all } \ell \in \mathbb{N}, \quad (\text{G1})$$

where $E(\rho_1, \dots, \rho_r)$ is the optimal error exponent defined in (31).

Proof. First, we have that

$$E(\rho_1, \dots, \rho_r) \leq \frac{1}{\ell} E(\rho_1^{\otimes \ell}, \dots, \rho_r^{\otimes \ell}) \quad \text{for all } \ell \in \mathbb{N}, \quad (\text{G2})$$

because $\{-\frac{1}{n\ell} \ln \text{Err}(\mathcal{E}^{n\ell})\}_{n \in \mathbb{N}}$ is a subsequence of $\{-\frac{1}{n} \ln \text{Err}(\mathcal{E}^n)\}_{n \in \mathbb{N}}$. We now prove the converse inequality to (G2). Let $\{M_{k,\ell}(1), \dots, M_{k,\ell}(r)\}$ be a POVM attaining $\text{Err}(\mathcal{E}^{k\ell})$ for all $k, \ell \in \mathbb{N}$. Then for any $n \in \mathbb{N}$ with $n \geq \ell$, we have

$$\text{Err}(\mathcal{E}^n) \leq \sum_{i \in [r]} \eta_i \text{Tr} \left[\rho_i^{\otimes n} \left(M_{\lfloor \frac{n}{\ell} \rfloor, \ell}(i) \otimes \mathbb{I}^{\otimes (n - \lfloor \frac{n}{\ell} \rfloor \ell)} \right) \right] \quad (\text{G3})$$

$$= \sum_{i \in [r]} \eta_i \operatorname{Tr} \left[\rho_i^{\otimes \lfloor \frac{n}{\ell} \rfloor \ell} M_{\lfloor \frac{n}{\ell} \rfloor \ell}(i) \right] \quad (\text{G4})$$

$$= \operatorname{Err}(\mathcal{E}^{\lfloor \frac{n}{\ell} \rfloor \ell}). \quad (\text{G5})$$

This implies

$$\liminf_{n \rightarrow \infty} -\frac{1}{n} \ln \operatorname{Err}(\mathcal{E}^n) \geq \liminf_{n \rightarrow \infty} -\frac{1}{\lfloor \frac{n}{\ell} \rfloor \ell} \ln \operatorname{Err}(\mathcal{E}^{\lfloor \frac{n}{\ell} \rfloor \ell}) = \frac{1}{\ell} \liminf_{k \rightarrow \infty} -\frac{1}{k} \ln \operatorname{Err}(\mathcal{E}^{k\ell}). \quad (\text{G6})$$

This completes the proof. \blacksquare

Appendix H: Limit of the regularized maximal multivariate quantum Chernoff divergence

Here we provide a proof of equation (173). We first observe that the multivariate classical Chernoff divergence is subadditive; i.e.,

$$\xi_{\text{cl}}(P_1 \otimes Q_1, \dots, P_r \otimes Q_r) \leq \xi_{\text{cl}}(P_1, \dots, P_r) + \xi_{\text{cl}}(Q_1, \dots, Q_r) \quad (\text{H1})$$

for any sets of probability densities $\{P_1, \dots, P_r\}$ and $\{Q_1, \dots, Q_r\}$ on a measurable space (Ω, \mathcal{A}) . This follows easily from the definitions of the Hellinger transform (19) and multivariate Chernoff divergence (23). So, from the definition (168), we have for $\ell, m \in \mathbb{N}$ that

$$\begin{aligned} & \xi_{\max}(\rho_1^{\otimes(\ell+m)}, \dots, \rho_r^{\otimes(\ell+m)}) \\ &= \inf_{\substack{(\mathcal{P}^{(\ell+m)}, \{P_i^{(\ell+m)}\}_{i \in [r]}) \\ \mathcal{P}^{(\ell+m)}(P_i^{(\ell+m)}) = \rho_i^{\otimes \ell} \otimes \rho_i^{\otimes m}}} \xi_{\text{cl}}(P_1^{(\ell+m)}, \dots, P_r^{(\ell+m)}) \end{aligned} \quad (\text{H2})$$

$$\leq \inf_{\substack{(\mathcal{P}^{(\ell)} \otimes \mathcal{P}^{(m)}, \{P_i^{(\ell)} \otimes P_i^{(m)}\}_{i \in [r]}) \\ \mathcal{P}^{(\ell)}(P_i^{(\ell)}) = \rho_i^{\otimes \ell}, \mathcal{P}^{(m)}(P_i^{(m)}) = \rho_i^{\otimes m}}} \xi_{\text{cl}}(P_1^{(\ell)} \otimes P_1^{(m)}, \dots, P_r^{(\ell)} \otimes P_r^{(m)}) \quad (\text{H3})$$

$$\leq \inf_{\substack{(\mathcal{P}^{(\ell)} \otimes \mathcal{P}^{(m)}, \{P_i^{(\ell)} \otimes P_i^{(m)}\}_{i \in [r]}) \\ \mathcal{P}^{(\ell)}(P_i^{(\ell)}) = \rho_i^{\otimes \ell}, \mathcal{P}^{(m)}(P_i^{(m)}) = \rho_i^{\otimes m}}} \left(\xi_{\text{cl}}(P_1^{(\ell)}, \dots, P_r^{(\ell)}) + \xi_{\text{cl}}(P_1^{(m)}, \dots, P_r^{(m)}) \right) \quad (\text{H4})$$

$$= \inf_{\substack{(\mathcal{P}^{(\ell)}, \{P_i^{(\ell)}\}_{i \in [r]}) \\ \mathcal{P}^{(\ell)}(P_i^{(\ell)}) = \rho_i^{\otimes \ell}}} \left(\xi_{\text{cl}}(P_1^{(\ell)}, \dots, P_r^{(\ell)}) \right) + \inf_{\substack{(\mathcal{P}^{(m)}, \{P_i^{(m)}\}_{i \in [r]}) \\ \mathcal{P}^{(m)}(P_i^{(m)}) = \rho_i^{\otimes m}}} \left(\xi_{\text{cl}}(P_1^{(m)}, \dots, P_r^{(m)}) \right) \quad (\text{H5})$$

$$= \xi_{\max}(\rho_1^{\otimes \ell}, \dots, \rho_r^{\otimes \ell}) + \xi_{\max}(\rho_1^{\otimes m}, \dots, \rho_r^{\otimes m}). \quad (\text{H6})$$

We have thus proved that the sequence $(\xi_{\max}(\rho_1^{\otimes \ell}, \dots, \rho_r^{\otimes \ell}))_{\ell \in \mathbb{N}}$ is subadditive. It then follows from Fekete's subadditive lemma [Fek23] that the limit $\lim_{\ell \rightarrow \infty} \xi_{\max}(\rho_1^{\otimes \ell}, \dots, \rho_r^{\otimes \ell})/\ell$ exists and is given by

$$\lim_{\ell \rightarrow \infty} \frac{1}{\ell} \xi_{\max}(\rho_1^{\otimes \ell}, \dots, \rho_r^{\otimes \ell}) = \inf_{\ell \in \mathbb{N}} \frac{1}{\ell} \xi_{\max}(\rho_1^{\otimes \ell}, \dots, \rho_r^{\otimes \ell}). \quad (\text{H7})$$

Appendix I: Proof of Equation (208)

Let $\omega \in \mathcal{D}$ be arbitrary and $(s_1, \dots, s_r) \in \mathbb{R}^r$ be any probability vector. Since the quantum states ρ_1, \dots, ρ_r have full support, we have

$$\sum_{i \in [r]} s_i D(\omega \| \rho_i)$$

$$= \sum_{i \in [r]} s_i \operatorname{Tr}[\omega(\ln \omega - \ln \rho_i)] \quad (11)$$

$$= \operatorname{Tr}[\omega \ln \omega] - \operatorname{Tr} \left[\omega \left(\sum_{i \in [r]} s_i \ln \rho_i \right) \right] \quad (12)$$

$$= \operatorname{Tr}[\omega \ln \omega] - \operatorname{Tr} \left[\omega \ln \exp \left(\sum_{i \in [r]} s_i \ln \rho_i \right) \right] \quad (13)$$

$$= \operatorname{Tr}[\omega \ln \omega] - \operatorname{Tr} \left[\omega \ln \left(\frac{\exp \left(\sum_{i \in [r]} s_i \ln \rho_i \right)}{\operatorname{Tr} \left[\exp \left(\sum_{i \in [r]} s_i \ln \rho_i \right) \right]} \cdot \operatorname{Tr} \left[\exp \left(\sum_{i \in [r]} s_i \ln \rho_i \right) \right] \right) \right] \quad (14)$$

$$= \operatorname{Tr}[\omega \ln \omega] - \operatorname{Tr} \left[\omega \ln \left(\frac{\exp \left(\sum_{i \in [r]} s_i \ln \rho_i \right)}{\operatorname{Tr} \left[\exp \left(\sum_{i \in [r]} s_i \ln \rho_i \right) \right]} \right) \right] - \ln \operatorname{Tr} \left[\exp \left(\sum_{i \in [r]} s_i \ln \rho_i \right) \right] \quad (15)$$

$$= D \left(\omega \left\| \frac{\exp \left(\sum_{i \in [r]} s_i \ln \rho_i \right)}{\operatorname{Tr} \left[\exp \left(\sum_{i \in [r]} s_i \ln \rho_i \right) \right]} \right) - \ln \operatorname{Tr} \left[\exp \left(\sum_{i \in [r]} s_i \ln \rho_i \right) \right] \quad (16)$$

$$\geq -\ln \operatorname{Tr} \left[\exp \left(\sum_{i \in [r]} s_i \ln \rho_i \right) \right], \quad (17)$$

where the inequality follows from the non-negativity of quantum relative entropy for quantum states. The lower bound is achieved by picking $\omega = \frac{\exp(\sum_{i \in [r]} s_i \ln \rho_i)}{\operatorname{Tr}[\exp(\sum_{i \in [r]} s_i \ln \rho_i)]}$, so that

$$\begin{aligned} & \inf_{\omega \in \mathcal{D}} \sum_{i \in [r]} s_i D(\omega \| \rho_i) \\ &= \inf_{\omega \in \mathcal{D}} D \left(\omega \left\| \frac{\exp \left(\sum_{i \in [r]} s_i \ln \rho_i \right)}{\operatorname{Tr} \left[\exp \left(\sum_{i \in [r]} s_i \ln \rho_i \right) \right]} \right) - \ln \operatorname{Tr} \left[\exp \left(\sum_{i \in [r]} s_i \ln \rho_i \right) \right] \end{aligned} \quad (18)$$

$$= -\ln \operatorname{Tr} \left[\exp \left(\sum_{i \in [r]} s_i \ln \rho_i \right) \right]. \quad (19)$$

This directly gives (208).