

# THE LIMITS OF DETERMINACY IN SECOND ORDER ARITHMETIC

ANTONIO MONTALBÁN AND RICHARD A. SHORE

ABSTRACT. We establish the precise bounds for the amount of determinacy provable in second order arithmetic. We show that for every natural number  $n$ , second order arithmetic can prove that determinacy holds for Boolean combinations of  $n$  many  $\mathbf{\Pi}_3^0$  classes, but it cannot prove that all finite Boolean combinations of  $\mathbf{\Pi}_3^0$  classes are determined. More specifically, we prove that  $\Pi_{n+2}^1\text{-CA}_0 \vdash n\text{-}\mathbf{\Pi}_3^0\text{-DET}$ , but that  $\Delta_{n+2}^1\text{-CA} \not\vdash n\text{-}\mathbf{\Pi}_3^0\text{-DET}$ , where  $n\text{-}\mathbf{\Pi}_3^0$  is the  $n$ th level in the difference hierarchy of  $\mathbf{\Pi}_3^0$  classes.

We also show some conservativity results that imply that reversals for the theorems above are not possible. We prove that for every true  $\Sigma_4^1$  sentence  $T$  (as for instance  $n\text{-}\mathbf{\Pi}_3^0\text{-DET}$ ) and every  $n \geq 2$ ,  $\Delta_n^1\text{-CA}_0 + T + \Pi_\infty^1\text{-TI} \not\vdash \Pi_n^1\text{-CA}_0$  and  $\Pi_{n-1}^1\text{-CA}_0 + T + \Pi_\infty^1\text{-TI} \not\vdash \Delta_n^1\text{-CA}_0$ .

## 1. INTRODUCTION

The general enterprise of calibrating the strength of classical mathematical theorems in terms of the axioms (typically of set existence) needed to prove them was begun by Harvey Friedman in [1971]. In that paper he worked primarily in the set theoretic settings of subsystems (and extensions) of ZFC. Actually, almost all of classical mathematics can be formalized in the language of second order arithmetic. This language consists of that of ordinary (first order) arithmetic and membership only for numbers in sets of numbers. It also has variables for, and quantification over, numbers and subsets of the natural numbers  $\mathbb{N}$ . (In terms of representing classical mathematics in this setting, we are restricting our attention to the countable case for algebraic and combinatorial results and, analogously, the separable situation for analytic or topological ones.) Moreover, the standard theorems of classical (countable) mathematics can be established in systems requiring only the basic axioms for arithmetic and set existence axioms just for subsets of  $\mathbb{N}$ . Realizing this, Friedman [1975] moved to the setting of second order arithmetic and subsystems of its full theory  $Z_2$  which assumes some basic axioms of arithmetic and the existence of each subset of  $\mathbb{N}$  defined by a formula of second order arithmetic. (See §2.3 for the precise definitions of our language, structures, axiom systems and their models.) Many researchers have since contributed to this endeavor but the major systematic developer and expositor since Friedman has been Stephen Simpson and the basic source for both background material and extensive results is his book [2009].

Five subsystems of  $Z_2$  of strictly increasing strength emerged as the core of the subject with the vast majority of classical mathematical theorems being provable in one of them. (Formal definitions of the subsystems of second order arithmetic used in this paper are in §2.3.)

---

*Date:* Submitted Jan 22nd, 2010.

Compiled: January 22, 2010.

Montalbán was partially supported by NSF grant DMS-0901169, and by the AMS centennial fellowship. Shore was partially supported by NSF Grants DMS-0554855 and DMS-0852811, Grant 13408 from the John Templeton Foundation and as a visitor by the University of Chicago.

Indeed, relative to the weakest of them ( $\text{RCA}_0$ , which corresponds simply to computable mathematics) almost all the theorems studied turned out to be equivalent to one of these five systems. Here the equivalence of a theorem  $T$  to a system  $S$  means that not only is the theorem  $T$  provable in  $S$  but that  $\text{RCA}_0 + T$  proves all the axioms of  $S$  as well. Thus the system  $S$  is precisely what is needed to establish  $T$  and gives a characterization of its (proof theoretic) strength. It is this approach that gives the subject the name of Reverse Mathematics. In standard mathematics one proves a theorem  $T$  from axioms  $S$ . Here one then tries to reverse the process (over a weak base theory) by proving the axioms of  $S$  from  $T$  (and  $\text{RCA}_0$ ). In fact, for those theorems that are not true effectively (essentially the same as provable in  $\text{RCA}_0$ ), the vast majority turn out to be equivalent in the sense of reverse mathematics to one of the two weakest of these systems ( $\text{WKL}_0$  or  $\text{ACA}_0$ ). Only a handful are equivalent to one of the two stronger systems ( $\text{ATR}_0$  or  $\Pi_1^1\text{-CA}_0$ ) and just a couple lie beyond them. (See Simpson [2009] for examples.) There is one now known (Mummert and Simpson [2005]) to be at the next level ( $\Pi_2^1\text{-CA}_0$ ) and (as far as we know) none that are not known to be provable a bit beyond this level but still at one less than  $\Pi_3^1\text{-CA}_0$ .

In this paper we supply a natural hierarchy of theorems that require, respectively, each of the natural levels of set existence assumptions going from the previous known systems all the way up to full second order arithmetic. These theorems come from the realm in which the subject began, axioms of determinacy. The subject here is that of two person games with complete information. Our *games* (at least in this section) are played by two players I and II. They alternate *playing* natural numbers with I playing first to produce a *play of the game* which is a sequence  $x \in \omega^\omega$ . A *game*  $G_A$  is specified by a subset  $A$  of  $\omega^\omega$ . We say that I *wins* a play  $x$  of the game  $G_A$  specified by  $A$  if  $x \in A$ . Otherwise II wins that play. A *strategy* for I (II) is a function  $\sigma$  from strings  $p$  of even (odd) length into  $\omega$ . We say that the game  $G_A$  is *determined* if there is a winning strategy for I or II in this game. (More details and terminology can be found in §2.1 where we switch from Baire space,  $\omega^\omega$ , to Cantor space,  $2^\omega$  to better match the standard setting for reverse mathematics and for other technical reasons. Basic references are Moschovakis [2009] and Kechris [1995].)

Now, in its full form, the Axiom of Determinacy says that all games  $G_A$  are determined. It has many surprising implications (e.g. all sets of reals are Lebesgue measurable) that contradict the Axiom of Choice. So instead, one considers restricted classes of games: If  $\Gamma$  is a class of sets  $A$  (of reals), then we say that  $\Gamma$  is *determined* if  $G_A$  is determined for every  $A \in \Gamma$ . We denote the assertion that  $\Gamma$  is determined by  $\Gamma$  *determinacy* or  $\Gamma\text{-DET}$ . One is then interested in determinacy for “simple” or easily definable classes  $\Gamma$ . One hopes that, for such  $\Gamma$ , determinacy will be provable (in some system of set theory including the axiom of choice) and that many of the consequences of the full axiom will then follow for sets in  $\Gamma$ . This has, indeed, turned out to be the case and a remarkably rich and extensive theory with applications in measure theory, descriptive set theory, harmonic analysis, ergodic theory, dynamical systems and other areas has been developed. (In addition to the basic texts mentioned above, one might refer, for example, to Kechris and Becker [1996], Kechris and Louveau [1987], Kechris and Miller [2004] and Foreman [2000] for sample applications.) A crowning achievement of this theory has been the calibration of the higher levels of determinacy for  $\Gamma$  in the projective hierarchy of sets of reals that begins with the analytic sets and progresses by complementation and projection. Here work of Martin, Steel and Woodin (Martin and Steel [1989], Woodin [1988]) has precisely characterized the large

cardinal assumptions needed to prove each level of determinacy in the projective hierarchy (and beyond).

We are here concerned with lower levels of determinacy. Friedman's first foray [1971] into the area that grew into reverse mathematics dealt with these issues. He famously proved that Borel determinacy is not provable in ZFC without the power set axiom. Indeed, he showed that one needed  $\aleph_1$  many iterations of the power set to prove it. Martin [1975], then showed that Borel determinacy is provable in ZFC and provided a level by level analysis of Borel hierarchy and the number of iterations of the power set needed to establish determinacy at those levels. Moving from set theory to second order arithmetic and the realm of what is now called reverse mathematics, Friedman [1971] also showed that  $\Sigma_5^0$  determinacy (i.e. for  $G_{\delta\sigma\delta\sigma}$  sets or see §2.2) is not provable in full second order arithmetic. Martin [1974a], [n.d., Ch. 1] improved this to  $\Sigma_4^0$  determinacy (i.e.  $F_{\sigma\delta\sigma}$  sets). He also presented [1974] [n.d., Ch. 1] a proof of  $\Delta_4^0$  determinacy (sets that are both  $F_{\sigma\delta\sigma}$  and  $G_{\delta\sigma\delta}$ ) that he said could be carried out in  $Z_2$ . This seemed to fully determine the boundary of determinacy that is provable in second order arithmetic and left only the first few levels of the Borel hierarchy to be analyzed from the viewpoint of reverse mathematics.

The first very early result (essentially Steel [1976] see also Simpson [2009 V.8]) was that  $\Sigma_1^0$  (open) determinacy is equivalent (over  $\text{RCA}_0$ ) to  $\text{ATR}_0$ . Tanaka [1990] then showed that  $\Pi_1^1\text{-CA}_0$  is equivalent to  $\Sigma_1^0 \wedge \Pi_1^0$  (intersections of open and closed sets) determinacy. Moving on to  $\Sigma_2^0$  ( $F_\sigma$ ) determinacy, Tanaka [1991] showed that it was equivalent to an unusual system based on closure under monotonic  $\Sigma_1^1$  definitions. At the level of  $\Delta_3^0$  (sets in both  $F_{\sigma\delta}$  and  $G_{\delta\sigma}$ ) determinacy, MedSalem and Tanaka [2007] showed that each of  $\Pi_2^1\text{-CA}_0 + \Pi_3^1\text{-TI}_0$  (an axiom system of transfinite induction) and  $\Delta_3^1\text{-CA}_0 + \Sigma_3^1\text{-IND}_0$  (induction for  $\Sigma_3^1$  formulas) prove  $\Delta_3^0\text{-DET}$  but  $\Delta_3^1\text{-CA}_0$  alone does not. They improve these results in [2008] by showing that  $\Delta_3^0\text{-DET}$  is equivalent (over  $\Pi_3^1\text{-TI}_0$ ) to another system based on transfinite combinations of  $\Sigma_1^1$  inductive definitions. Finally, Welch [2009] has shown that  $\Pi_3^1\text{-CA}_0$  proves not only  $\Pi_3^0$  ( $F_{\delta\sigma}$ ) determinacy but even that there is a  $\beta$ -model (Definition 2.9) of  $\Delta_3^1\text{-CA}_0 + \Pi_3^0\text{-DET}$ . In the other direction, he has also shown that, even augmented by an axiom about the convergence of arithmetical quasi-inductive definitions,  $\Delta_3^1\text{-CA}_0$  does not prove  $\Pi_3^0\text{-DET}$ . The next level of determinacy is then  $\Delta_4^0$ .

Upon examining Martin's proof of  $\Delta_4^0$ -determinacy as sketched in [1974] and then later as fully written out in [n.d., Ch. 1], it seemed to us that one cannot actually carry out his proof in  $Z_2$ . Essentially, the problem is that the proof proceeds by a complicated induction argument over an ordering whose definition seems to require the full satisfaction relation for second order arithmetic. This realization opened up anew the question of determining the boundary line for determinacy provable in second order arithmetic. We answer that question in this paper by analyzing the strength of determinacy for the finite levels of the difference hierarchy on  $\Pi_3^0$  ( $F_{\sigma\delta}$ ) sets, the  $m\text{-}\Pi_3^0$  sets. (As defined in §2.2, these give a natural hierarchy for the finite Boolean combinations of  $\Pi_3^0$  sets.)

In the positive direction (§4), we carefully present a variant of Martin's proof specialized and simplified to the finite levels of the difference hierarchy on  $\Pi_3^0$  along with the analysis needed to determine the amount of comprehension used in the proof for each level of the hierarchy. (It is a classical theorem of Kuratowski [1958] that extending the hierarchy into the transfinite to level  $\aleph_1$  gives all the  $\Delta_4^0$  sets. Martin's proof proceeds by an induction

encompassing all these levels.) This establishes an upper bound for the provability of the  $m\text{-}\Pi_3^0$  sets in  $Z_2$ .

**Theorem 1.1.** *For each  $m \geq 1$ ,  $\Pi_{m+2}^1\text{-CA}_0 \vdash m\text{-}\Pi_3^0\text{-DET}$ .*

In the other direction, we prove that this upper bound is sharp in terms of the standard subsystems of second order arithmetic.

**Theorem 1.2.** *For every  $m \geq 1$ ,  $\Delta_{m+2}^1\text{-CA}$  does not prove  $m\text{-}\Pi_3^0\text{-DET}$ .*

**Corollary 1.3.** *Determinacy for the class of all finite Boolean combinations of  $\Pi_3^0$  classes of reals ( $\omega\text{-}\Pi_3^0\text{-DET}$ ) cannot be proved in second order arithmetic. As these classes are all (well) inside  $\Delta_4^0$ ,  $Z_2$  does not prove  $\Delta_4^0\text{-DET}$ .*

Note that by Theorem 1.1, any model of second order arithmetic in which the natural numbers are the standard ones (i.e.  $\mathbb{N}$ ) does satisfy  $\omega\text{-}\Pi_3^0\text{-DET}$  and so the counterexample for its failure to be a theorem of  $Z_2$  must be nonstandard. Of course, it can be constructed by a compactness argument or, equivalently, as an ultraproduct of the  $L_{\alpha_n}$  defined below. In contrast, the counterexamples from Friedman [1971] and Martin [1974a], [n.d., Ch. 1] are all even  $\beta$ -models, so not only with its numbers standard but all its “ordinals” (well orderings) are true ordinals (well orderings) as well.

We can reformulate this limitative result in the setting of set theory by noting the following conservation result.

**Proposition 1.4.**  *$ZFC^-$  ( $ZFC$  with collection but without the power set axiom) and even with a definable well ordering of the universe assumed as well, is a  $\Pi_4^1$  conservative extension of  $Z_2$ .*

**PROOF:** (Sketch) This fact should be “well known” and certainly follows from the extensive analysis of the relations between models of (subsystems of) second order arithmetic and those of the form  $L(X)$  in Simpson [2009, VII]. Basically, given a model  $\mathcal{M}$  of  $Z_2$  and an  $X$  a set in  $\mathcal{M}$ , one defines an interpretation  $L^{\mathcal{M}}(X)$  (defined over the well orderings of  $\mathcal{M}$ ) of  $L(X)$  in  $\mathcal{M}$  and checks that it is a model of  $ZFC^-$ . This is detailed work but fairly straight forward for  $Z_2$ . Much of the work in Simpson [2009 VII.7] (whose ideas are described there as “probably well known but we have been unable to find bibliographic references for them”) is devoted to showing that the facts required for the interpretation to satisfy the basic axioms of set theory can be derived in  $\text{ATR}_0$  and the Shoenfield absoluteness theorem (VII.4.14) in  $\Pi_1^1\text{-CA}_0$ . As a guide we note that the basic translations and interpretations between second order arithmetic and a simple set theory are in VII.3. The relations with  $L$  and  $L(X)$  and their basic properties are in VII.4. The material needed to verify the comprehension axioms is in VII.5 with Theorem VII.5.9(10) and the proof of its Corollary VII.5.11 being the closest to what is needed here. The material for replacement is in VII.6 with Lemma VII.6.15 and Theorem VII.6.16 being the closest to what is needed here. Finally, as  $L^{\mathcal{M}}(X)$  is a model of  $V = L(X)$ ,  $L^{\mathcal{M}}(X)$  satisfies even global choice, i.e. there is a definable well ordering of the universe. Thus  $L^{\mathcal{M}}(X)$  is a model of  $ZFC^-$  (and global definable choice).

Now if  $T = \forall X \exists Y \forall Z \exists W \varphi$  is  $\Pi_4^1$  and false in some  $\mathcal{M} \models Z_2$ , we take  $X$  to be the counterexample and consider  $L^{\mathcal{M}}(X)$  and any  $Y \in L^{\mathcal{M}}(X)$ . By assumption,  $\mathcal{M} \models \exists Z \forall W \neg \varphi(X, Y)$  and so by Shoenfield absoluteness,  $L^{\mathcal{M}}(X) \models \exists Z \forall W \neg \varphi(X, Y)$ . Thus  $T$  fails in  $L^{\mathcal{M}}(X)$  for the desired contradiction.  $\square$

It is worth pointing out that  $\Pi_4^1$  is as far as conservation results can go for  $ZFC^-$  over  $Z_2$ . As Simpson [2009, VII.6.3] points out, Feferman and Levy have produced a model of  $Z_2$  in which a  $\Sigma_3^1$ -AC axiom fails. Now each  $\Sigma_3^1$ -AC axiom is clearly a  $\Sigma_5^1$  sentence and a theorem of  $ZFC^-$ . Now if  $T = \forall X \exists Y \forall Z \exists W \varphi$  is  $\Pi_4^1$  and false in some  $\mathcal{M} \models Z_2$ , we take  $X$  to be the counterexample and consider  $L^{\mathcal{M}}(X)$  and any  $Y \in L^{\mathcal{M}}(X)$ . By assumption,  $\mathcal{M} \models \exists Z \forall W \neg \varphi(X, Y)$  and so by Shoenfield absoluteness,  $L^{\mathcal{M}}(X) \models \exists Z \forall W \neg \varphi(X, Y)$ . Thus  $T$  fails in  $L^{\mathcal{M}}(X)$  which is a model of  $ZFC^-$ .

**Corollary 1.5.** *Determinacy for the class of all finite Boolean combinations of  $\Pi_3^0$  classes of reals ( $\omega$ - $\Pi_3^0$ -DET) and so, a fortiori,  $\Delta_4^0$ -DET cannot be proved in  $ZFC^-$ .*

In fact, our counterexamples that establish Theorem 1.2 (the games with no strategy in  $L_{\alpha_n}$  as described below) are all given by effective versions of the  $m$ - $\Pi_3^0$  sets where the initial  $\Pi_3^0$  sets are effectively defined, i.e. given by recursive unions and intersections starting with closed sets that are “lightfaced”, i.e. defined by  $\Pi_1^0$  formulas of first order arithmetic without real (second order) parameters. (See the remarks in the last paragraph of §2.2. We use notations such as  $\Pi_1^0$  and  $\Pi_3^0$  to denote these “lightfaced” versions of the Borel classes.) This gives rise to a Gödel like phenomena for second order arithmetic with natural mathematical  $\Sigma_2^1$  statements saying that specific games have strategies and containing no references to provability.

**Theorem 1.6.** *There is a  $\Sigma_2^1$  formula  $\varphi(x)$  with one free number variable  $x$ , such that, for each  $n \in \omega$ ,  $Z_2 \vdash \varphi(\underline{n})$  but  $Z_2 \not\vdash \forall n \varphi(n)$ .*

Of course, on their face Theorems 1.1 and 1.2 along with Corollary 1.3 produce a sequence of  $\Pi_3^1$  formulas  $\psi(n)$  that have the same proof theoretic properties while eliminating the references to syntax and recursion theory present in the  $\varphi$  of Theorem 1.6. They simply state the purely mathematical propositions that all  $n$ - $\Pi_3^0$  games are determined.

The plan for the first few sections of this paper is as follows. We provide basic definitions and notations about games and determinacy, hierarchies of sets and formulas and subsystems of second order arithmetic in §2. Basic facts about Gödel’s constructible universe  $L$  not found in the standard texts are given in §3. The proof of Theorem 1.1 is in §4.

To prove Theorem 1.2, we will work in set theory or more specifically in fragments of  $ZFC + V=L$  instead of directly in second order arithmetic. We can do this as Simpson [2009, VII] shows how one can move back and forth between systems of second order arithmetic and subsystems of  $ZFC + V=L(X)$ , once one has  $ATR_0$  (or at times  $\Pi_1^1$ - $CA_0$ ) as a base theory. (Note also that, as mentioned above, these systems are equivalent to even weaker forms of determinacy than any we consider and so are provable even in  $\Pi_3^0$ -DET.)

Let  $\alpha_n$  denote the first  $n$ -admissible ordinal, and  $\mathbb{R}$  the set of subsets of  $\omega$ . (See §3 for the basic definitions.) By Lemma 3.2 and Simpson [2009, VII.5.3],  $\mathbb{R} \cap L_{\alpha_n}$  is a  $\beta$ -model of  $\Delta_{n+1}^1$ -CA for  $n \geq 2$ . Let  $Th_{\alpha_n}$  denote the true theory of  $L_{\alpha_n}$ . We think of  $Th_{\alpha_n}$  as a subset of  $\omega$ , or as the characteristic function (in  $2^\omega$ ) for the set of indices of sentences of set theory (without parameters) true in  $L_{\alpha_n}$ . As every element of  $L_{\alpha_n}$  is definable in  $L_{\alpha_n}$  without parameters (Lemma 3.6), the Gödel-Tarski undefinability of truth (a simple diagonal argument at this point) says that  $Th_{\alpha_n} \notin L_{\alpha_n}$ . Our plan is to show that, for  $n \geq 2$ ,  $(n-1)$ - $\Pi_3^0$ -DET does not hold in  $L_{\alpha_n}$  by showing that, if it did,  $Th_{\alpha_n}$  would be a member of  $L_{\alpha_n}$ .

As the  $L_{\alpha_n}$  are  $\beta$ -models, Theorem 1.2 can be immediately improved in various ways by including in the base theory any axioms true in all  $\beta$ -models, as for example:

**Theorem 1.7.** *For every  $n \geq 1$ ,  $\Sigma_{n+2}^1\text{-DC} + \Pi_\infty^1\text{-TI}$  does not prove  $n\text{-}\Pi_3^0\text{-DET}$ .*

PROOF: Every  $\beta$ -model is obviously a model of  $\Pi_\infty^1\text{-TI}$  and, as  $L_{\alpha_{n+1}} \models \Delta_{n+1}^1\text{-CA}$ , it also satisfies  $\Sigma_{n+2}^1\text{-DC}$  (which is stronger than  $\Delta_{n+1}^1\text{-CA}$ ) by Simpson [2009, VII.6.18].  $\square$

The key result needed to prove Theorem 1.2 is the following.

**Lemma 1.8.** *For every  $n \geq 2$ , there is a game  $G$  that is  $(n-1)\text{-}\Pi_3^0$ , such that, if we interpret the play of each player as the characteristic function of a set of sentences in the language of set theory, then*

- (1) *If I plays  $Th_{\alpha_n}$ , he wins.*
- (2) *If I does not play  $Th_{\alpha_n}$  but II does, then II wins.*

We prove this Lemma in §5 but we now show how it implies our main result, Theorem 1.2.

PROOF OF THEOREM 1.2: As we mentioned above,  $\mathbb{R} \cap L_{\alpha_n}$  is a  $\beta$ -model of  $\Delta_{n+1}^1\text{-CA}$  and so it suffices to show that  $L_{\alpha_n} \not\models (n-1)\text{-}\Pi_3^0\text{-DET}$ . Let  $G$  be as in the Lemma; and suppose it is determined in  $L_{\alpha_n}$ . Player II cannot have a winning strategy for  $G$  in  $L_{\alpha_n}$  because if II has a winning strategy  $\sigma$  in  $L_{\alpha_n}$ ,  $\sigma$  would also be a winning strategy in  $V$  as  $\mathbb{R} \cap L_{\alpha_n}$  is a  $\beta$ -model (and  $\sigma$  being a winning strategy for  $G$  is a  $\Pi_1^1$  property). But, I has a winning strategy for  $G$  in  $V$  by clause (1) of the Lemma. So, I must have a winning strategy  $\sigma$  for  $G$  in  $L_{\alpha_n}$ . Again, as  $\mathbb{R} \cap L_{\alpha_n}$  is a  $\beta$ -model,  $\sigma$  is truly a winning strategy for I (in  $V$ ). We claim that if II plays so as to simply copy I's moves, then  $\sigma$  has to play  $Th_{\alpha_n}$ . If not, then at some first point I plays a bit that is different from  $Th_{\alpha_n}$ . At this point II could stop copying I and just continue playing  $Th_{\alpha_n}$  and he would win (by clause (2) of the Lemma). Thus  $\sigma$  would not be a truly winning strategy for I (in  $V$ ). We conclude that  $\sigma$  computes  $Th_{\alpha_n}$  as the sequence of I's plays against II's copying his moves and so  $Th_{\alpha_n} \in L_{\alpha_n}$  for the desired contradiction.  $\square$

In the spirit of reverse mathematics one should now ask for reversals showing that  $m\text{-}\Pi_3^0\text{-DET}$  implies  $\Pi_{m+2}^1\text{-CA}_0$  or something along these lines. Nothing like this is, however, possible. MedSalem and Tanaka [2007] have shown that even full Borel ( $\Delta_1^1$ ) determinacy does not imply even  $\Delta_2^1\text{-CA}_0$ . (Indeed they also show that it does not imply either induction for  $\Sigma_3^1$  formulas or  $\Pi_2^1\text{-TI}_0$ .) Their proofs proceed via constructing countably coded  $\beta$ -models and appealing to the second Gödel incompleteness theorem.

In §6, we provide a very different approach that applies to any true (or even consistent with ZFC)  $\Sigma_4^1$  sentence  $T$  and shows that no such sentence can imply  $\Delta_2^1\text{-CA}_0$  even for  $\beta$ -models. The counterexamples are all initial segments of  $L(X)$  (for  $X$  a witness to the  $\Sigma_4^1$  sentence). We also show that for any such  $T$  and  $n \geq 2$ ,  $\Delta_n^1\text{-CA}_0 + T + \Pi_\infty^1\text{-TI} \not\models \Pi_n^1\text{-CA}_0$  and  $\Pi_n^1\text{-CA}_0 + T + \Pi_\infty^1\text{-TI} \not\models \Delta_{n+1}^1\text{-CA}_0$  (even for  $\beta$ -models). These results are also suitably generalized to  $\Sigma_m^1$  theorems of ZFC in an optimal way. As Borel and  $m\text{-}\Pi_3^0$  determinacy are  $\Pi_3^1$  theorems of ZFC, these general conservation results apply and we have the following results:

**Borel-DET** +  $\Pi_\infty^1\text{-TI} \not\models \Delta_2^1\text{-CA}_0$ .

For  $n \geq 0$ ,  $\Delta_{n+2}^1\text{-CA}_0 + n\text{-}\Pi_3^0\text{-DET} + \Pi_\infty^1\text{-TI} \not\models \Pi_{n+2}^1\text{-CA}_0$ .

Thus, by Theorem 1.1, for  $n \geq 0$ ,  $n\text{-}\Pi_3^0\text{-DET}$  is a consequence of  $\Pi_{n+2}^1\text{-CA}_0$  incomparable (in terms of provability) with  $\Delta_{k+2}^1\text{-CA}_0$  for all  $k \leq n$  while  $\Delta_{n+2}^1\text{-CA}_0 + n\text{-}\Pi_3^0\text{-DET}$

is a system strictly between  $\Delta_{n+2}^1\text{-CA}_0$  and  $\Pi_{n+2}^1\text{-CA}_0$  given by a mathematically natural proposition.

## 2. DEFINITIONS AND NOTATIONS

**2.1. Games and Determinacy.** From now on, our basic playing field is not Baire space but Cantor space,  $2^\omega$ , the set of infinite (length  $\omega$ ) binary sequences (reals),  $x$  which we identify with the set  $X \subseteq \mathbb{N}$  with its characteristic function  $x$ . Our *games* are played by two players I and II. They alternate *playing* 0 or 1 with I playing first to produce a *play of the game* which is a real  $x$ . A *game*  $G_A$  is specified by a subset  $A$  of  $2^\omega$ . We say that I *wins* a play  $x$  of the game  $G_A$  specified by  $A$  if  $x \in A$ . Otherwise II wins that play. A *strategy* for I (II) is a function  $\sigma$  from binary strings  $p$  of even (odd) length into  $\{0, 1\}$ . The intuition here is that at any finite string of even (odd) length, a *position in the game* at which it is I's (II's) *turn to play*, the strategy  $\sigma$  instructs I (II) to play  $\sigma(p)$ . We say that a position  $q$  (play  $x$ ) is *consistent with*  $\sigma$  if, for every  $p \subset q$  ( $x$ ) of even (odd) length,  $\sigma \hat{\ } p \subseteq q$  ( $x$ ). (We use  $\hat{\ }$  for concatenation of strings and confuse a number  $i$  with the string  $\langle i \rangle$ .) The strategy  $\sigma$  is a winning strategy for I (II) in the game  $G_A$  if every play consistent with  $\sigma$  is in  $A$  ( $\bar{A}$ ). (We use  $\bar{A}$  for  $2^\omega \setminus A$ , the *complement* of  $A$  in Cantor space.)

We say that the game  $G_A$  is *determined* if there is a winning strategy for I or II in this game. If  $\Gamma$  is a class of sets  $A$  (of reals), then we say that  $\Gamma$  is *determined* if  $G_A$  is determined for every  $A \in \Gamma$ . We denote the assertion that  $\Gamma$  is determined by  $\Gamma$  *determinacy* or  $\Gamma$ -DET.

Note that easy codings translate between games on Baire space to ones on Cantor space. Given an  $A \subseteq \omega^\omega$  we code it by  $\hat{A} \subseteq 2^\omega$  with elements precisely the ones of the form  $\hat{f} = 0^{f(0)}10^{f(1)}10^{f(2)} \dots 10^{f(n)}1 \dots$  where we use  $0^k$  to denote the sequence consisting of  $k$  many 0's. A strategy for  $G_A$  can be gotten effectively from one for  $\hat{A} = \{x \in 2^\omega \mid [\exists n \forall m > n(x(2m+1) = 0) \text{ or } \forall n \exists m > n(x(2m) = 1) \ \& \ \text{for the } f \text{ such that } x = \hat{f}, f \in A]\}$ . In the other direction, given an  $A \subseteq 2^\omega$  a strategy for  $G_A$  can be found effectively from one for  $G_{\hat{A}}$  where  $\hat{A} = \{f \in \omega^\omega \mid \exists n(f(2n+1) \notin \{0, 1\} \text{ or } f \in A)\}$ . Thus if  $\Gamma$  is rich enough to be closed under these operations, it makes no difference whether we play in Baire space or Cantor space. It is clear that once  $\Gamma$  is at least at the level of  $\Pi_3^0$  ( $F_{\sigma\delta}$ ) (or even  $\Delta_3^0$ )  $\Gamma$  determinacy in one space is equivalent to it in the other, and in this paper we consider only classes from  $\Pi_3^0$  and above. For the record, we note that there are significant differences at levels below  $\Delta_3^0$ . In contrast to the standard low level results for Baire space mentioned in §1, Nemoto, MedSalem and Tanaka [2007] show that in Cantor space  $\Delta_1^0$ -DET is equivalent (over  $\text{RCA}_0$ ) to each of  $\Sigma_1^0$ -DET and  $\text{WKL}_0$ .  $\text{ACA}_0$  is equivalent to  $(\Sigma_2^0 \wedge \Pi_2^0)$ -DET.  $\Delta_2^0$ -DET is equivalent to each of  $\Sigma_2^0$ -DET and  $\text{ATR}_0$  while, as indicated above,  $\Delta_3^0$ -DET is equivalent to the same level of determinacy in Baire space.

As usual, we let  $\check{\Gamma} = \{\bar{A} \mid A \in \Gamma\}$ . Idiosyncratically, and only for the purposes of the next Lemma, we let  $A^* = \{0 \hat{\ } x \mid x \in A\} \cup \{1 \hat{\ } x \mid x \in A\}$  and  $\Gamma^* = \{A^* \mid A \in \Gamma\}$ .

**Lemma 2.1.** ( $\text{RCA}_0$ ) *If  $\Gamma^* \subseteq \Gamma$  and  $\Gamma$  is determined then so is  $\check{\Gamma}$ .*

PROOF: A winning strategy  $\sigma$  for I (II) in  $G_{A^*}$  is easily converted into one  $\tau$  for II (I) in  $G_{\bar{A}}$ :  $\tau(p) = \sigma(\tau(p) = \sigma(\emptyset) \hat{\ } p) (\sigma(0 \hat{\ } p))$ .  $\square$

This is as much as we need about determinacy for our proof of its failure for various  $\Gamma$  in specified models of  $\Delta_{n+2}^1\text{-CA}_0$  in §5. To show that determinacy holds for these  $\Gamma$  in  $\Pi_{n+2}^1\text{-CA}_0$  in §4, we need to generalize these notions a bit.

We now allow games to be played on arbitrary binary trees  $T$ . The idea here is that we replace  $2^\omega$  by  $[T]$ , the *set of paths through  $T$* ,  $\{x \mid \forall n(x \upharpoonright n \in T)\}$ . (So the basic notion takes  $T$  to be  $2^{<\omega}$ , the tree of all finite binary sequences.) A *(binary) tree* is a set of finite binary sequences closed under initial segments and with no dead ends, i.e.  $\forall p \in T(p \hat{\ } 0 \in T$  or  $p \hat{\ } 1 \in T)$ . If  $T$  is a tree and  $p \in T$  then  $T_p$  is the subtree of all  $q \in T$  comparable with  $p$ ,  $T_p = \{q \in T \mid q \subseteq p \text{ or } p \subseteq q\}$ . (A *subtree of a tree  $T$*  is a subset  $T$  which is also a tree.) A *play (real)  $x$  is one in the game on  $T$*  if  $x \in [T]$ . A *game  $G(A, T)$*  is now specified by an  $A \subseteq [T]$ . The other notions defined above are generalized accordingly with the positions restricted to  $T$  and the plays to  $[T]$ . A *quasistrategy for I (II) in  $G(A, T)$*  is a subtree  $S$  of  $T$  such that if  $p \in S$  is of odd (even) length then the immediate successors in  $S$  are precisely the same as the ones in  $T$ . Thus a strategy for I (II) can be viewed as a quasistrategy in which each  $p \in S$  of even (odd) length has precisely one immediate successor in  $S$ . A position  $p$  or play  $x$  in  $G(A, T)$  being *consistent with  $S$*  now just means that  $p \in S$  or  $x \in [S]$ . We say that  $S$  is a *winning quasistrategy for I (II)* if every  $x$  consistent with  $S$  is in  $A$  ( $\bar{A}$ ). If I (II) does not have a winning strategy in  $G(A, T)$  then we define II's (I's) *nonlosing quasistrategy  $W$*  in this game to be the set of all  $p \in T$  such that I (II) has no winning strategy in  $G(A, T_p)$ . If this set exists (e.g. we have enough comprehension as when  $A$  is arithmetic and we have at least  $\Pi_2^1\text{-CA}_0$ ), it is easily seen to be a quasistrategy for II (I) by induction on the length of  $p \in W$ .

**2.2. Classes and Hierarchies.** We have to deal with several classes and hierarchies of both formulas and sets that have standard but at times conflicting (or overlapping) designations. We try to use as single unambiguous system through out the paper while choosing notations whose meaning in specific contexts should be evident to readers familiar with work in the appropriate areas.

We begin with formulas. Here we have formulas of both second order arithmetic and set theory. We use  $\Sigma_n^0$ ,  $\Pi_n^0$ ,  $\Sigma_n^1$  and  $\Pi_n^1$  in the usual way for the formulas of second order arithmetic with the understanding (typical in reverse mathematics) that the second order quantifiers range over subsets of  $\mathbb{N}$  (or equivalently reals) and the formulas may include free set variables (i.e. ones over reals) at no cost to the quantifier complexity. For formulas of set theory, we use the notations  $\Sigma_n$  and  $\Pi_n$  in the usual way with the understanding that at the bottom level we allow bounded quantifiers in the  $\Sigma_0 = \Pi_0$  formulas and may also include free variables.

We will be concerned with sets of reals and not sets of numbers and so define the relevant classes (of sets) only that setting. From the set theoretic point of view we have the usual topologically based Borel hierarchy beginning with the open or closed sets and then progressing by alternating applications of countable intersections and unions. Recursion theoretically, the open (closed) sets are those defined by  $\Sigma_1^0$  ( $\Pi_1^0$ ) formulas where the free set variables other than the one representing the reals in the set being defined are replaced by specific real parameters. (It is worth noting that the closed sets are precisely those of the form  $[T]$  for some tree  $T$ .) One then moves up the Borel hierarchy as usual. We are only concerned with the first few levels and, in particular, the  $\mathbf{\Pi}_3^0$  sets are those  $A$  of the form  $\cap A_k$  where the  $A_k \in \mathbf{\Sigma}_2^0$ , i.e. they are of the form  $A_k = \cup A_{k,j}$  where the  $A_{k,j}$  are  $\mathbf{\Pi}_1^0$ , i.e. closed, sets. They are also the classes of sets of reals defined by formulas at the indicated level of the arithmetic hierarchy with real parameters.

The sets that we will be analyzing from the viewpoint of their determinacy are the finite Boolean combinations of  $\mathbf{\Pi}_3^0$  sets. These are typically laid out in the difference hierarchy of the  $m\text{-}\mathbf{\Pi}_3^0$  sets.

**Definition 2.2.** A set  $A$  (of reals) is  $m\text{-}\mathbf{\Pi}_3^0$  if there are  $\mathbf{\Pi}_3^0$  sets  $A_0, A_1, \dots, A_{m-1}, A_m = \emptyset$  such that

$$x \in A \Leftrightarrow \text{the least } i \text{ such that } x \notin A_i \text{ is odd.}$$

We say that *the sequence*  $\langle A_i | i \leq m \rangle$  *represents*  $A$  (as an  $m\text{-}\mathbf{\Pi}_3^0$  set).

*Remark 2.3.* Without loss of generality, we may assume that the  $A_i$  in Definition 2.2 are nested downward, i.e.  $A_i \supseteq A_{i+1}$  by simply replacing  $A_i$  by  $\bigcap_{j \leq i} A_j$ . We assume that any given representation (as an  $m\text{-}\mathbf{\Pi}_3^0$  set) is of this form.

This hierarchy is called the difference hierarchy since the sets at level  $m$  are generated by taking unions of differences of  $\mathbf{\Pi}_3^0$  sets. More specifically, they are Boolean combinations of  $\mathbf{\Pi}_3^0$  sets of the form  $(A_0 \setminus A_1) \cup (A_2 \setminus A_3) \dots \cup (A_{m-2} \setminus A_{m-1})$  if  $m$  is even and  $(A_0 \setminus A_1) \cup (A_2 \setminus A_3) \dots \cup (A_{m-3} \setminus A_{m-2}) \cup A_{m-1}$  if  $m$  is odd. (We write  $A \setminus B$  for  $A \cap \bar{B}$ .) Of course, when  $m$  is even,  $A_m = \emptyset$  and so  $A_{m-1} = A_{m-1} \setminus A_m$  and we can view it as being of the first form as well. The converse holds as well, i.e. every set  $A$  of the form  $(A_0 \setminus A_1) \cup (A_2 \setminus A_3) \dots \cup (A_{m-2} \setminus A_{m-1})$  is  $m\text{-}\mathbf{\Pi}_3^0$  for  $m$  even and each of the form  $(A_0 \setminus A_1) \cup (A_2 \setminus A_3) \dots \cup A_{m-1}$  is  $m\text{-}\mathbf{\Pi}_3^0$  for  $m$  odd. One can see this as analogous to the difference hierarchy on r.e. sets.

It is easy to see that all finite Boolean combinations of  $\mathbf{\Pi}_3^0$  sets are  $m\text{-}\mathbf{\Pi}_3^0$ . The sets that are a union of differences of  $\mathbf{\Pi}_3^0$  sets are clearly closed under union and include the  $\mathbf{\Pi}_3^0$  sets. On the other hand, if  $A$  is represented by  $\langle A_i | i \leq m \rangle$ , then  $\bar{A}$  is represented by  $\langle 2^\omega, A_i | i \leq m \rangle$ . Thus  $\cup_m m\text{-}\mathbf{\Pi}_3^0$  contains the  $\mathbf{\Pi}_3^0$  sets and is closed under union and complementation.

An advantage of Definition 2.2 as a hierarchy on the finite Boolean combinations of  $\mathbf{\Pi}_3^0$  sets is that it can immediately be extended into the transfinite simply by allowing sequences  $A_i$  for  $i < \alpha$  with the understanding that limit ordinals are even. It is a classical theorem of Kuratowski [1958] that as  $\alpha$  ranges over the countable ordinals these sets are precisely the  $\Delta_4^0$  ones, i.e. those that are both  $\Sigma_4^0$  and  $\Pi_4^0$ . (See MedSalem and Tanaka [2007] for a proof in  $\text{ACA}_0$ .)

In §4, we prove, in  $\Pi_{m+2}^1\text{-CA}_0$ , that all  $m\text{-}\mathbf{\Pi}_3^0$  sets are determined for each  $m \in \mathbb{N}$ . Our negative results in §5, however, use a different, dual representation of the finite Boolean combinations of  $\mathbf{\Pi}_3^0$  sets.

**Definition 2.4.** A set  $A \subseteq 2^\omega$  is  $\mathbf{\Pi}_{3,n}^0$  if there exist  $\mathbf{\Pi}_3^0$  sets  $A_0, \dots, A_n$  such that  $A_n = 2^\omega$  and

$$x \in A \iff \text{the least } i \text{ such that } x \in A_i \text{ is even.}$$

We say that *the sequence*  $\langle A_i | i \leq m \rangle$  *represents*  $A$  (as a  $\mathbf{\Pi}_{3,n}^0$  set).

*Remark 2.5.* Note that without loss of generality we may assume that the  $A_i$  in Definition 2.4 are nested upward, i.e.  $A_i \subseteq A_{i+1}$  by simply replacing  $A_i$  by  $\bigcup_{j \geq i} A_j$ . We assume that any given representation (as a  $\mathbf{\Pi}_{3,n}^0$  set) is of this form.

This formulation does not extend into the transfinite to capture the  $\Delta_4^0$  sets as did that of the  $m\text{-}\mathbf{\Pi}_3^0$  sets. Indeed, it is easy to see that at level  $\omega$  one gets all the  $\Sigma_4^0$  sets represented by sequences  $\langle A_i | i < \omega \rangle$  of  $\mathbf{\Pi}_3^0$  sets. (The issue here is that the  $\mathbf{\Pi}_3^0$  sets are closed under countable intersections and so  $\{x | \forall i < \beta (x \in A_i)\}$  is still  $\mathbf{\Pi}_3^0$  while they are not closed

under countable (as opposed to finite) unions and so  $\{x \mid \exists i < \beta(x \in A_i)\}$  is only  $\Sigma_4^0$ . Thus Definition 2.4 is suited to a difference hierarchy built on the  $\Sigma_3^0$  rather than the  $\Pi_3^0$  sets. Nonetheless, at the finite levels of the two hierarchies are closely enough related to allow us to use both in our calculations of the limits of determinacy provable in fragments of second order arithmetic.

**Lemma 2.6.**  $(RCA_0) \Pi_{3,2n+1}^0 = (2n+1)\text{-}\Pi_3^0$  and  $\check{\Pi}_{3,2n}^0 = (2n)\text{-}\Pi_3^0$ .

PROOF: Consider any sequence  $A_0 \subseteq A_1 \subseteq \dots \subseteq A_{m-1}$  of  $\Pi_3^0$  sets. Let  $A$  be the  $\Pi_{3,m}^0$  set represented by  $\langle A_0, \dots, A_{m-1}, 2^\omega \rangle$ . Let  $B$  be the  $m\text{-}\Pi_3^0$  set represented by  $\langle A_{m-1}, \dots, A_1, \emptyset \rangle$ . We claim that  $A = B$  when  $m$  is odd and  $A = \bar{B}$  when  $m$  is even. This would prove the lemma as  $A$  or  $B$  could be any  $m\text{-}\Pi_3^0$  or  $\Pi_{3,m}^0$  set.

Consider  $x \in 2^\omega$ . Let  $i_x$  be the least  $i$  such that  $x \in A_i$ . So,  $x \in A$  if and only if  $i_x$  is even. Let  $j_x$  be the least  $j$  such that  $x \notin A_{m-1-j}$ . So,  $x \in B$  if and only if  $j_x$  is odd. Since  $i_x - 1$  is the greatest  $i$  such that  $x \notin A_i$ , we have that  $m - 1 - j_x = i_x - 1$ . It then follows that  $i_x$  and  $j_x$  have the same parity if  $m$  is even and different parities if  $m$  is odd. Hence  $x \in A \iff x \notin B$  if  $m$  is even and  $x \in A \iff x \in B$  if  $m$  is odd.  $\square$

**Corollary 2.7.** For each  $m \geq 1$ ,  $RCA_0 \vdash m\text{-}\Pi_3^0\text{-DET} \leftrightarrow \Pi_{3,m}^0\text{-DET}$ .

PROOF: It is immediate that  $(\Pi_1^0)^* \subseteq \Pi_1^0$  and so the same property clearly holds for  $\Pi_3^0$  as it is generated from  $\Pi_1^0$  by unions and intersections which preserves the desired closure property. The required closure property then holds for  $m\text{-}\Pi_3^0$ ,  $\Pi_{3,m}^0$ ,  $m\text{-}\check{\Pi}_3^0$  and  $\check{\Pi}_{3,m}^0$  (simply replace each  $A_i$  in the representation by  $A_i^*$ ). Thus the corollary holds by Lemmas 2.6 and 2.1.  $\square$

We also point out that the classes we have considered all have “light faced” versions, i.e. ones defined by the same classes of formulas but without any reals as parameters. In terms of our definitions, the light faced (or recursive) closed sets are those of the form  $[T]$  for  $T$  a recursive tree. The class of these sets is denoted by  $\Pi_1^0$ . The light faced class  $\Pi_3^0$  is defined from  $\Pi_1^0$  as in the boldface version except that now the closed sets  $A_{k,j}$  required to define  $A \in \Pi_3^0$  must be uniformly recursive, i.e. there is a set of uniformly recursive trees  $T_{k,j}$  (i.e. the relation  $p \in T_{k,j}$  is a recursive one of the triple  $\langle p, k, j \rangle$ ) such that  $A_{k,j} = [T_{k,j}]$ . The light faced  $m\text{-}\Pi_3^0$  and  $\Pi_{3,n}^0$  sets are then defined from the  $\Pi_3^0$  sets exactly as in the bold faced versions in Definitions 2.2 and 2.4, respectively. Of course, each of these light faced classes are contained in the corresponding bold face class. Our negative results about the failure of determinacy in various subsystems of second order arithmetic as exhibited in the particular models of §5 all provide light face counterexamples.

**2.3. Systems of Reverse Mathematics.** We first briefly review the five standard systems of reverse mathematics. For the analysis of even the lowest levels of determinacy one begins near the top of this list and moves well beyond. Details, general background, and results, as well as many examples of reversals, can be found in Simpson [2009], the standard text on reverse mathematics. Each of the systems is given in the (two sorted) language of second order arithmetic, that is, the usual first order language of arithmetic augmented by set variables with their quantifiers and the membership relation  $\in$  between numbers and sets. A structure for this language is one of the form  $\mathcal{M} = \langle M, S, +, \times, <, 0, 1, \in \rangle$  where  $M$  is a set (the set of “numbers” of  $\mathcal{M}$ ) over which the first order quantifiers and variables of

our language range;  $S \subseteq 2^M$  is the collection of subsets of “numbers” in  $\mathcal{M}$  over which the second order quantifiers and variables of our language range;  $+$  and  $\times$  are binary functions on  $M$ ;  $<$  is a binary relation on  $M$  while  $0$  and  $1$  are members of  $M$ .

Each subsystem of second order arithmetic that we consider contains the standard basic axioms for  $+$ ,  $\cdot$ , and  $<$  (which say that  $\mathbb{N}$  is an ordered semiring). In addition, they all include a form of induction that applies only to sets (that happen to exist):

$$(I_0) \quad (0 \in X \wedge \forall n (n \in X \rightarrow n + 1 \in X)) \rightarrow \forall n (n \in X).$$

We call the system consisting of  $I_0$  and the basic axioms of ordered semirings  $P_0$ . All five standard systems are defined by adding various types of set existence axioms to  $P_0$ .

( $RCA_0$ ) Recursive Comprehension Axioms: This is a system just strong enough to prove the existence of the computable sets. In addition to  $P_0$  its axioms include the schemes of  $\Delta_1^0$  comprehension and  $\Sigma_1^0$  induction:

$$\begin{aligned} (\Delta_1^0\text{-CA}_0) \quad & \forall n (\varphi(n) \leftrightarrow \psi(n)) \rightarrow \exists X \forall n (n \in X \leftrightarrow \varphi(n)) \text{ for all } \Sigma_1^0 \text{ formulas} \\ & \varphi \text{ and } \Pi_1^0 \text{ formulas } \psi \text{ in which } X \text{ is not free.} \\ (I\Sigma_1^0) \quad & (\varphi(0) \wedge \forall n (\varphi(n) \rightarrow \varphi(n + 1))) \rightarrow \forall n \varphi(n) \text{ for all } \Sigma_1^0 \text{ formulas } \varphi. \end{aligned}$$

The next system says that every infinite binary tree has an infinite path.

( $WKL_0$ ) Weak König’s Lemma: This system consists of  $RCA_0$  plus the statement that every infinite subtree of  $2^{<\omega}$  has an infinite path.

We next move up to arithmetic comprehension.

( $ACA_0$ ) Arithmetic Comprehension Axioms: This system consists of  $RCA_0$  plus the axioms  $\exists X \forall n (n \in X \leftrightarrow \varphi(n))$  for every arithmetic formula (i.e.  $\Sigma_n^0$  for some  $n$ )  $\varphi$  in which  $X$  is not free.

The next system says that arithmetic comprehension can be iterated along any countable well order. As noted in §1, it is equivalent to  $\Sigma_1^0$ -determinacy in Baire space (Steel [1976]).

( $ATR_0$ ) Arithmetical Transfinite Recursion: This system consists of  $RCA_0$  plus the following axiom. If  $X$  is a set coding a well order  $<_X$  with domain  $D$  and  $Y$  is a code for a set of arithmetic formulas  $\varphi_x(z, Z)$  (indexed by  $x \in D$ ) each with one free set variable and one free number variable, then there is a sequence  $\langle K_x \mid x \in D \rangle$  of sets such that if  $y$  is the immediate successor of  $x$  in  $<_X$ , then  $\forall n (n \in K_y \leftrightarrow \varphi_x(n, K_x))$ , and if  $x$  is a limit point in  $<_X$ , then  $K_x$  is  $\bigoplus \{K_y \mid y <_X x\}$ .

The systems that we actually need climb up to full second order arithmetic,  $Z_2$ , (i.e. comprehension for all formulas). They are characterized by the syntactic level of the second order formulas for which they include comprehension axioms.

( $\Pi_n^1\text{-CA}_0$ ) These systems consists of  $RCA_0$  plus the  $\Pi_n^1$  comprehension axioms:  $\exists X \forall k (k \in X \leftrightarrow \varphi(k))$  for every  $\Pi_n^1$  formula  $\varphi$  in which  $X$  is not free.

The first of these,  $\Pi_1^1\text{-CA}_0$ , is the last of the five standard systems. As noted in §1, it is equivalent to  $(\Sigma_1^0 \wedge \Pi_1^0)$ -determinacy in Baire space (Tanaka [1990]). If one assumes  $\Pi_n^1\text{-CA}_0$  for every  $n$  then one has full second order arithmetic.

( $\Pi_\infty^1\text{-CA}_0$ ) or ( $Z_2$ ) This system consists of  $\Pi_n^1\text{-CA}_0$  for every  $n$ .

We also need the intermediate  $\Delta_n^1$  classes and corresponding axioms.

( $\Delta_n^1$ -CA<sub>0</sub>)  $\Delta_n^1$  Comprehension Axioms:  $\forall k (\psi(k) \leftrightarrow \neg\varphi(k)) \rightarrow \exists X \forall k (k \in X \leftrightarrow \varphi(k))$  for every  $\Sigma_n^1$  formulas  $\varphi$  and  $\psi$  in which  $X$  is not free.

The other set existence axioms that play a role in our analysis are various versions of the axiom of choice in the setting of second order arithmetic.

( $\Sigma_n^1$ -AC<sub>0</sub>) This system consists of ACA<sub>0</sub> plus the  $\Sigma_n^1$  choice axioms  $\forall n \exists Y \varphi(n, Y) \rightarrow \exists Z \forall n \varphi(n, (Z)_n)$  for every  $\Sigma_n^1$  formula  $\varphi$  and where  $(Z)_n = \{i \mid \langle i, n \rangle \in Z\}$ .

( $\Sigma_n^1$ -DC<sub>0</sub>) This system consists of ACA<sub>0</sub> plus the  $\Sigma_n^1$  dependent choice axioms  $\forall n \forall X \exists Y \varphi(n, X, Y) \rightarrow \exists Z \varphi(n, (Z)^n, (Z)_n)$  for every  $\Sigma_n^1$  formula  $\varphi$  and where  $(Z)^n = \{\langle i, m \rangle \mid \langle i, m \rangle \in Z \ \& \ m < n\}$

(Strong  $\Sigma_n^1$ -DC<sub>0</sub>) This system consists of ACA<sub>0</sub> plus the strong  $\Sigma_n^1$  dependent choice axioms  $\exists Z \forall n \forall Y (\varphi(n, (Z)^n, Y) \rightarrow \varphi(n, (Z)^n, (Z)_n))$  for every  $\Sigma_n^1$  formula  $\varphi$ .

See Simpson [2009, VII.6] for the relations among these choice principles and various comprehension axioms.

If we strengthen the basic induction axiom  $I_0$  by replacing it with induction for all formulas  $\varphi$  of second order arithmetic we get full induction

(I)  $(\varphi(0) \wedge \forall n (\varphi(n) \rightarrow \varphi(n+1))) \rightarrow \forall n (\varphi(n))$  for every formula  $\varphi$  of second order arithmetic.

Each of the systems above has an analog in which  $I_0$  is replaced by I. It is designated by the same letter sequence as above but without the subscript 0, as for example, RCA in place of RCA<sub>0</sub>.

**Definition 2.8.** If  $\mathcal{M}$  is a structure for second order arithmetic and its first order part  $M$  is  $\mathbb{N}$  then  $\mathcal{M}$  is an  $\omega$ -model.

Obviously if an  $\omega$ -model  $\mathcal{M}$  is a model of one of the systems above, such as  $\Pi_n^1$ -CA<sub>0</sub>, then it is also a model of the analogous system, such as  $\Pi_n^1$ -CA.

Restricted versions of the full induction axioms are designated as follows:

( $\Sigma_n^1$ -IND)  $(\varphi(0) \wedge \forall n (\varphi(n) \rightarrow \varphi(n+1))) \rightarrow \forall n (\varphi(n))$  for every  $\Sigma_n^1$  formula  $\varphi$  of second order arithmetic.

**Definition 2.9.** An  $\omega$ -model  $\mathcal{M}$  is a  $\beta$ -model if, for every  $\Sigma_1^1$  sentence  $\varphi$  with parameters in  $\mathcal{M}$ ,  $\varphi \Leftrightarrow \mathcal{M} \models \varphi$ .

In particular, if  $\mathcal{M}$  is a  $\beta$ -model then any well ordering in  $\mathcal{M}$ , i.e. a linear ordering in  $S$  with no infinite descending chain in  $S$ , is actually a well ordering (where  $S$  is the second order part of  $\mathcal{M}$ ). There are also some induction axioms and systems (even stronger than  $\Sigma_n^1$ -IND<sub>0</sub>) for transfinite procedures that serve as proof theoretical approximations to being a  $\beta$ -model.

( $\Pi_\infty^1$ -TI<sub>0</sub>) This system consists of ACA<sub>0</sub> plus all the transfinite induction axioms  $\forall X (X \text{ is a well-ordering} \ \& \ \forall j (\forall i (i <_X j \rightarrow \psi(i)) \rightarrow \psi(j)) \rightarrow \forall j \psi(j)$ , where by “ $X$  is a well-ordering” we mean that the relation  $<_X = \{\langle i, j \rangle \mid \langle i, j \rangle \in X\}$  given by the pairs of numbers in  $X$  is a well ordering, i.e. has no infinite descending chains, and  $\psi$  is any formula of second order arithmetic in which  $X$  is not free.

Obviously, every  $\beta$ -model satisfies  $\Pi_\infty^1$ -TI<sub>0</sub>. The systems  $(\Pi_n^1$ -TI<sub>0</sub>) are defined analogously with  $\psi$  restricted to  $\Pi_n^1$  formulas.

## 3. FACTS ABOUT L

We assume a familiarity with the basic facts about  $L$  and  $\Sigma_1$  admissible sets of the form  $L_\alpha$  as can be found, for example, in Devlin [1984, I-II] or Barwise [1975, I-II]. Generalizing from 1 to  $n$  we say that an  $L_\alpha$  is  $\Sigma_n$  *admissible* if it satisfies  $\Sigma_n$  bounding: For any  $\delta < \alpha$  and any  $\Pi_{n-1}/L_\alpha$  formula  $\varphi(x, y)$ , if  $L_\alpha \models \forall \gamma < \delta \exists y \varphi(\gamma, y)$ , then there is a  $\lambda < \alpha$  such that  $L_\alpha \models \forall \gamma < \delta \exists y \in L_\lambda \varphi(\gamma, y)$ . This property is equivalent (for initial segments of  $L$ ) to the requirements that the set satisfy (in addition to the standard (KP) axioms of admissibility)  $\Delta_{n-1}$  collection and  $\Sigma_{n-1}$  separation (also called comprehension). It is also equivalent to there not being a function  $f$  with domain some  $\delta < \alpha$  which is  $\Sigma_n$  (or equivalently  $\Pi_{n-1}$ ) over  $L_\alpha$  whose range is unbounded in  $\alpha$  or  $L_\alpha$ . One more equivalent is that for any  $\Sigma_n/L_\alpha$  function  $f$ ,  $f \upharpoonright \gamma \in L_\alpha$  for every  $\gamma < \alpha$ . Thus we have the usual bounded quantifier manipulation rules: for  $\varphi \in \Pi_{n-1}$ ,  $\forall x \in t \exists y \varphi$  is equivalent to a  $\Sigma_n$  formula. We also say that  $\alpha$  is  $n$ -*admissible* if  $L_\alpha$  is  $\Sigma_n$  admissible.

We need a few facts about  $\Sigma_n$  Skolem functions and projections which are deep and difficult theorems of Jensen's fine structure theory in general (as in Devlin [1984, VI]) but become quite simple via standard arguments when enough admissibility is assumed. We also need a couple of special facts about the ordinals less than  $\beta_0$ , the first ordinal which is  $\Sigma_n$  admissible for all  $n$ . All of these ought to be at least known folklore but seem to be missing from the couple of standard texts we consulted.

**Proposition 3.1.** *If  $\alpha$  is  $\Sigma_k$  admissible then  $L_\alpha$  has a parameterless  $\Sigma_{k+1}$  Skolem function.*

PROOF: For a  $\Sigma_{k+1}$  formula  $\exists x \varphi(x, \bar{y})$  with Gödel number  $n$  we let  $f(n, \bar{y}) = x \Leftrightarrow \varphi(x, \bar{y}) \wedge (\forall x' <_L x)(\neg \varphi(x', \bar{y}))$ . The first conjunct is  $\Pi_k$  while the second one is equivalent to a  $\Sigma_k$  formula by  $\Sigma_k$  admissibility.  $\square$

**Lemma 3.2.** *If  $L_\alpha$  is  $\Sigma_n$  admissible then it satisfies  $\Delta_n$  comprehension, i.e. for any  $u \in L_\alpha$  and  $\Sigma_n$  formulas  $\varphi(z)$  and  $\psi(z)$  such that  $L_\alpha \models \forall z(\varphi(z) \leftrightarrow \neg \psi(z))$ ,  $\{z \in u \mid \varphi(z)\} \in L_\alpha$ .*

PROOF: For  $n = 1$  this is a standard fact. We now proceed by induction on  $n \geq 2$ . Consider any  $\Sigma_{n-2}$  formulas  $\varphi(x, y, z)$  and  $\psi(x, y, z)$  such that  $L_\alpha \models \forall z(\exists x \forall y \varphi(x, y, z) \leftrightarrow \neg \exists x \forall y \psi(x, y, z))$  and any  $u \in L_\alpha$ . We must show that  $\{z \in u \mid L_\alpha \models \exists x \forall y \varphi(x, y, z)\} \in L_\alpha$ . Define in  $L_\alpha$  a function on every  $z \in u$  by  $f(z) = x \Leftrightarrow [\forall y \varphi(x, y, z) \wedge (\forall x' <_L x) \exists y' \neg \varphi(x', y', z)] \vee [\forall y \psi(x, y, z) \wedge (\forall x' <_L x) \exists y' \neg \psi(x', y', z)]$ . By  $\Sigma_n$  admissibility, the formula defining  $f$  is equivalent to one that is  $\Delta_n$  and so its range is bounded, say by  $L_\gamma$ . We then have that  $\{z \in u \mid L_\alpha \models \exists x \forall y \varphi(x, y, z)\} = \{z \in u \mid L_\alpha \models \exists x \in L_\gamma \forall y \varphi(x, y, z)\} = \{z \in u \mid L_\alpha \models \forall x \in L_\gamma \exists y \neg \psi(x, y, z)\}$ . Again by  $\Sigma_n$  admissibility these last two definitions are equivalent to ones that are  $\Pi_{n-1}$  and  $\Sigma_{n-1}$ , respectively and so by  $\Delta_{n-1}$  comprehension which we have by induction, they belong to  $L_\alpha$  as required.  $\square$

**Lemma 3.3.** *If  $\alpha$  is  $(k+1)$ -admissible, then there are unboundedly many  $\gamma < \alpha$  such that  $L_\gamma \preceq_k L_\alpha$ . (We write  $M \preceq_k N$  to denote the assertion that  $M$  is a  $\Sigma_k$  elementary submodel of  $N$ , i.e. for any  $\Sigma_k$  formula  $\varphi$  with parameters in  $M$ ,  $M \models \varphi \Leftrightarrow N \models \varphi$ .)*

PROOF: Consider any  $\delta < \alpha$ . We want to show there is a  $\gamma$  between  $\delta$  and  $\alpha$  such that  $L_\gamma \preceq_k L_\alpha$ . Let  $h$  be a (parameterless)  $\Sigma_k$ -Skolem function for  $L_\alpha$ . Let  $H_1$  be the  $\Sigma_k$ -Skolem hull of  $L_\delta$  in  $L_\alpha$ . Since  $\alpha$  is  $(k+1)$ -admissible,  $H_1$  is bounded in  $L_\alpha$ ; let  $\delta_1 < \alpha$  be the least ordinal such that  $H_1 \subseteq L_{\delta_1}$ . We now iterate this process. Let  $H_l$  be the  $\Sigma_k$ -Skolem hull of

$L_{\delta_{l-1}}$  in  $L_\alpha$ , and let  $\delta_l$  be the least ordinal such that  $H_l \subseteq L_{\delta_l}$ . As for  $\delta_1$ , one can show that all the  $\delta_l$  are  $< \alpha$ . Let  $\gamma = \sup_{l \in \omega} \delta_l$ . First, let us show that  $L_\gamma \preceq_k L_\alpha$ . Suppose that  $i < k$  and that we know that  $L_\gamma \preceq_i L_\alpha$ ; we claim that  $L_\gamma \preceq_{i+1} L_\alpha$ . Let  $\varphi$  be a  $\Pi_i$  formula with parameters in  $L_{\delta_l}$  for some  $l$  such that  $L_\alpha \models \exists x \varphi(x)$ . Then, there exists an  $x \in H_{l+1}$  such that  $L_\alpha \models \varphi(x)$ , and since  $L_\gamma \preceq_i L_\alpha$  we have that  $L_\gamma \models \varphi(x)$ . This proves our claim, and hence  $L_\gamma \preceq_k L_\alpha$ .

Next, we need to show that  $\gamma < \alpha$ . The reason is that the function  $l \mapsto \delta_l: \omega \rightarrow L_\alpha$  is  $\Sigma_{k+1}$ :

$$\delta_l = \mu\beta(\exists\beta_0, \dots, \beta_l \leq \beta(\beta_0 = \delta \wedge \beta_l = \beta \wedge \bigwedge_{i < l} \forall x \in L_{\beta_i}(h(x) \in L_{\beta_{i+1}}))).$$

Since  $\alpha$  is  $(k+1)$ -admissible and  $h$  is  $\Sigma_k$ , the formula in parenthesis is  $\Sigma_k$ . Finding the least such  $\beta$  is  $\Delta_{k+1}$ . Therefore, again by  $(k+1)$ -admissibility, this function has to be bounded below  $\alpha$ , and hence  $\gamma = \sup_{l \in \omega} \delta_l < \alpha$ .  $\square$

**Lemma 3.4.** *If  $L_\gamma \preceq_k L_\beta$  and  $\beta$  is  $\Sigma_{k-1}$  admissible then  $\gamma$  is  $\Sigma_k$  admissible.*

PROOF: We proceed by induction on  $k \geq 1$ . Consider  $k$  and  $L_\gamma \preceq_k L_\beta$  with  $\beta$   $\Sigma_{k-1}$  admissible. (By convention,  $\Sigma_0$  admissibility is just transitivity and so every  $L_\beta$  is  $\Sigma_0$  admissible.) If the Lemma fails, there is a  $\Pi_{k-1}/L_\gamma$  function  $f$  with domain some  $\delta < \gamma$  and range unbounded in  $L_\gamma$ . Suppose, for  $\eta < \delta$ ,  $f(\eta) = y \Leftrightarrow \psi(\eta, y)$  where  $\psi$  is  $\Pi_{k-1}/L_\gamma$ . For each  $\eta < \delta$  and  $y = f(\eta)$ ,  $L_\beta \models \psi(\eta, y)$  since  $L_\gamma \preceq_k L_\beta$ . Thus  $L_\beta \models \exists x \forall \eta < \delta \exists y <_L x(\psi(\eta, y))$ . By the usual rules for manipulating bounded quantifiers, there is a  $\Pi_{k-1}$  formula  $\theta(x)$  which is equivalent to  $\forall \eta < \delta \exists y <_L x(\psi(\eta, y))$  in any  $\Sigma_{k-1}$  admissible  $L_\zeta$  for  $\zeta \geq \gamma$ . As  $L_\beta$  is  $\Sigma_{k-1}$  admissible by assumption,  $L_\beta \models \exists x \theta(x)$ . As  $L_\gamma \preceq_k L_\beta$ ,  $L_\gamma \models \exists x \theta(x, \eta, y)$ . Finally, as  $L_\gamma$  is  $\Sigma_{k-1}$  admissible by induction,  $L_\gamma \models \exists x \forall \eta < \delta \exists y <_L x(\psi(\eta, y))$  for the desired contradiction.  $\square$

**Lemma 3.5.** *For  $\gamma < \beta_0$ ,  $L_\gamma$  is countable in  $L_{\gamma+1}$ .*

PROOF: For  $\gamma = \omega$ , the conclusion is immediate. We now proceed by induction on  $\gamma$ . For a successor ordinal  $\gamma + 1$ , the conclusion follows simply from the countability of the formulas paired with the (inductively given) countability of the parameter space  $L_\gamma$ . If  $\gamma$  is a limit ordinal less than  $\beta_0$ , it is not  $\Sigma_n$  admissible for some  $n$ , i.e. there is a  $\Sigma_n/L_\gamma$  map  $f$  from some  $\delta < \gamma$  with range unbounded in  $\gamma$ . We now define a counting of  $L_\gamma$  by combining the countings of  $\delta$  and  $L_{f(\zeta)}$  for each  $\zeta < \delta$  (which are uniformly definable as they are each in  $L_{f(\zeta)+1}$  by induction).  $\square$

**Lemma 3.6.** *Every member of  $L_{\alpha_n}$  is  $\Sigma_{n+1}$  definable over  $L_{\alpha_n}$  without parameters.*

PROOF: By Lemma 3.1 we may take  $f$  to be a parameterless  $\Sigma_{n+1}$  Skolem function for  $L_{\alpha_n}$ . Let  $H$  be the Skolem hull of the empty set under  $f$ . As  $L_{\alpha_n}$  is not  $(n+1)$ -admissible (by Lemmas 3.3 and 3.4),  $H$  is unbounded in  $L_{\alpha_n}$ . On the other hand, By Lemma 3.5,  $H$  is transitive. (Say  $\delta \in H$  and so a counting of  $\delta$  also belongs to  $H$  but then so does the value of this counting on every  $k \in \omega$ .) Thus  $H = L_{\alpha_n}$  and so every element of  $L_{\alpha_n}$  is  $\Sigma_{n+1}$  definable (as the value of  $f$  on some  $k \in \omega$ ) over  $L_{\alpha_n}$  without parameters.  $\square$

#### 4. PROVING $m$ - $\Pi_3^0$ DETERMINACY IN $\Pi_{m+2}^1$ -CA $_0$

This section is devoted to proving Theorem 1.1: For each  $m \geq 1$ ,  $\Pi_{m+2}^1$ -CA $_0 \vdash m$ - $\Pi_3^0$ -DET. Our proof is a variant of Martin's proof [1974], [n.d., Ch. 1] of  $\Delta_4^0$ -DET specialized

and simplified to the finite levels of the difference hierarchy on  $\mathbf{\Pi}_3^0$ . Our proof of the main Lemma (Lemma 4.9) is perhaps somewhat different from his. In addition, we supply the analysis needed to see precisely how much comprehension is needed at every level.

We fix  $m \in \omega$ ,  $m \geq 1$ . As  $m$ - $\mathbf{\Pi}_3^0$  determinacy is a  $\mathbf{\Pi}_3^1$  sentence and strong  $\Sigma_{m+2}^1$ -DC<sub>0</sub> is  $\mathbf{\Pi}_4^1$  conservative over  $\mathbf{\Pi}_{m+2}^1$ -CA<sub>0</sub> by Simpson [2009, VII.6.20], we assume strong  $\Sigma_{m+2}^1$ -DC<sub>0</sub> as well as  $\mathbf{\Pi}_{m+2}^1$ -CA<sub>0</sub>. In particular, we fix an  $m$ - $\mathbf{\Pi}_3^0$  set  $A$  given by the nested sequence  $A_0 \supseteq A_1 \supseteq \dots \supseteq A_m = \emptyset$  of  $\mathbf{\Pi}_3^0$  sets. We let  $A_i = \cap A_{i,k}$  for  $A_{i,k} \in \Sigma_2^0$  and  $A_{i,k} = \cup A_{i,k,j}$  for  $A_{i,k,j} \in \mathbf{\Pi}_1^0$ .

We note that this proof does not use any determinacy beyond open determinacy, that (as we have seen) can be proved in systems much weaker than those we are using here.

We consider strings  $s \in \omega^{\leq m}$  (so of length at most  $m$ ) and subgame trees  $S$  of the full binary tree. When  $s$  is specified, we let  $l = m - |s|$  and frequently make definitions and proofs depend on the parity of  $l$  (or equivalently of  $m$  when  $s = \emptyset$  or  $|s|$  is otherwise fixed).

**Definition 4.1.** We define relations  $P^s(S)$  by induction on  $|s| \leq m$ :

- (1)  $P^\emptyset(S) \Leftrightarrow$  there is a winning strategy for I (II) if  $l$  (or equivalently  $m$ ) is even (odd) in  $G(A, S)$ .

For  $|s| = n + 1$  we also divide into cases based on the parity of  $l$ .

- (1)
- (2) If  $l$  is even,  $P^s(S) \Leftrightarrow$  there is a quasistrategy  $U$  for I in  $S$  such that
  - (a)  $[U] \subseteq A \cup A_{m-n-1, s(n)}$  and
  - (b)  $P^{s \uparrow n}(U)$  fails.
- (3) If  $l$  is odd,  $P^s(S) \Leftrightarrow$  there is a quasistrategy  $U$  for II in  $S$  such that
  - (a)  $[U] \subseteq \bar{A} \cup A_{m-n-1, s(n)}$  and
  - (b)  $P^{s \uparrow n}(U)$  fails.

A quasistrategy  $U$  witnesses  $P^s(S)$  if  $U$  is as required in the appropriate clause ((1), (2) or (3)) of the above definition.

It is not hard to show that for any  $s$ , if  $l$  is even (odd), and I (II) has a winning strategy in  $G(A, S)$ , then  $P^s(S)$  holds. The converse is also true, but we would need to use  $m$ - $\mathbf{\Pi}_3^0$  determinacy to prove it. Our first goal is to prove that for every  $s$  of length  $n + 1$  and any  $S$ , either  $P^s(S)$  holds or  $P^{s \uparrow n}(S)$  holds (Lemma 4.9). Then, in the proof of Theorem 1.1, we will use this result to show that, if  $P^\emptyset(T)$  does not hold, we can build a winning strategy for II (I) in the case when  $m$  is even (odd).

**Definition 4.2.** A quasistrategy  $U$  locally witnesses  $P^s(S)$  if,  $|s| = n + 1$  and  $U$  is a quasistrategy for I (II) if  $l$  is even (odd) and there is a  $D \subseteq S$  such that for every  $d \in D$  there is a quasistrategy  $R^d$  for II (I) if  $l$  is even (odd) in  $S_d$  such that

- (i)  $\forall d \in D \cap U(U_d \cap R^d$  witnesses  $P^s(R^d)$ )
- (ii)  $[U] \setminus \cup_{d \in D} [R^d] \subseteq A (\bar{A})$  if  $l$  is even (odd)
- (iii)  $\forall p \in S \exists \leq 1 d \in D (d \subseteq p \ \& \ p \in R^d)$

*Remark 4.3.* Simple quantifier counting shows that the relations  $P^s(S)$ ,  $U$  witnesses  $P^s(S)$  and  $U$  locally witnesses  $P^s(S)$  are  $\Sigma_{|s|+2}^1$ ,  $\mathbf{\Pi}_{|s|+1}^1$  and  $\Sigma_{|s|+2}^1$ , respectively.

We now prove a number of Lemmas in  $\Pi_{m+2}^1\text{-CA}_0 + \text{strong } \Sigma_{m+2}^1\text{-DC}_0$  to establish our theorem. Note that when we perform inductions on something like  $|s|$  we do not have to worry about what induction axioms we need as  $|s| \leq m \in \omega$ .

**Lemma 4.4.** *If  $U$  locally witnesses  $P^s(S)$  then  $U$  witnesses  $P^s(S)$ .*

PROOF: Suppose  $l$  is even. We show that clause (2a) of Definition 4.1 is satisfied. Consider  $x \in [U]$ . If  $x \in A$  there is nothing to prove. If not, by Definition 4.2(ii),  $x \in [R^d]$  for some  $d \in D$ . Then by (i)  $U_d \cap R^d$  witnesses  $P^s(R^d)$  and so by Definition 4.1 (2a)  $x \in A_{m-n-1, s(n)}$  as required. If  $l$  is odd, the same argument with  $\bar{A}$  in place of  $A$  and clause (3a) in place of (2a) shows that clause (3a) is satisfied as required.

We now proceed by induction on  $|s| = n+1 \leq m$  to show that part (b) of clause (2) or (3) of Definition 4.1 holds. We begin with  $n = 0$ . Suppose that  $m$  is odd and so the associated  $l$  with  $s$  in the definitions of  $P^s(S)$  and  $U$  locally witnessing  $P^s(S)$  is even. If (2b) does not hold, i.e.  $P^0(U)$  holds, there is a winning strategy  $\tau$  for II in  $G(A, U)$  (as  $m$  is odd). We claim there is a  $d \in D$  consistent with  $\tau$  such that every  $x \supseteq d$  in  $[U]$  and consistent with  $\tau$  is in  $[R^d]$ . Suppose not. Now note that every  $e \notin \cup_{d \in D, d \subseteq e} R^d$  in  $U$  and consistent with  $\tau$  has a  $\hat{d} \supseteq e$  in  $D$  (and so a minimal one) consistent with  $\tau$ , for if not then, for any  $x \supseteq e$  in  $[U]$  and consistent with  $\tau$ ,  $x \notin \cup_{d \in D} R^d$ . By Definition 4.2(ii) we would then have  $x \in A$  contradicting our choice of  $\tau$  as a winning strategy for II in  $G(A, U)$ . Next, note that, by our assumption, any such  $\hat{d}$  has a (minimal) extension  $\hat{e}$  consistent with  $\tau$  and not in  $R^{\hat{d}}$ . By Definition 4.2(iii), no  $e' \supset \hat{d}$  with  $e' \subset \hat{e}$  is in  $D$  and so  $\hat{e}$  has the same properties assumed about  $e$ . We can continue this procedure to build a sequence  $e_j$  each consistent with  $\tau$  such that  $\cup e_j = x \notin \cup_{d \subseteq x, d \in D} R^d$  and so  $x \in U \setminus \cup_{d \in D} R^d \subseteq A$  for the same contradiction to our choice of  $\tau$ .

So we have a  $d \in D$  consistent with  $\tau$  such that every  $x \supseteq d$  in  $[U]$  and consistent with  $\tau$  is in  $[R^d]$ . Thus the natural restriction of  $\tau$  is a winning strategy for II in  $G(A, U_d \cap R^d)$ , i.e.  $P^0(U_d \cap R^d)$  contradicting Definition 4.2(i) and so establishing (2b) of Definition 4.1 as required.

If  $m$  is even and so the associated  $l$  for  $s$  in the definition of  $P^s(S)$  is odd, the same proof works with the following pairs interchanged: I and II; 2 and 3;  $A$  and  $\bar{A}$  (except that  $G(A, U)$  is unchanged).

Next, we consider  $n > 1$  and assume for definiteness that  $m - n (= m - |s \upharpoonright n|)$  is odd and, for the sake of a contradiction, that  $P^{s \upharpoonright n}(U)$  holds. Let  $\hat{S}$  be II's quasistrategy in  $U$  witnessing  $P^{s \upharpoonright n}(U)$ . We will build  $\hat{U}$ ,  $\hat{D}$  and  $\{\hat{R}^d : d \in \hat{D}\}$  locally witnessing  $P^{s \upharpoonright n-1}(\hat{S})$ . By the induction hypothesis this would imply  $P^{s \upharpoonright n-1}(\hat{S})$ , contradicting that  $\hat{S}$  witnesses  $P^{s \upharpoonright n}(U)$ . Roughly speaking, to build I's quasistrategy  $\hat{U}$ , player I would try to fall out of  $\cup_{d \in D} [R^d]$  and hence end up in  $A$ . For the  $d \in D$  such that I has no strategy to exit  $R^d$ , we will add  $d \in \hat{D}$  and define  $\hat{R}^d \subseteq R^d$  such that  $P^{s \upharpoonright n-1}(\hat{R}^d)$ .

Let  $\hat{D} = \{d \in \hat{S} \cap D \mid G(\overline{[R^d]}, \hat{S}_d) \text{ is a win for II}\}$  (a  $\Sigma_2^1$  set). For  $d \in \hat{D}$  we let  $\hat{R}^d$  be II's nonlosing quasistrategy in  $G(\overline{[R^d]}, \hat{S}_d)$  (each a  $\Pi_2^1$  set and so the indexed collection is given by  $\Pi_3^1\text{-CA}_0$ ) so, of course,  $\hat{R}^d \subseteq R^d$  and  $\hat{R}^d$  is a quasistrategy for II in  $U_d$ . As  $\hat{S}$  witnesses  $P^{s \upharpoonright n}(U)$ ,  $[\hat{S}] \subseteq \bar{A} \cup A_{m-n, s(n-1)}$  and so  $[\hat{R}^d] \subseteq \bar{A} \cup A_{m-n, s(n-1)}$ . By Definition 4.2(i),  $U_d \cap R^d$  witnesses  $P^s(R^d)$  and so, in particular,  $P^{s \upharpoonright n}(U_d \cap R^d)$  fails and  $\hat{R}^d$  is not a witness for it. As condition (3a) has already been verified for  $[U] \supseteq [\hat{R}^d]$ , (3b) must fail with  $\hat{R}^d$  and so

there is a witness  $\hat{U}^d$  for each  $P^{s \upharpoonright n-1}(\hat{R}^d)$ . The indexed sequence  $\{\hat{U}^d : d \in \hat{D}\}$  then exists by  $\Sigma_{|s|}^1\text{-AC}_0$  by Remark 4.3. Finally, we use  $\Sigma_2^1\text{-CA}_0$  and  $\Sigma_2^1\text{-AC}_0$  to define the sequence  $\sigma_{q,d}$  of winning strategies for I in  $G(\overline{[R^d]}, \hat{S}_q)$  for  $d \in D$  and  $q \notin \hat{R}^d$  (which is taken to be  $\emptyset$  for  $d \in D \setminus \hat{D}$ ) when one exists.

We now (arithmetically in the above parameters) define a quasistrategy  $\hat{U}$  for I in  $\hat{S}$ :

- (1) If  $p \in \hat{U}$  and there is no  $d \in D$  such that  $d \subseteq p$  and  $p \in R^d$  then the immediate extensions of  $p$  in  $\hat{U}$  are the same as those in  $\hat{S}$ .
- (2) For each  $q \in \hat{U}$  and  $d \in D$  such that  $q$  is a minimal extension of  $d$  in  $R^d \setminus \hat{R}^d$ , the play in  $\hat{U}$  following  $q$  agrees with that of  $\sigma_{q,d}$  until we reach a  $p \notin R^d$ .
- (3) For  $d \in \hat{D} \cap \hat{U}$ , let  $\hat{U}_d \cap \hat{R}^d = \hat{U}^d$ .

We claim that  $\hat{U}$  (with  $\hat{D}$  and  $\hat{R}^d$ ) locally witnesses  $P^{s \upharpoonright n-1}(\hat{S})$ . Property (i) of Definition 4.2 follows from (3) in the definition of  $\hat{U}$  and our choice of  $\hat{U}^d$  for  $d \in \hat{D}$ . For (ii), note that, by (2) in the definition of  $\hat{U}$ ,  $[\hat{U}] \cap [R^d] \subseteq [\hat{R}^d]$  while  $\hat{U}$  is a quasistrategy in  $\hat{S}$  which is one in  $U$ . Thus  $[\hat{U}] \setminus \cup_{d \in D} [\hat{R}^d] \subseteq [U] \setminus \cup_{d \in D} [R^d] \subseteq A$  by Definition 4.2(ii) as required. Property (iii) is immediate from the definitions and the corresponding condition for  $U$  being a local witness for  $P^s(S)$  as  $\hat{D} = D \cap \hat{S}$  and  $\hat{R}^d \subseteq R^d$ . Thus by induction,  $\hat{U}$  witnesses  $P^{s \upharpoonright n-1}(\hat{S})$  for the desired contradiction (with clause (3b) in the definition of  $\hat{S}$  being a witness for  $P^{s \upharpoonright n(U)}$ ).

Finally, when  $n = 1$  (still assuming that  $m - n$  is odd), the proof is much the same as when  $n > 1$  except that when we choose a witness  $\hat{U}^d$  for  $P^{s \upharpoonright n-1}(\hat{R}^d)$  we must take a winning strategy for I in  $G(A, \hat{R}^d)$  and we need to show that  $[\hat{U}] \subseteq A$  to see that it witnesses  $P^\emptyset(\hat{S})$  for the desired contradiction (with clause (3b) in the definition of  $\hat{S}$  being a witness for  $P^{s \upharpoonright n(U)}$ ). The point here is that if we stay in some  $\hat{R}^d$  then we follow  $\hat{U}^d$  which is a winning strategy for I in  $G(A, \hat{R}^d)$ . If we leave  $\hat{R}^d$  then we leave  $R^d$  by (2) in the definition of  $\hat{U}$ . If we leave every  $R^d$ , we follow  $\hat{S}$  and so stay in  $U$  and also wind up  $A$  by clause (ii) of the definition of  $U$  being a local witness for  $P^s(S)$ .

For  $m - n$  even, the proof is the same except that we interchange I and II, clauses (2) and (3) of Definition 4.1,  $A$  and  $\bar{A}$  and  $G(\overline{[R^d]}, \hat{S}_d)$  and  $G([R^d], \hat{S}_d)$ .  $\square$

**Definition 4.5.**  $P^s(S)$  fails everywhere if  $P^s(S_p)$  fails for every  $p \in S$ . By Remark 4.3, this relation is  $\Pi_{|s|+2}^1$ .

**Lemma 4.6.** *If  $P^s(S)$  fails then there is a quasistrategy  $W$  for I if  $l$  is odd (for II if  $l$  is even) in  $S$  such that  $P^s(W)$  fails everywhere.*

**PROOF:** For  $|s| = 0$ , if  $l$  is odd (even), II (I) does not have a winning strategy in  $G(A, S)$ . Let  $W$  be I's (II's) nonlosing quasistrategy in this game. It exists by  $\Pi_2^1\text{-CA}_0$  and, by definition, II (I) does not have a winning strategy in  $G(A, W_p)$  for any  $p \in W_p$  and so  $P^s(W)$  fails everywhere.

Now suppose  $|s| = n+1$  and, for definiteness, that  $l$  is even. Let  $D$  be the  $\Sigma_{|s|+2}^1 \wedge \Pi_{|s|+2}^1$  set of minimal  $d \in S$  for which  $P^s(S_d)$ . (If there are none, then  $P^s(S)$  already fails everywhere.) By  $\Sigma_{|s|+2}^1\text{-AC}_0$  let  $U^d$  witness  $P^s(S_d)$  for every  $d \in D$ .

Consider now the game  $G(B, S)$  where  $B = \{x \in [S] \mid \exists d \in D (d \subseteq x)\}$ . We claim that I has no winning strategy in this game. If there were one  $\sigma$ , then we could define a quasistrategy  $U$  for I in  $S$  by following  $\sigma$  until a position  $d \in D$  is reached at which point we move into

$U^d$ . With  $D$  and  $R^d = S_d$  for  $d \in D$ , it is easy to see that  $U$  locally witnesses  $P^s(S)$  which by the previous Lemma is the desired contradiction.

Thus we can let  $W$  be II's nonlosing quasistrategy in  $G(B, S)$  and  $\sigma_p$  be a winning strategy for I for  $p \in S \setminus W$ . (These exist as above.) If  $W$  is not as required, there is a  $q \in W$  and a witness  $\hat{U}$  that  $P^s(W_q)$ . Define a quasistrategy  $U$  for I in  $S_q$  by  $U \cap W_q = \hat{U}$ . If we ever reach a position  $p \notin W$ , we switch to following  $\sigma_p$  until we reach a position  $d \in D$  at which point we switch to  $U^d$ . If we now let  $\hat{D} = D \cup \{q\}$ ,  $R^q = W_q$  and  $R^d = S_d$  for  $d \in D$ , it is easy to see that together with  $U$  we have a local witness for  $P^s(S_q)$ . By the previous Lemma then  $P^s(S_q)$  and so by the definition of  $D$  there is a  $p \subseteq q$  in  $D$  contradicting the choice of  $W$  as a nonlosing strategy for II in  $G(B, S)$ .

The argument for  $s = n + 1$  and odd  $l$  is the same except that we interchange I and II and  $G(B, S)$  and  $G(\bar{B}, S)$ .  $\square$

**Definition 4.7.** For  $n + 1 = |s|$ ,  $W$  *strongly witnesses*  $P^s(S)$  if, for all  $p \in W$ ,  $W_p$  witnesses  $P^s(S_p)$ , i.e.  $W$  witnesses  $P^s(S)$  and  $P^{s \upharpoonright n}(W)$  fails everywhere. By Remark 4.3, this relation is  $\Pi_{|s|+1}^1$ .

**Lemma 4.8.** *If  $P^s(S)$  then there is a  $W$  that strongly witnesses it.*

PROOF: Assume, for definiteness, that  $l$  is even. Let  $U$  witness  $P^s(S)$ . By Definition 4.1(2b),  $P^{s \upharpoonright n}(U)$  fails. By Lemma 4.6, let  $W$  be a quasistrategy for I in  $U$  such that  $P^{s \upharpoonright n}(W)$  fails everywhere. As  $[W] \subseteq [U]$ , Definition 4.1(2a) holds of  $W$  and so it strongly witnesses  $P^s(S)$  as required. The proof for  $l$  odd is the same except that we interchange I and II as well as clauses (2) and (3) of Definition 4.1.  $\square$

**Lemma 4.9.** *If  $|s| = n + 1$  then at least one of  $P^s(S)$  and  $P^{s \upharpoonright n}(S)$  holds.*

PROOF: We prove the Lemma by reverse induction on  $n < m$ . Suppose for definiteness that  $m - n$  is odd. If  $P^s(S)$  fails we define by induction on the length of positions (using strong  $\Sigma_{m+2}^1$ -DC<sub>0</sub>) a quasistrategy  $U$  for II in  $S$  along with  $D \subseteq S$  and  $R^d$  for  $d \in D$  locally witnessing  $P^{s \upharpoonright n}(S)$  if  $n > 0$  and witnessing  $P^\emptyset(S)$  if  $n = 0$ . Thus, by Lemma 4.4 when  $n > 0$ ,  $P^{s \upharpoonright n}(S)$  holds as required. We will decide which  $q$  are in  $U$  by recursion on the length of  $q$ , starting with  $\emptyset \in U$  and going up the tree  $S$ . Along the way we will enumerate some of these  $q$  into  $D$  and define a quasistrategy  $R^q$  for I. Inside each  $R^q$  we have to make sure to define  $U$  such that  $U \cap R^q$  witnesses  $P^{s \upharpoonright n}(R^q)$ . For  $x \in [U] \setminus \bigcup_{d \in D} [R^d]$  we need to make sure  $x \in \bar{A}$ , and we will do this as follows. On the one hand, we will define  $U$  so that  $x$  belongs to  $\bar{A} \cup A_{m-n-2,j}$  for every  $j \in \omega$ , and hence  $x \in \bar{A} \cup A_{m-n-2}$ . On the other hand, we will make sure that  $x \notin A_{m-n-1,s(n),j}$  for any  $j$ , and hence  $x \notin A_{m-n-1}$ . Then we use that  $\bar{A} \cup A_{m-n-2} \setminus A_{m-n-1} \subseteq \bar{A}$  to show that  $x \in \bar{A}$ . We will go along the path  $x$  satisfying the requirements  $x \in \bar{A} \cup A_{m-n-2,j}$  and  $x \notin A_{m-n-1,s(n),j}$  one  $j$  at the time. To keep track of which requirements need to be satisfied, along the construction we will define the notion of a position  $q \in U$  marking a  $j \in \omega$  (with the  $j$ 's marked in order along any path). Simultaneously we also define certain auxiliary quasistrategies  $W^q$  and  $\hat{R}^q$ .

Of course,  $\emptyset \in U$  and we say that it marks 0. If  $n = m - 1$ , by lemma 4.6, we let  $W^\emptyset$  be a quasistrategy for II in  $S$  such that  $P^s(W^\emptyset)$  fails everywhere. If  $n < m - 1$ , we know by (reverse induction) that  $P^{s \upharpoonright 0}(S)$  holds. In this case we apply Lemma 4.8 (and strong  $\Sigma_{m+2}^1$ -DC<sub>0</sub>) to choose a  $W^\emptyset$  strongly witnessing this fact (a  $\Pi_{|s|+2}^1$  and so at worst  $\Pi_{m+1}^1$  relation). (Note that in this case,  $P^s(W^\emptyset)$  also fails everywhere by Definition 4.7.)

Now assume by induction (on positions in  $U$ ) that  $q \in U$  marks  $j$  and that  $q$  belongs to a quasistrategy  $W^q$  for II in  $S_q$  such that  $P^s(W^q)$  fails everywhere and if  $n < m - 1$  then  $W^q$  strongly witnesses  $P^{s \uparrow j}(S_q)$ .

Recall that  $A_{m-n-1,s(n),j}$  is a closed set. If  $G(A_{m-n-1,s(n),j}, W^q)$  is not a win for II (a  $\Pi_2^1$  relation), put  $q$  in  $D$  and let  $\hat{R}^q$  be I's nonlosing quasistrategy in this game (a  $\Pi_2^1$  set in the parameters). Define  $R^q$  to be  $\hat{R}^q$  on  $W^q$  and to agree with  $S$  elsewhere in  $S_q$ . Now  $[\hat{R}^q] \subseteq A_{m-n-1,s(n),j}$  since it is a nonlosing quasistrategy for this closed set. Next note that, by definition,  $A_{m-n-1,s(n),j} \subseteq A_{m-n-1,s(n)}$ . Thus if  $P^{s \uparrow n}(\hat{R}^q)$  fails,  $\hat{R}^q$  would witness  $P^s(W^q)$  contrary to our assumption that  $P^s(W^q)$  fails everywhere. So we may take  $U^q$  to be a witness for  $P^{s \uparrow n}(\hat{R}^q)$  (a  $\Pi_{|s|}^1$  relation). We now continue to define  $U$  by letting it agree with  $U^q$  on  $\hat{R}^q$  and let no  $p \in \hat{R}^q$  into  $D$  nor mark any  $j'$ .

If beginning in this way with  $q$  we ever first reach a  $p \notin \hat{R}^q$  (which we take to be  $q$  if  $G(A_{m-n-1,s(n),j}, W^q)$  is a win for II), we continue in  $U$  by following a winning strategy  $\tau_p$  for II in  $G(A_{m-n-1,s(n),j}, W^q)$  (a  $\Pi_1^1$  relation) until we first reach a  $q' \supset p$  with  $[W_{q'}^q] \cap A_{m-n-1,s(n),j} = \emptyset$ . We must reach such a  $q'$  since  $\tau_p$  is a winning strategy for II in  $G(A_{m-n-1,s(n),j}, W^q)$  and  $A_{m-n-1,s(n),j}$  is a closed set. We do not mark any  $q^*$  with  $p \subseteq q^* \subset q'$  nor put any such into  $D$ . We now say that  $q'$  marks  $j + 1$ .  $P^s(W_{q'}^q)$  fails everywhere since  $P^s(W^q)$  does. If  $n = m - 1$ , we let  $W^{q'} = W_{q'}^q$ . If  $n < m - 1$ ,  $P^{s \uparrow j+1}(W_{q'}^q)$  holds by our reverse induction on  $n$  and we let  $W^{q'}$  strongly witness this fact (a  $\Pi_{|s|+2}^1$  and so at worst  $\Pi_{m+1}^1$  relation). Of course, it also strongly witnesses  $P^{s \uparrow (j+1)}(S_{q'})$  as required for our inductive definition of  $U$ .

We now prove that, if  $n > 0$ ,  $U$  (with  $D$  and  $R^d$ ) locally witnesses  $P^{s \uparrow n}(S)$ . First note that, by our definitions,  $U_d \cap R^d = U_d \cap \hat{R}^d = U^d$  for  $d \in D$ . Thus condition (i) of Definition 4.2 holds as  $U^d$  witnesses  $P^{s \uparrow n}(\hat{R}^d)$ . As we never put any  $q \supset d \in D$  into  $D$  until we have left  $\hat{R}^d$ , condition (iii) holds as well. We now turn to condition (ii) and consider any  $x \in [U]$ . Let  $\emptyset = q_0 \subset q_1 \dots \subset q_i \subset \dots$  be the strictly increasing sequence of  $q$ 's contained in  $x$  such that  $q_j$  marks  $j$ . By construction, each  $q_j \in D$ . If the sequence terminates at some  $d = q_k$ , then, by definition,  $x$  never leaves  $\hat{R}^d$  and so  $x \in [R^d]$  and (ii) holds trivially for this  $x$ . If not, then, for every  $j$ ,  $x \notin A_{m-n-1,s(n),j}$  as  $x \in [W_{q_{j+1}}^{q_j}]$  and  $[W_{q_{j+1}}^{q_j}] \cap A_{m-n-1,s(n),j} = \emptyset$ . Thus  $x \notin A_{m-n-1,s(n)} = \bigcup_j A_{m-n-1,s(n),j}$ . As  $A_{m-n-1,s(n)} \supseteq A_{m-n-1}$ ,  $x \notin A_{m-n-1}$ . Now if  $n + 1 = m$ ,  $x \notin A_0$  and so  $x \notin A$  and condition (ii) is satisfied. If  $n + 1 < m$  then, as  $W^{q_j}$  witnesses  $P^{s \uparrow j}(S_{q_{j+1}})$ ,  $x \in \bar{A} \cup A_{m-n-2,j}$  for each  $j$ . (Note that as  $m - n$  is odd so is  $m - |s \uparrow j|$  and we are in case (3) of Definition 4.1.) Thus  $x \in \bar{A} \cup A_{m-n-2}$  as  $A_{m-n-2} = \bigcap_j A_{m-n-2,j}$ . As, by our case assumptions,  $m - n - 1$  is even,  $A_{m-n-2} \setminus A_{m-n-1} \subseteq \bar{A}$  by the definition of  $A$  in terms of the sequence  $A_i$ . Thus  $x \in \bar{A}$  and condition (ii) is again satisfied.

Finally, if  $n = 0$  we argue that  $U$  is a winning quasistrategy for II in  $G(A, S)$ . Consider any  $x \in [U]$ . If there is a  $d \in D$  such that  $x \in [\hat{R}^d]$  then  $x \in [U^d]$  by construction. Now  $U^d$  is a witness for  $P^\emptyset(\hat{R}^d)$  (as  $n = 0$ ,  $s \uparrow n = \emptyset$ ), i.e.  $U^d$  is a winning strategy for II in  $G(A, \hat{R}^d)$ . Thus  $x \in \bar{A}$  as required. On the other hand, if  $x$  leaves every  $\hat{R}^d$  then, by the argument above,  $x \in \bar{A}$  as well.

For  $m - n$  even we interchange I and II as well as  $G(A_{m-n-1,s(n),j}, W^q)$  and  $G(\bar{A}_{m-n-1,s(n),j}, W^q)$  in the definition of  $U$ . In the verification that  $U$  is as desired note that the case that  $n + 1 = m$  cannot occur. Otherwise, we interchange I and II, cases (2) and (3) of Definition 4.1, even and odd as well as  $A$  and  $\bar{A}$ .  $\square$

PROOF: (of Theorem 1.1) If  $G(A, T)$  is not a win for II (I) if  $m$  is odd (even),  $P^\emptyset(T)$  fails. By Lemma 4.6, there is a quasistrategy  $W^\emptyset$  for I (II) such that  $P^\emptyset(W^\emptyset)$  fails everywhere. We define a quasistrategy  $U$  for I (II) in  $W^\emptyset$  by induction on  $|p|$  for  $p \in U$ . For  $|p| = j + 1$ , we simultaneously define (using strong  $\Sigma_3^1\text{-DC}_0$ ) a quasistrategy  $W^p$  for I (II) which strongly witnesses  $P^{(j)}(W_p^{p \upharpoonright j})$ . Of course,  $\emptyset \in U$  and we already have  $W^\emptyset$ . Suppose then that  $p \in U$ ,  $|p| = j + 1$ , and  $W^p$  has been defined. The immediate successors  $q$  of  $p$  in  $U$  are precisely its immediate successors in  $W^p$ . By our choice of  $W^p$ ,  $P^\emptyset(W^p)$  fails everywhere and so, in particular,  $P^\emptyset(W_q^p)$  fails for each immediate successor  $q$  of  $p$  in  $U$ . By Lemma 4.9, then  $P^{(j)}(W_q^p)$  holds and, by Lemma 4.8 (and strong  $\Sigma_3^1\text{-DC}_0$ ) we may choose a  $W^q$  that strongly witnesses it to continue our induction.

Consider now any play  $x \in [U]$ . By construction  $x \in [W^{x \upharpoonright j+1}]$  for every  $j$  and  $W^{x \upharpoonright j+1}$  witnesses  $P^{(j)}(W_{x \upharpoonright j+1}^{x \upharpoonright j})$ . By Definition 4.1 (2a) (4.1 (3a)) if  $m$  is odd (even),  $x \in A \cup A_{m-1, j}$  ( $\bar{A} \cup A_{m-1, j}$ ) for every  $j$ . As  $\bigcap_j A_{m-1, j} = A_{m-1} \subseteq A$  ( $\bar{A}$ ) (by the definition of  $A$  in terms of the sequence  $A_i$ ),  $x \in A$  ( $\bar{A}$ ). Thus  $U$  is a winning quasistrategy for I (II) in  $G(A, T)$  as desired.  $\square$

## 5. THE GAME: FAILURES OF DETERMINACY IN $L$

This section is dedicated to proving Lemma 1.8. Let  $T_n$  consist of  $\text{KP} + \Sigma_n\text{-admissibility} + \mathbf{V=L} + \forall \alpha \in \text{Ord}(L_\alpha \text{ is not } \Sigma_n\text{-admissible})$ . So  $L_{\alpha_n}$  is the only wellfounded model of  $T_n$ . Each player will have to play the characteristic function of a complete consistent extension of  $T_n$ , or otherwise they loose. The goal now is to be able to identify the player playing the true theory of  $L_{\alpha_n}$  using a Boolean combination of  $\Pi_3^0$  formulas.

To define  $G$  we first define  $2n + 4$  many  $\Pi_3^0$  conditions in order:

$$(R_{\text{I}0}), (R_{\text{II}0}), (R_{\text{II}1}), (R_{\text{II}1}), \dots, (R_{\text{II}}(n+1)), (R_{\text{II}}(n+1)).$$

Then, if the first condition satisfied by the play is of the form  $(R_{\text{I}k})$ , I wins, and if it is of the form  $(R_{\text{II}k})$ , then II wins. If no condition is ever satisfied, we will let one of the two players win; for now it does not matter which.

We now define these conditions one at a time, so we can make the motivations and the pertinent definitions along the way. We will later rearrange the conditions to reduce the level of the game in the  $\Pi_{3,n}^0$  hierarchy. Every time we define a new pair of conditions  $(R_{\text{I}k})$  and  $(R_{\text{II}k})$  we will show that they are  $\Pi_3^0$ , and that if any of these conditions is the first one to be satisfied, the choice of the winner is compatible with (1) and (2) of Lemma 1.8. At the end, we will show that, if one of the players is actually playing  $Th_{\alpha_n}$ , then some condition along the way will have to be satisfied, and hence (1) and (2) of Lemma 1.8 hold.

We begin with Conditions  $(R_{:,0})$ .

$$\begin{aligned} (R_{\text{I}0}): & \quad \text{II does not play a complete consistent extension of } T_n \\ (R_{\text{II}0}): & \quad \text{I does not play a complete consistent extension of } T_n \end{aligned}$$

*Claim 5.1.* Conditions  $(R_{\text{I}1})$  and  $(R_{\text{II}1})$  are  $\Sigma_1^0$ .

PROOF: Saying that the set of sentences played form a complete and consistent theory is  $\Pi_1^0$ . Saying that the axioms of  $T_n$  are included in these sets is also  $\Pi_1^0$ .  $\square$

If either of these conditions are satisfied, the game ends and a player playing  $Th_{\alpha_n}$  cannot loose at this stage. In this case it does not matter what the rest of the conditions say. So,

from now on, when we describe the rest of the conditions, we will assume neither of these two conditions apply.

The term model of a complete theory  $T$  consistent with  $T_n$  is built as follows: Consider the set of formulas with one free variable  $\varphi(x)$  such that  $T \vdash \exists!x\varphi(x)$ . Think of this formula  $\varphi$  as a term  $t_\varphi$  representing the unique element satisfying  $\varphi$ . So, given two such formulas  $\varphi$  and  $\psi$ , we say that  $t_\varphi = t_\psi$  if  $T$  proves that  $\varphi$  and  $\psi$  are equivalent. Also, we let  $t_\varphi \in t_\psi$  if  $T$  proves that the witness of  $\varphi$  belongs to the witness of  $\psi$ . This defines the term model of  $T$ . Since  $T_n$  proves that every nonempty class has a  $<_L$ -least element, every sentence  $\exists x\varphi(x)$  in  $T$  has a term witnessing it. So, this term model we just built is a model of  $T$  and, furthermore, from  $T$  we can compute its full elementary diagram. Note that the term model of  $Th_{\alpha_n}$  is isomorphic to  $L_{\alpha_n}$  because every element of  $L_{\alpha_n}$  is definable without parameters by Lemma 3.6.

Let  $\mathcal{M}_I$  be the term model of the theory built by I and  $\mathcal{M}_{II}$  the one built by II. So, from the play, we can compute the full elementary diagrams of  $\mathcal{M}_I$  and  $\mathcal{M}_{II}$ . These models might be wellfounded or not. But recall that the only wellfounded model of  $T_n$  is  $L_{\alpha_n}$ . All the other models of  $T_n$  are ill-founded. We now continue with the next pair of conditions.

- ( $R_I1$ ):  $\mathcal{M}_{II}$  is not an  $\omega$ -model.  
 ( $R_{II}1$ ):  $\mathcal{M}_I$  is not an  $\omega$ -model.

*Claim 5.2.* Conditions ( $R_I1$ ) and ( $R_{II}1$ ) are  $\Sigma_2^0$ .

PROOF: To say that a term model  $\mathcal{M}$  is an  $\omega$ -model, we have to say that for every term  $t$ , if  $\mathcal{M} \models t \in \omega$ , then there exists a number  $n$  such that  $\mathcal{M} \models t = 1 + 1 + \dots + 1$ , where 1 is added  $n$  times. This is a  $\Pi_2^0$  formula about  $\mathcal{M}$ .  $\square$

We remark that the proof of the claim above works when both players have actually played complete consistent extensions of  $T_n$ . Formally, we think of ( $R_{II}1$ ) and ( $R_I1$ ) as the  $\Sigma_2^0$  formulas given in the proof of the claim. The interpretation of these  $\Sigma_2^0$  formulas will match what we first called ( $R_I1$ ) and ( $R_{II}1$ ) above the claim, when conditions ( $R_I0$ ) and ( $R_{II}0$ ) are not satisfied, but that is the only case where we care. Similar remarks will apply to the conditions ( $R_{:,k}$ ) below. That is, we supply an intuitive description of sets in our list and then a formal  $\Pi_3^0$  definition that makes the intuition true in the cases of interest. As we continue to define more conditions on the winning set, keep in mind that our goal as we go along is to guarantee for each condition that if I plays the theory of  $L_{\alpha_n}$  then he does not loose. If he does not play  $Th_{\alpha_n}$  and II does, then II does not loose. At the end we will prove that someone wins the game.

$L_{\alpha_n}$  is an  $\omega$ -model, so a player playing  $Th_{\alpha_n}$ , cannot loose at this step either. From now on assume (for the intuitive descriptions of our clauses) that both  $\mathcal{M}_I$  and  $\mathcal{M}_{II}$  are  $\omega$ -models of  $T_n$  and that at least one of them is  $L_{\alpha_n}$ . Let us move on to the third pair of conditions.

- ( $R_I2$ ):  $\mathbb{R}_{\mathcal{M}_I} \subseteq \mathbb{R}_{\mathcal{M}_{II}}$ .  
 ( $R_{II}2$ ):  $\mathbb{R}_{\mathcal{M}_{II}} \subseteq \mathbb{R}_{\mathcal{M}_I}$ .

If both players are playing  $Th_{\alpha_n}$ , then ( $R_I2$ ) holds, and hence I wins as needed for (1) of Lemma 1.8. So, from now on, let us assume that one player is playing  $Th_{\alpha_n}$  and the other one is not. Next, note that  $L_{\alpha_n}$  has no proper sub-models satisfying  $T_n$ , and hence the player playing  $Th_{\alpha_n}$  cannot loose by either of these conditions.

*Claim 5.3.* Conditions  $(R_I2)$  and  $(R_{II}2)$  are  $\Pi_3^0$ .

**PROOF:** For  $(R_I2)$ , it is enough to say that every ordinal in  $\mathcal{M}_I$ , is also an ordinal in  $\mathcal{M}_{II}$ : If a linear ordering  $\alpha$  belongs to both  $On^{\mathcal{M}_I}$  and  $On^{\mathcal{M}_{II}}$ , it has to actually be wellfounded as we are assuming one of  $\mathcal{M}_I$  or  $\mathcal{M}_{II}$  is wellfounded. So, by transfinite induction, this implies that  $L_\alpha$  also belongs to both  $\mathcal{M}_I$  and  $\mathcal{M}_{II}$ . Now, note that since every ordinal  $\alpha$  in  $\mathcal{M}_I$  is countable in  $\mathcal{M}_I$  (Lemma 3.5) there is always a real representing a linear ordering that is isomorphic to  $\alpha$ .

To say that two ordinals  $\alpha$  and  $\beta$  (even in different models) are the same, we say that there is a real  $x$  coding  $\alpha$  in  $\mathcal{M}_I$  and a real  $y$  coding an ordinal  $\beta$  in  $\mathcal{M}_{II}$ , such that  $x = y$ . Even if  $x$  and  $y$  belong to different models, we can say they represent the same real by saying that for every natural number  $n$ ,  $\mathcal{M}_I \models n \in x \iff \mathcal{M}_{II} \models n \in y$ . Therefore, the  $\Pi_3^0$  sentence we need is

$$(5.1) \quad \forall \alpha \in On^{\mathcal{M}_I} \exists x \in \mathbb{R}^{\mathcal{M}_I}, y \in \mathbb{R}^{\mathcal{M}_{II}}, \beta \in On^{\mathcal{M}_{II}} [(\mathcal{M}_I \models x \text{ codes } \alpha) \\ \wedge (\mathcal{M}_{II} \models y \text{ codes } \beta) \wedge \forall n \in \omega (\mathcal{M}_I \models n \in x \iff \mathcal{M}_{II} \models n \in y)].$$


□

We remark again, that the claim should actually say: If  $\mathcal{M}_I$  and  $\mathcal{M}_{II}$  are  $\omega$ -models of  $T_n$  and at least one of them is wellfounded, then  $(R_I2)$  and  $(R_{II}2)$  are equivalent to  $\Pi_3^0$  properties. In any other case, we think of  $(R_I2)$  and  $(R_{II}2)$  as given formally by these  $\Pi_3^0$  properties, which, in cases not of interest, might not have the intended meaning.

To this point, our proof follows the ideas in Martin [1974a] and [n.d., Ch. 1] where he shows that  $\Sigma_4^0$ -determinacy is not true in the least  $\beta$ -model of second order arithmetic  $\mathbb{R} \cap L_{\beta_0}$ . These, in turn, are based on those of H. Friedman's [1971] proof that  $\Sigma_5^0$  determinacy fails in  $\mathbb{R} \cap L_{\beta_0}$ . The ideas in the rest of the proof are new.

From now on, we assume that no condition listed so far holds and so, in particular, not both of  $\mathcal{M}_I$  and  $\mathcal{M}_{II}$  are well founded. Again, we are interested in the plays where one is well founded and so actually  $L_{\alpha_n}$ .

**Definition 5.4.** We use  $\mathcal{M}$  to denote the wellfounded model of the pair  $\mathcal{M}_I$  and  $\mathcal{M}_{II}$  and  $\mathcal{N}$  for the ill-founded one. The player playing  $\mathcal{M} \cong L_{\alpha_n}$  should win the game but we do not know which one it is. There is a least ordinal  $\alpha \in On^{\mathcal{M}}$  that is not in  $\mathcal{N}$ . Let  $\mathcal{A}$  be the well founded part of  $\mathcal{N}$ . So, we have that  $L_\alpha \cong \mathcal{A}$ . Let us call  $\mathcal{A}_I$  and  $\mathcal{A}_{II}$  the initial segments of  $\mathcal{M}_I$  and  $\mathcal{M}_{II}$  that correspond to  $L_\alpha$  and  $\mathcal{A}$  in  $\mathcal{M}$  and  $\mathcal{N}$ .



We abuse notation slightly by using  $\mathcal{A}$  to also denote the isomorphism between  $\mathcal{A}_I$  and  $\mathcal{A}_{II}$ . That is,  $\mathcal{A}$  is the set of pairs of terms  $(z, w) \in \mathcal{M}_I \times \mathcal{M}_{II}$  representing elements  $z \in \mathcal{A}_I$  and  $w \in \mathcal{A}_{II}$  which are the same set under the isomorphism  $\mathcal{A}_I \cong \mathcal{A}_{II}$ .

*Claim 5.5.*  $\mathcal{A}$  is a  $\Sigma_2^0$  set. Also,  $\mathcal{A}_I$  and  $\mathcal{A}_{II}$  are  $\Sigma_2^0$  sets.

PROOF: For  $\mathcal{A}$  we have the following  $\Sigma_2^0$  definition.

$$(z, w) \in \mathcal{A} \iff \exists x \in \mathbb{R}^{\mathcal{M}_I}, y \in \mathbb{R}^{\mathcal{M}_{II}} \left( \mathcal{M}_I \models x \text{ codes } z \wedge \mathcal{M}_{II} \models y \text{ codes } w \wedge \forall n \in \omega (\mathcal{M}_I \models n \in x \iff \mathcal{M}_{II} \models n \in y) \right).$$

For  $\mathcal{A}_I$  we have that  $z \in \mathcal{A}_I \iff \exists w ((z, w) \in \mathcal{A})$ , and analogously for  $\mathcal{A}_{II}$ .  $\square$

(At this point, Martin's proof ([1974a] and [n.d., Ch. 1]) that  $\Sigma_4^0$ -DET fails in  $L_{\beta_0}$  has a  $\Sigma_4^0$  winning condition that says that there exists a  $\alpha \in \mathcal{M}_I$  such that  $L_\alpha^{\mathcal{M}_I} = \mathcal{A}_I$ .)

The next  $n - 1$  pairs of conditions all have basically the same form. The conditions  $(R_{\cdot};3)$  are a bit simpler than the rest and we describe them first. We will then generalize them and define the conditions  $(R_{\cdot};(2+k))$  for  $k > 2$  all at once.

We start by defining the sets of ordinals in one model that have witnesses for  $\Sigma_1$  formulas with parameters in  $\mathcal{A}$  that have no witnesses in the other model. The expectation is that the witnesses are in the nonwellfounded model  $\mathcal{N}$ . We will show later that either there are many such witnesses and the set of witnesses shows that the  $On^{\mathcal{N}} \setminus \mathcal{A}$  has no least element; or there are few such witnesses and there is an ordinal  $\beta \in \mathcal{N}$  such that  $\mathcal{A} \preceq_1 L_\beta^{\mathcal{N}}$ . In the former case, we have identified the nonwellfounded model and we can end the game. In the latter case, we continue with the next pair of conditions, now assuming the existence of such a  $\beta$ .

Let

$$C_{\mathcal{M}_{II},1} = \{ \beta \in On^{\mathcal{M}_{II}} : \exists (x_1, x_2) \in \mathcal{A}, \varphi \in \Delta_0, ((\exists z \in L_\beta^{\mathcal{M}_{II}} \mathcal{M}_{II} \models \varphi(z, x_2)) \wedge (\mathcal{M}_I \models \neg \exists y \varphi(y, x_1))) \}.$$

Note that this condition says more than that  $\mathcal{A}$  is not a  $\Sigma_1$  substructure of  $L_\beta^{\mathcal{M}_{II}}$  as it asserts that there are no witnesses for the formula  $\varphi$  even in  $\mathcal{M}_I$ . Thus its negation does not imply that  $\mathcal{A}$  is a  $\Sigma_1$  substructure of  $L_\beta^{\mathcal{M}_{II}}$ .  $C_{\mathcal{M}_I,1}$  is defined analogously. These sets are  $\Sigma_2^0$  because, as we noted before,  $\mathcal{A}$  is  $\Sigma_2^0$  and the second line of the definition is computable from the elementary diagrams of  $\mathcal{M}_I$  and  $\mathcal{M}_{II}$ . We are now ready to define the next pair of conditions.

$$(R_I3): \quad C_{\mathcal{M}_{II},1} \text{ has no least element.}$$

$$(R_{II}3): \quad C_{\mathcal{M}_I,1} \text{ has no least element.}$$

Clearly a player playing a wellfounded model cannot lose because of these conditions. Also, these conditions are  $\Pi_3^0$  as they say, for example, that  $\forall \beta (\beta \notin C_{\mathcal{M}_{II},k} \vee \exists \gamma < \beta (\gamma \in C_{\mathcal{M}_{II},k}))$  and  $C_{\mathcal{M}_{II},k}$  is  $\Sigma_2^0$ .

Suppose now that neither of these conditions is satisfied.

*Claim 5.6.* There is a  $\beta \in On^{\mathcal{N}} \setminus \mathcal{A}$  such that  $\mathcal{A} \preceq_1 L_\beta^{\mathcal{N}}$ .

PROOF: Suppose not, that is, for every  $\gamma \in On^{\mathcal{N}} \setminus \mathcal{A}$ , there is a  $\Sigma_1$  formula with parameters in  $\mathcal{A}$ , that is true in  $L_\gamma$  but not in  $\mathcal{A}$ . Since conditions  $(R_I3)$  and  $(R_{II}3)$  are not satisfied,  $C_{\mathcal{N},1}$  has a least element  $\delta$ . Notice that  $\delta$  cannot be in  $\mathcal{A}$  because if the witness  $z$  for a  $\Delta_0$

formula  $\varphi(z, x_2)$  is in  $L_{\mathcal{N}_\delta} \subseteq \mathcal{A}$ , then  $\exists y \varphi(y, x_1)$  also holds in  $\mathcal{M}$ . Let  $\delta > \gamma_0 > \gamma_1 > \gamma_2 > \dots$  be a descending sequence in  $On^{\mathcal{N}} \setminus \mathcal{A}$ , converging down to the cut  $(On^{\mathcal{A}}, On^{\mathcal{N}} \setminus \mathcal{A})$ . For each  $i$ , there is a  $\Delta_0$  formula  $\varphi_i$  with parameters from  $\mathcal{A}$  and a  $<_L$ -least witness  $z_i \in L_{\gamma_i}$  such that  $\mathcal{N} \models \varphi_i(z_i)$  but  $\mathcal{A} \models \neg \exists y \varphi_i(y)$ . By thinning out the sequence if necessary, we may assume that  $z_i \in L_{\gamma_i}$  but  $z_i \notin L_{\gamma_{i+1}}$ . So  $\{z_i : i \in \omega\}$  is an  $<_L^{\mathcal{N}}$ -descending sequence. Since  $\gamma_i \notin C_{\mathcal{N},1}$ ,  $\mathcal{M} \models \exists y \varphi_i(y)$ . Let  $y_i$  be the  $<_L^{\mathcal{M}}$ -least such witness. Since  $\mathcal{M}$  is wellfounded, the sequence  $\{y_i : i \in \omega\}$  cannot be a  $<_L^{\mathcal{M}}$ -descending sequence. So, there exists  $i < j$  such that  $z_j <_L^{\mathcal{N}} z_i$  but  $y_i <_L^{\mathcal{M}} y_j$ . Therefore,  $L_{\gamma_j}$  is a witness in  $L_{\gamma_{j+1}}$  for  $x$  showing the truth in  $\mathcal{N}$  of the  $\Delta_0$  formula

$$\psi(x) \equiv \exists z \in x \varphi_j(z) \wedge \forall z \in x \neg \varphi_i(z)$$

but there is no witness for  $\psi(x)$  in  $\mathcal{M}$ . This shows that  $\gamma_j + 1$  is in  $C_{\mathcal{N},1}$ , contradicting our choice of  $\delta$  as the least element of  $C_{\mathcal{N},1}$ .  $\square$

We now turn to defining the rest of the conditions. Again, the idea is to define conditions such that they hold only if they have identified the nonwellfounded model, and if they do not hold, they give us information about the models that we can assume when we define the following conditions. We will show that if for some  $k$ , none of the conditions  $(R_{::1}), \dots, (R_{::(2+k)})$  is satisfied, then  $\alpha$  is  $(k+1)$ -admissible. Since  $L_{\alpha_n}$  has no  $n$ -admissible ordinals, we will be able to argue that some condition  $(R_{::(2+k)})$  will have to be satisfied for some  $k < n$ . This will be proved, of course, by induction on  $k$ , but the induction hypothesis needs to be stronger. We show that if no condition  $(R_{::1}), \dots, (R_{::(2+k)})$  holds then there exists  $\beta_1$  and  $\beta_2$  such that

$$\begin{aligned} (\star_k)(\beta_1, \beta_2) : \quad & \beta_1 \in On^{\mathcal{M}_I} \setminus \mathcal{A}_I \wedge \mathcal{M}_I \models \beta_1 \text{ is } (k-1)\text{-admissible} \wedge \\ & \beta_2 \in On^{\mathcal{M}_{II}} \setminus \mathcal{A}_{II} \wedge \mathcal{M}_{II} \models \beta_2 \text{ is } (k-1)\text{-admissible} \wedge \\ & L_{\beta_1}^{\mathcal{M}_I} \equiv_{k, \mathcal{A}} L_{\beta_2}^{\mathcal{M}_{II}}. \end{aligned}$$

Here  $\equiv_{k, \mathcal{A}}$  means  $\Sigma_k$ -elementary equivalence with parameters from  $\mathcal{A}$ . In other words,  $L_{\beta_1}^{\mathcal{M}_I} \equiv_{k, \mathcal{A}} L_{\beta_2}^{\mathcal{M}_{II}}$  means that for every  $\Sigma_k$  formula  $\varphi(z)$ , and every  $x_1 \in \mathcal{A}_I$  and  $x_2 \in \mathcal{A}_{II}$ , if  $x_1$  and  $x_2$  represent the same set (i.e.  $(x_1, x_2) \in \mathcal{A}$ ), then  $\mathcal{M}_I \models \varphi(x_1) \iff \mathcal{M}_{II} \models \varphi(x_2)$ .

By Claim 5.6, we already know there are ordinals satisfying  $(\star_1)$ : Just let  $\beta_1 = \alpha \in \mathcal{M}$  and let  $\beta_2 \in \mathcal{N}$  be such that  $\mathcal{A} \preceq_1 L_{\beta_2}^{\mathcal{N}}$ .

Property  $(\star_k)$  is useful because, on the one hand, it is simple to define (it is  $\Pi_2^0$  as we prove below) and, on the other hand, it has strong consequences as we will show in Lemma 5.10. For instance, we will show that the existence of  $\beta_1$  and  $\beta_2$  satisfying it implies that  $\alpha$  is  $(k+1)$ -admissible.

**Lemma 5.7.**  $(\star_k)$  is a  $\Pi_2^0$  property.

PROOF: We already showed that  $\beta_1 \notin \mathcal{A}_I \wedge \beta_2 \notin \mathcal{A}_{II}$  is a  $\Pi_2^0$  relation. Whether  $\beta_i$  is  $(k-1)$ -admissible is just one sentence in the theory of the structure. To check that  $L_{\beta_1}^{\mathcal{M}_I} \equiv_{k, \mathcal{A}} L_{\beta_2}^{\mathcal{M}_{II}}$ , we need to say that, for every  $\Sigma_k$  formula  $\varphi(x)$ , and for every pair of parameters  $(x_1, x_2) \in \mathcal{A}$ ,  $\mathcal{M}_I \models \varphi(x_1) \iff \mathcal{M}_{II} \models \varphi(x_2)$ . Recall that  $\mathcal{A}$  is  $\Sigma_2^0$  (Claim 5.5). Counting quantifiers we see that  $(\star_k)$  is  $\Pi_2^0$ .  $\square$

**Definition 5.8.**  $S_k$  is the set of formulas that are Boolean combinations of formulas of the form  $(\forall x \in z) \psi(z, \bar{y})$ , where  $\psi$  is  $\Sigma_k$ .

**Lemma 5.9.** *If  $L_{\beta_1}^{\mathcal{M}_I} \equiv_{k,\mathcal{A}} L_{\beta_2}^{\mathcal{M}_{II}}$ , then  $L_{\beta_1}^{\mathcal{M}_I}$  and  $L_{\beta_2}^{\mathcal{M}_{II}}$  satisfy the same  $S_k$ -sentences with parameters from  $\mathcal{A}$  substituted for the free variables  $z$  and  $\bar{y}$ .*

PROOF: Let  $\varphi$  be a sentence of the form  $\forall x \in z \psi(x, z, \bar{y})$  with parameters from  $\mathcal{A}$  where  $\psi$  is  $\Sigma_k$ . Since  $\varphi$  has no free variables,  $z$  and  $\bar{y}$  are parameters from  $\mathcal{A}$ , and so every  $x \in z$  is also in  $\mathcal{A}$ . Since  $L_{\beta_1}^{\mathcal{M}_I} \equiv_{k,\mathcal{A}} L_{\beta_2}^{\mathcal{M}_{II}}$ , we have that for every  $x \in z$ ,  $L_{\beta_1}^{\mathcal{M}_I} \models \psi(x, z, \bar{y}) \Leftrightarrow L_{\beta_2}^{\mathcal{M}_{II}} \models \psi(x, z, \bar{y})$ . So, of course,  $L_{\beta_1}^{\mathcal{M}_I} \models \varphi \Leftrightarrow L_{\beta_2}^{\mathcal{M}_{II}} \models \varphi$ . Finally, if  $\phi$  is a Boolean combination of sentences of this form, then we also have  $L_{\beta_1}^{\mathcal{M}_I} \models \phi \Leftrightarrow L_{\beta_2}^{\mathcal{M}_{II}} \models \phi$ .  $\square$

Now we define sets that generalizes  $C_{\mathcal{M}_I,1}$  and  $C_{\mathcal{M}_{II},1}$ . This time, the sets consist of the ordinals in one model ( $\mathcal{N}$ ) that have witnesses for  $S_k$  formulas with parameters in  $\mathcal{A}$  that have no witnesses in the other model ( $\mathcal{M}$ ). We will show later that: either there are lots of such witnesses and the set  $C_{\mathcal{N},k}^{\beta_1,\beta_2}$  shows that  $On^{\mathcal{N}} \setminus \mathcal{A}$  has no least element, in which case we have identified the nonwellfounded model, or there are few such witnesses and we can find ordinals satisfying  $(\star_k)$ .

For  $\beta_1 \in \mathcal{M}_I$  and  $\beta_2 \in \mathcal{M}_{II}$ , let

$$C_{\mathcal{M}_{II},k}^{\beta_1,\beta_2} = \{\beta \in \beta_2 : \exists(x_1, x_2) \in \mathcal{A}, \exists\varphi \in S_{k-1} \\ (\exists z \in L_{\beta}^{\mathcal{M}_{II}} L_{\beta_2}^{\mathcal{M}_{II}} \models \varphi(z, x_2)) \wedge (L_{\beta_1}^{\mathcal{M}_I} \models \neg\exists y \varphi(y, x_1))\}.$$

and define  $C_{\mathcal{M}_I,k}^{\beta_1,\beta_2}$  analogously. As in the case  $k = 1$ , these sets are  $\Sigma_2^0$ .

$(R_I(2+k))$ : For every  $\beta_1, \beta_2$ , if  $(\star_{k-1})(\beta_1, \beta_2)$ , then  $C_{\mathcal{M}_{II},k}^{\beta_1,\beta_2}$  has no least element.

$(R_{II}(2+k))$ : For every  $\beta_1, \beta_2$ , if  $(\star_{k-1})(\beta_1, \beta_2)$ , then  $C_{\mathcal{M}_I,k}^{\beta_1,\beta_2}$  has no least element.

The conditions  $(R_{II}3)$  and  $(R_I3)$  are  $\Pi_3^0$ : We already showed that  $(\star_{k-1})$  is a  $\Pi_2^0$  property, and, as for  $k = 1$ ,  $C_{\mathcal{M}_{II},k}^{\beta_1,\beta_2}$  having no least element is  $\Pi_3^0$  because  $C_{\mathcal{M}_{II},k}^{\beta_1,\beta_2}$  is  $\Sigma_2^0$ . Also observe that if there are  $\beta_1, \beta_2$  satisfying  $(\star_{k-1})$ , then the player playing  $Th_{\alpha_n}$  cannot loose by a condition  $(R_{\cdot}(2+k))$ .

**Lemma 5.10.** *Suppose that  $\beta_1, \beta_2$  satisfy  $(\star_k)$ . For the sake of definiteness, suppose that  $\beta_1 \in \mathcal{M}$  and  $\beta_2 \in \mathcal{N}$ . Then*

- (1)  $L_{\alpha} \preceq_k L_{\beta_1}$  and  $\mathcal{A} \preceq_k L_{\beta_2}$ .
- (2)  $\alpha$  is  $(k+1)$ -admissible.
- (3) There exists a descending sequence of  $\mathcal{N}$  ordinals  $\gamma$  converging down to  $On^{\mathcal{A}}$  such that  $L_{\gamma} \preceq_k L_{\beta_2}$ .

PROOF: First, we note that  $\{\alpha\}$  is not  $\Sigma_k$  definable in  $L_{\beta_1}$  with parameters from  $L_{\alpha}$ . Suppose otherwise. Since  $\alpha \in \mathcal{M} \models T$ ,  $\alpha$  is not  $\Sigma_n$ -admissible and every  $\beta \in \mathcal{M}$  is countable in  $\mathcal{M}$ . Thus there is a  $\Sigma_n/L_{\alpha}$  map from  $\omega$  onto  $\alpha$  (possibly with parameters). This defines in  $L_{\alpha}$ , a  $\Sigma_n$  ordering of  $\omega$  of order type  $\alpha$ . This ordering cannot belong to  $\mathcal{N}$  as it would then define its wellfounded part. In  $L_{\beta_1}$  we can define this ordering with a  $\Sigma_k$  formula using the  $\Sigma_k$  definition of  $\alpha$  and quantification over  $L_{\alpha}$ . Since  $L_{\beta_1}^{\mathcal{M}_I} \equiv_{k,\mathcal{A}} L_{\beta_2}^{\mathcal{M}_{II}}$ , this ordering is definable in  $L_{\beta_2}$  by the same formula, and hence belongs  $\mathcal{N}$ . Contradiction.

Now for part (1), since  $\beta_1$  is  $(k-1)$ -admissible,  $L_{\beta_1}$  has a  $\Sigma_k$ -Skolem function without parameters (Lemma 3.1). Let  $H$  be the  $\Sigma_k$ -Skolem hull of  $L_{\alpha}$  in  $L_{\beta_1}$ . We show that  $H = L_{\alpha}$ . Note that this implies that  $L_{\alpha} \preceq_k L_{\beta_1}$  (as desired) and also  $\mathcal{A} \preceq_k L_{\beta_2}$  since  $L_{\beta_1}^{\mathcal{M}_I} \equiv_{k,\mathcal{A}} L_{\beta_2}^{\mathcal{M}_{II}}$ .

Suppose that  $H \supseteq L_\alpha$ . Let  $L_\gamma$  be the Mostowski collapse of  $H$ , so  $\alpha < \gamma \leq \beta_1$ . Let  $\alpha'$  be the ordinal in  $H$  that is sent to  $\alpha \in L_\gamma$  by the collapse. As  $\alpha'$  is in the image of the  $\Sigma_k$  Skolem function,  $\{\alpha'\}$  is  $\Sigma_k$  definable in  $H$  with parameters from  $L_\alpha$ . The same formula then gives a definition of  $\{\alpha\}$  in  $L_\gamma$  as the collapse is the identity on members of  $L_\alpha$ . Thus  $\{\alpha\}$  is  $\Sigma_k$  definable in  $L_\gamma$  with parameters from  $L_\alpha$  but clearly

$$L_\gamma^{\mathcal{M}} \equiv_{k, \mathcal{A}} H \equiv_{k, \mathcal{A}} L_{\beta_1}^{\mathcal{M}}.$$

So  $\{\alpha\}$  would be  $\Sigma_k$  definable in  $L_{\beta_1}^{\mathcal{M}}$  contradicting our first observation.

(2) Suppose  $\alpha$  is not  $(k+1)$ -admissible and so (since every ordinal is countable in  $\mathcal{M}$ ) there is a  $\Pi_k/L_\alpha$  function  $f$  from  $\omega$  onto  $\alpha$ . Since  $\mathcal{A} \preceq_k L_{\beta_2}$ , the same formula defining  $f$  over  $L_\alpha$  defines a function from  $\omega$  onto  $\alpha$  in  $L_{\beta_2}$ . But then, in  $\mathcal{N}$  we could define its wellfounded part,  $L_\alpha$ , for a contradiction.

Now since  $\alpha$  is  $(k+1)$ -admissible, there are unboundedly many  $\gamma < \alpha$  such that  $L_\gamma \preceq_k L_\alpha$  (Lemma 3.3). For all these  $\gamma$  we also have  $L_\gamma^{\mathcal{N}} \preceq_k L_{\beta_2}^{\mathcal{N}}$ . The set of  $\gamma < \beta_2$  such that  $L_\gamma^{\mathcal{N}} \preceq_k L_{\beta_2}^{\mathcal{N}}$  is definable in  $\mathcal{N}$ . If for some  $\delta \in On^{\mathcal{N}} \setminus \mathcal{A}$ , the set of  $\gamma < \delta$  such that  $L_\gamma^{\mathcal{N}} \preceq_k L_{\beta_2}^{\mathcal{N}}$  had supremum  $\alpha$ , then  $\alpha$  would be definable in  $\mathcal{N}$  and we know it is not. So for every  $\delta \in On^{\mathcal{N}} \setminus \mathcal{A}$ , there exists  $\gamma < \delta$ ,  $\gamma \in On^{\mathcal{N}} \setminus \mathcal{A}$  such that  $L_\gamma^{\mathcal{N}} \preceq_k L_{\beta_2}^{\mathcal{N}}$ .  $\square$

**Lemma 5.11.** *If there is a play of our game such that for all  $i \leq 2+k$ , the resulting real does not satisfy any condition  $(R_{\text{I}i})$  or  $(R_{\text{II}i})$ , then there are  $\beta_1$  and  $\beta_2$  satisfying  $(\star_k)$ .*

PROOF: We already showed that there are  $\beta_1$  and  $\beta_2$  satisfying  $(\star_1)$ . The proof now proceeds by induction, so, without loss of generality, we assume that there exists  $\beta_1, \beta_2$  satisfying  $(\star_{k-1})$ . We fix such  $\beta_1, \beta_2$ .

We claim that no  $\delta \in \mathcal{A}$  is in  $C_{\mathcal{N}, k}^{\beta_1, \beta_2}$ . Consider  $\delta \in \mathcal{A}$  and any  $S_{k-1}$  formula  $\forall x \in z\varphi(z, \bar{y})$  and  $z_2, \bar{y}_2 \in L_\delta^{\mathcal{N}} \subseteq \mathcal{A}$  such that  $L_{\beta_2}^{\mathcal{N}} \models \forall x \in z_2\varphi(z_2, \bar{y}_2)$ . Since  $L_{\beta_1}$  and  $L_{\beta_2}$  satisfy the same  $\Sigma_{k-1}$  formulas with parameters from  $\mathcal{A}$  (by  $(\star_{k-1})$ ),  $L_{\beta_1}^{\mathcal{M}} \models \forall x \in z_1\varphi(z_1, \bar{y}_1)$ , where  $z_1$  and  $\bar{y}_1$  are the images of  $z_2, \bar{y}_2$  in  $\mathcal{M}$ . Thus  $\delta \notin C_{\mathcal{N}, k}^{\beta_1, \beta_2}$  by its definition.

Since the conditions  $(R_{\text{I}}(2+k))$  and  $(R_{\text{II}}(2+k))$  are not satisfied,  $C_{\mathcal{N}, k}^{\beta_1, \beta_2}$  has a least element  $\delta$ , necessarily not in  $\mathcal{A}$ . By clause (3) of the previous lemma, there is a descending sequence  $\delta > \gamma_0 > \gamma_1 > \gamma_2 > \dots$  in  $On^{\mathcal{N}}$ , converging down to  $\alpha = On^{\mathcal{A}}$ , such that for each  $i$   $L_{\gamma_i}^{\mathcal{N}} \preceq_{k-1} L_{\beta_2}^{\mathcal{N}}$ . We now argue much as for Claim 5.6. We claim that, for some  $i$ ,  $L_\alpha \preceq_k L_{\gamma_i}^{\mathcal{N}}$ . If not, there is, for each  $i$ , a  $\Pi_{k-1}$  formula  $\varphi_i(z)$  with parameters from  $\mathcal{A}$  and a  $<_L^{\mathcal{N}}$ -least  $z_i \in L_{\gamma_i}^{\mathcal{N}}$  such that  $L_{\gamma_i}^{\mathcal{N}} \models \varphi_i(z_i)$  and  $z_i \notin \mathcal{A}$ . As  $L_{\gamma_i}^{\mathcal{N}} \preceq_{k-1} L_{\beta_2}^{\mathcal{N}}$ ,  $L_{\beta_2}^{\mathcal{N}} \models \varphi_i(z_i)$ . By thinning out the sequence if necessary, we may assume that  $z_i \in L_{\gamma_i}^{\mathcal{N}}$  but  $z_i \notin L_{\gamma_{i+1}}^{\mathcal{N}}$ . So  $\{z_i : i \in \omega\}$  is an  $<_L^{\mathcal{N}}$ -descending sequence. Since  $\gamma_i \notin C_{\mathcal{N}, k}^{\beta_1, \beta_2}$ ,  $L_{\beta_1}^{\mathcal{M}} \models \exists y\varphi_i(y)$ . Let  $y_i$  be the  $<_L^{\mathcal{M}}$ -least such witness. Since  $\mathcal{M}$  is wellfounded, the sequence  $\{y_i : i \in \omega\}$  cannot be a  $<_L^{\mathcal{M}}$ -descending sequence. So, there exist  $i < j$  such that  $z_j <_L^{\mathcal{N}} z_i$  but  $y_i <_L^{\mathcal{M}} y_j$ . Therefore,  $L_{\gamma_j}^{\mathcal{N}} \in L_{\gamma_{j+1}}^{\mathcal{N}}$  is a witness for  $x$  showing the truth in  $L_{\beta_2}^{\mathcal{N}}$  of the  $S_{k-1}$  formula

$$\psi(x) \equiv \exists z \in x\varphi_j(z, x) \wedge \forall z \in x\neg\varphi_i(z, x)$$

while there is no such witness for the corresponding formula in  $L_{\beta_1}^{\mathcal{M}}$ . This contradicts our choice of  $\delta$  as the least element of  $C_{\mathcal{N}, k}^{\beta_1, \beta_2}$ . Thus we have an  $i$  such that  $L_\alpha \preceq_k L_{\gamma_i}^{\mathcal{N}} \preceq_{k-1} L_{\beta_2}^{\mathcal{N}}$ .

Now  $L_{\gamma_i}^{\mathcal{N}}$  is  $\Sigma_{k-1}$  admissible by Lemma 3.4 as  $L_{\beta_2}^{\mathcal{N}}$  is  $\Sigma_{k-2}$  admissible by  $(\star_{k-1})$  while  $\alpha$  is (even)  $\Sigma_k$  admissible by our induction hypothesis and the previous Lemma and so  $(\star_k)(\alpha, \gamma_i)$  as required.  $\square$

**Lemma 5.12.** *The game  $G$  satisfies (1) and (2) of Lemma 1.8.*

PROOF: Since  $\alpha \in \mathcal{M} \models Th_{\alpha_n}$ ,  $\alpha$  cannot be  $n$ -admissible. We now know, by Lemma 5.10, that there are no  $\beta_1, \beta_2$  satisfying  $(\star_{n-1})$ . So, by Lemma 5.11, there is a  $k < n$  such that either  $(R_I(2+k))$  or  $(R_{II}(2+k))$  holds. Suppose  $(R_I(2+k))$  is the first condition that holds and that I wins the game. Since  $\forall i < 2+k$ , no condition  $(R_{::i})$  holds, there are  $\beta_1, \beta_2$  satisfying  $(\star_{k-1})$ . So,  $(R_I(2+k))$  implies that  $\mathcal{M}_{II}$  is not wellfounded. An analogous argument works if  $(R_{II}(2+k))$  is the first condition that holds and II wins the game.  $\square$

To complete the proof of Lemma 1.8 and so of our main result, Theorem 1.2, we need to show that we can get the same results with a  $\Pi_{3,n-1}^0$  game. As now defined, the game is clearly  $\Pi_{3,2n+4}^0$  as there are six sets at the beginning and then  $2(n-1)$  sets of the form  $R_I(2+k)$  or  $R_{II}(2+k)$  for  $k < n$ . So we need to modify the game a bit. The conditions  $(R_I0)$ ,  $(R_{II}0)$ ,  $(R_I1)$  and  $(R_{II}1)$  are  $\Sigma_2^0$  and so can be absorbed into some later  $\Pi_3^0$  condition as indicated below. Thus, we first prove that the rest of the game is  $\Pi_{3,n-1}^0$ . To decrease the number of alternations, we will change the order of the conditions. We need to make a couple observations: First, if no condition  $(R_{::i})$  for  $i < 2+k$  holds, then only the player playing a nonwellfounded model can loose because of a condition  $(R_{::(2+k)})$ . So it does not matter which of  $(R_I(2+k))$  and  $(R_{II}(2+k))$  comes first in the list and we can re-order this part of the list as:

$$\dots, (R_I3), \quad (R_I4), (R_I5), \quad (R_I6), \dots \\ (R_{II}3), (R_{II}4), \quad (R_{II}5), (R_{II}6),$$

Second, independently of the outcomes of  $(R_{::2})$ , only the player playing a nonwellfounded model can loose with a condition  $(R_{::3})$ , so it does not matter in which order we place these conditions either. We can now optimize the number of alternations of  $\Pi_3^0$  conditions as follows:

$$\dots, (R_I2), (R_I3), \quad (R_I4), (R_I5), \quad (R_I6), \dots \\ (R_{II}2), (R_{II}3), (R_{II}4), \quad (R_{II}5), (R_{II}6),$$

The first  $\Pi_3^0$  condition is now  $(R_I2) \vee (R_I3)$  which makes I win. We add in the early conditions that are at most  $\Sigma_2^0$  to make our desired first condition  $(R_I0) \vee (R_I1) \vee (R_I2) \vee (R_I3)$  which makes I win. The desired second condition is  $(R_{II}0) \vee (R_{II}1) \vee (R_{II}2) \vee (R_{II}3) \vee (R_{II}4)$  which makes II win. Then, for  $i > 1$ , the  $(2i-1)$ th condition is  $(R_I2i) \vee (R_I(2i+1))$  which makes I win, and the  $2i$ th condition is  $(R_{II}(2i+1)) \vee (R_{II}(2i+2))$  which makes II win. We end the list with  $(R_I(n+1))$  if  $n$  is even and with  $(R_{II}(n+1))$  if  $n$  is odd. This gives us  $n$  many conditions. We finish the description of the game by adding on the the full space  $2^\omega$  as our  $n$ th and last set. The idea here is that this prescribes a win for I when  $n$  is even (and so II failed to win at the end by providing a descending chain in  $\mathcal{M}_I$ ) and a win for II when  $n$  is odd (and so I failed to win at the end by providing a descending chain in  $\mathcal{M}_{II}$ ). This provision includes the conditions for a win required at the last step of our arguments and then enlarges the winning set (for one player) but remember that once we have passed through all the conditions on our list we no longer care how the game is defined. Thus, we

have a  $\Pi_{3,n-1}^0$  subset of  $2^\omega$  that defines our game as desired. This completes the proof of Lemma 1.8 and so of Theorem 1.2.  $\square$

## 6. NO REVERSALS FROM $\Sigma_4^1$ SENTENCES

In this section we will show that a true (or even consistent)  $\Sigma_4^1$  sentence  $T$  cannot imply  $\Delta_2^1\text{-CA}_0$  over  $\Pi_1^1\text{-CA}_0$  even for  $\beta$ -models. At higher levels ( $n \geq 2$ ), even with the help of  $\Pi_n^1\text{-CA}_0$  or  $\Delta_n^1\text{-CA}_0$  such a  $T$  does not imply  $\Delta_{n+1}^1\text{-CA}_0$  or  $\Pi_n^1\text{-CA}_0$ , respectively, even for  $\beta$ -models. We also supply analogs for  $\Sigma_m^1$  sentences that are theorems of ZFC (or at least consistent with  $\text{ZFC} + \text{V=L}(X)$  for some  $X \subseteq \omega$ ).

Our models for the failure of such implications will all be  $\beta$ -models with second order part of the form  $\mathbb{R} \cap \mathbb{L}_\delta$ . As before, we use the equivalences given by Simpson [2009, VII.5.3] between such models satisfying  $\Pi_{n+1}^1\text{-CA}_0$  or  $\Delta_{n+1}^1\text{-CA}_0$  and  $L_\delta$  being  $\Sigma_n$  nonprojectable or  $\Sigma_n$  admissible, respectively for  $n \geq 1$  or  $n \geq 2$ , respectively. The equivalence between  $\Sigma_1$  admissibility and  $\Delta_2^1\text{-CA}_0$  works when  $L_\delta$  is a limit of admissibles. In fact, as long as  $\delta$  is a limit of admissibles (i.e. closed under the operation of going to the next admissible) then, in  $L_\delta$ , any  $\Sigma_{n+1}^1$  sentence is equivalent to one that is  $\Sigma_n$  for  $n \geq 1$ . (This is essentially Simpson [2009, VII.3.24].) Also note that a  $\Sigma_n^1$  sentence is true in  $L_\delta$  if and only if it is true in (the  $\beta$ -model of second order arithmetic with sets in)  $\mathbb{R} \cap \mathbb{L}_\delta$ .

**Theorem 6.1.** *If  $T$  is a true  $\Pi_3^1$  sentence, then there is an ordinal  $\delta$  such that  $L_\delta \models T \ \& \ \forall \gamma \exists \beta > \gamma (\beta \text{ is admissible})$  but  $L_\delta$  is not  $\Sigma_1$  admissible and so  $\mathbb{R} \cap \mathbb{L}_\delta \not\models \Delta_2^1\text{-CA}_0$ .*

PROOF: First note that by Shoenfield absoluteness (for  $\Sigma_2^1$  formulas between  $V$  and  $L$ ), any true  $\Pi_3^1$  sentence is true in  $L$ . Suppose  $T$  is  $\forall A \exists B \forall C \Phi(A, B, C)$  where  $\Phi$  is arithmetic. Define functions  $f_T : \mathbb{R}^L \rightarrow ON$  and  $f_a : ON \rightarrow ON$  by  $f_T(A) = \mu \alpha (\exists B \in L_\alpha) \forall C \Phi(A, B, C)$  and  $f_a(\alpha) = \alpha^+$ , the least admissible greater than  $\alpha$ . For any ordinal  $\alpha$  closed under  $f_a$ ,  $\mathbb{R} \cap L_\alpha$  is a  $\beta$ -model of  $\Pi_1^1\text{-CA}$  Simpson [2009, VII.1.8]. If  $\alpha$  is also closed under  $f_T$  then  $L_\alpha \models T$  as well by the definition of a  $\beta$ -model (it is absolute for  $\Pi_1^1$  formulas). Thus there are  $\alpha$  such that  $L_\alpha \models T \ \& \ \forall \gamma \exists \beta > \gamma (\beta \text{ is admissible})$ . Let  $\delta_n$  be the  $n^{\text{th}}$  such ordinal and let  $\delta = \cup \delta_n$ . It is clear that  $L_\delta \models T \ \& \ \forall \gamma \exists \beta > \gamma (\beta \text{ is admissible})$ . The sequence  $\delta_n$  is clearly  $\Sigma_1$  over  $L_\delta$  and so  $L_\delta$  is not  $\Sigma_1$  admissible as required to have  $\mathbb{R} \cap \mathbb{L}_\delta \not\models \Delta_2^1\text{-CA}_0$ .  $\square$

Simpson pointed out that the above proof relativizes, of course, to any real  $X$  and so applies to  $\Sigma_4^1$  sentences by working in  $L(X)$  where  $X$  is the witness for the  $\Sigma_4^1$  sentence.

**Corollary 6.2.** *If  $T$  is a true  $\Sigma_4^1$  sentence (e.g. a theorem of ZFC) then  $T + \Pi_1^1\text{-CA} + \Pi_\infty^1\text{-TI} \not\models \Delta_2^1\text{-CA}_0$ .*

PROOF: Let  $\delta$  be defined in  $L(X)$  as it was on  $L$  in the Theorem but relativized to  $X$  where  $X$  is the witness to the first quantifier in  $T$ . As  $L_\delta(X)$  is closed under the next admissible,  $\mathbb{R} \cap L_\delta(X)$  is a  $\beta$ -model of  $\Pi_1^1\text{-CA}_0 + \Pi_\infty^1\text{-TI}_0$  as well as  $T$ .  $\square$

**Corollary 6.3.** *Borel-DET +  $\Pi_1^1\text{-CA} + \Pi_\infty^1\text{-TI} \not\models \Delta_2^1\text{-CA}_0$ .*

PROOF: Borel-DET is equivalent to a  $\Pi_3^1$  sentence over  $\Pi_1^1\text{-CA}_0$ . The sentence says that for every Borel code  $A$  there is a strategy  $f$  such that, for every  $C$ , the result of applying  $f$  to  $C$  is in the set coded by  $A$ . (The only thing to point out is that being a Borel code is  $\Pi_1^1$  and, assuming  $\Pi_1^1\text{-CA}_0$  (which actually follows in  $\text{RCA}_0$  from  $(\Sigma_1^0 \wedge \Pi_1^0)\text{-DET}$ ), membership in the set coded is  $\Delta_1^1$  by Simpson [2009, V.3.3-4]).  $\square$

Indeed, if a  $\Sigma_4^1$  sentence  $T$  is even consistent with ZFC then it is true in some model  $M$  of ZFC and so also in  $L^M(X)$  for some  $X \in M$  within which we can carry out the proof of Theorem 6.1. Thus all the proof theoretic nonimplications of Corollary 6.2 apply to such  $T$  as well.

An often used measuring rod for determinacy assumptions as in Martin [n.d.] and Welch [2009] is associating a class  $\Gamma$  with the least ordinal  $\delta$  such that  $L_\delta \models \Gamma\text{-DET}$ . The proof of Theorem 6.1 shows that, by the standards of admissibility, such ordinals are not themselves large. The following Proposition captures this version of our results. It relativizes to  $\Sigma_4^1$  sentences true in  $L$  (e.g. theorems of ZFC) although not literally to all true ones.

**Proposition 6.4.** *If  $T$  is a true  $\Pi_3^1$  sentence which implies  $\Pi_1^1\text{-CA}_0$  such as  $\Gamma\text{-DET}$  for any of the standard classes of the Borel hierarchy from  $\Sigma_1^0 \wedge \Pi_1^0$  up to the class of all Borel sets, then, the least  $\delta$  such that  $L \models \Gamma\text{-DET}$ , while a limit of admissibles, is not itself  $\Sigma_1^1$  admissible. Indeed, the  $\Delta_1$  projectum of  $\delta$  is  $\omega$ .*

**PROOF:** Let  $T$  be  $\forall x \subseteq \omega \exists y \subseteq \omega \forall z \subseteq \omega [\phi(x, y, z)]$ . First note that, in  $L_\delta$ , every  $\beta$  is countable. If not, then there is a least  $\beta < \delta$  which is not countable in  $L_\delta$ . In particular no new subsets of  $\omega$  appear between  $L_\beta$  and  $L_\delta$ . Thus any  $\Pi_\infty^1$  sentence true in  $L_\delta$  is true in  $L_\beta$  for a contradiction to the leastness of  $\delta$ . Thus every  $L_\beta$  for  $\beta < \delta$  is coded as a subset of  $\omega$  in  $L_\delta$  and so the closure of  $\mathbb{R} \cap \mathbb{L}_\delta$  under  $\Pi_1^1$  comprehension implies that  $L_\delta$  is closed under the next admissible and so is a limit of admissibles.

Next, consider the function  $f$  which takes a subset  $x$  of  $\omega$  in  $L_\delta$  to the least ordinal  $\gamma$  such that there is an  $\alpha < \gamma$  with  $\alpha^+ < \gamma$  as well and a subset  $y$  of  $\omega$  in  $L_\alpha$  such that  $L_{\alpha^+} \models \forall z \subseteq \omega [\phi(x, y, z)]$  (or equivalently  $\forall z \subseteq \omega [\phi(x, y, z)]$ ). As  $L_\delta \models T$  and is closed under the next admissible, this function is total on  $L_\delta$  and obviously  $\Delta_0$ . We can then define the total  $\Delta_0$  map  $h$  taking  $\nu$  to the least  $\gamma$  such that  $f(x) \in L_\gamma$  for every  $x \subseteq \omega$  in  $L_\nu$ . If  $\gamma_0 = \omega$  and  $\gamma_{n+1} = h(\gamma_n)$  then  $\gamma = \cup \gamma_n \leq \delta$ . It is clear that  $L_\gamma \models T$  and so  $\gamma = \delta$  which is therefore, not only not admissible but actually has a total  $\Sigma_1$  map from  $\omega$  cofinal in it. As every  $\beta < \delta$  is countable in  $L_\delta$ , these countings (of the  $\gamma_n$ ) can be combined with the cofinal map  $n \mapsto \gamma_n$  to produce the desired one-one  $\Sigma_1$  map from all of  $\omega$  onto  $L_\delta$ .  $\square$

We can actually say more than Theorem 6.1 about the reverse mathematical weakness of true  $\Pi_3^1$  (or  $\Sigma_4^1$ ) sentences. True  $\Pi_3^1$  sentences  $T$  don't help to prove higher levels of comprehension.

**Proposition 6.5.** *If  $T$  is a true  $\Pi_3^1$  sentence then, for each  $n \geq 1$ , there is an ordinal  $\alpha$  which is  $\Sigma_n$  admissible, a limit of  $\Sigma_n$  admissibles,  $\Sigma_n$  projectable and satisfies  $T$ .*

**PROOF:** Choose  $\alpha$  such that  $L_\alpha$  is the  $\Sigma_n$  Skolem hull of  $\omega$  in  $L_{\omega_1}$  under the standard  $\Sigma_n$  Skolem function. (Note that any  $\Sigma_n$  hull of  $L_{\omega_1}$  is transitive since every ordinal is countable in  $L_{\omega_1}$ .) Thus  $L_\alpha$  is a  $\Sigma_n$  elementary substructure of  $L_{\omega_1} \models T$ . It is  $\Sigma_n$  admissible by Lemma 3.4 and is the  $\Sigma_n$  Skolem hull of  $\omega$  in  $L_\alpha$  as well and so  $\Sigma_n$  projectable into  $\omega$ .

If  $n \geq 2$ , it is immediate that  $L_\alpha$  satisfies the  $\Pi_3^1$  formula  $T$ . For  $n = 1$  we have to be a bit more careful. Suppose  $T = \forall x \exists y \forall z \phi(x, y, z)$ . Consider any  $x \in L_\alpha$ .  $L_{\omega_1} \models \exists y (L_\delta \models \forall z \phi(x, y, z))$  where  $\delta$  is the second admissible after  $x$  and  $y$  appear in  $L$ . Thus there is such a  $y \in L_\alpha$ . To conclude, note that if it is not the case that  $\forall z \phi(x, y, z)$  then there would be a counterexample  $z$  inside  $L_\delta$  (actually definable over the first admissible after  $x$  and  $y$ ).  $\square$

**Corollary 6.6.** *If  $T$  is a true  $\Pi_3^1$  formula then, for  $n \geq 2$ ,  $\Delta_n^1\text{-CA} + T + \Pi_\infty^1\text{-TI} \not\vdash \Pi_n^1\text{-CA}_0$ .*

**Corollary 6.7.** *For  $n \geq 0$ ,  $\Delta_{n+2}^1\text{-CA} + n\text{-}\Pi_3^0\text{-DET} + \Pi_\infty^1\text{-TI} \not\vdash \Pi_{n+2}^1\text{-CA}_0$ .*

As before, the Proposition and its first Corollary hold for true  $\Sigma_4^1$  sentences by working in  $L(X)$  where  $X$  is the witness for the first existential quantifier.

We complete the picture of the powerlessness of true  $\Pi_3^1$  ( $\Sigma_4^1$ ) sentences in the sense of reverse mathematics presented in Corollaries 6.2 and 6.6 by showing that they do not help  $\Pi_n^1\text{-CA}$  prove  $\Delta_{n+1}^1\text{-CA}_0$  even for  $\beta$ -models.

**Proposition 6.8.** *If  $T$  is a true  $\Pi_3^1$  sentence then, for each  $n \geq 1$ , there is an ordinal  $\alpha$  which is  $\Sigma_n$  nonprojectable, satisfies  $T$  but is not  $\Sigma_{n+1}$  admissible.*

PROOF: Let  $\gamma_j$  be the  $j^{\text{th}}$  ordinal  $\delta$  such that  $L_\delta \preceq_n L_{\omega_1}$  and  $L_\delta \models T$ . Let  $\gamma = \cup \gamma_j$  and so  $L_{\gamma_j} \preceq_n L_\gamma \preceq_n L_{\omega_1}$  for each  $j$ . In addition,  $L_\gamma \models T$  as for any  $x \in L_\gamma$ ,  $x \in L_{\gamma_j}$  for some  $j$  and the witness for  $x$  verifying  $T$  in  $L_{\gamma_j}$  is one in  $L_\gamma$  since, as  $\Sigma_1$  elementary substructures of  $L_{\omega_1}$ , both are  $\beta$ -models. In particular,  $L_\gamma$  is  $\Sigma_n$  admissible by Lemma 3.4.

First note that  $L_\gamma$  is not  $\Sigma_n$  projectable. The point here is that if  $\beta < \gamma$  and  $\theta$  defines a  $\Sigma_n$  subset of  $\beta$  over  $L_\gamma$  then for some  $j$ ,  $\gamma_j$  is larger than  $\beta$  and all parameters in  $\theta$  are also in  $L_{\gamma_j}$ . For such a  $j$ , the  $\Sigma_n$  definition  $\theta$  defines the same subset of  $\beta$  over  $L_{\gamma_j}$  as over  $L_\gamma$ . The subset is then in  $L_\gamma$  as required.

Finally,  $\gamma$  is not  $\Sigma_{n+1}$  admissible as the sequence  $\gamma_j$  is definable by recursion over  $L_\gamma$  by a  $\Sigma_{n+1}$  formula.  $\gamma_j$  is the  $j^{\text{th}}$  ordinal  $\delta$  such that  $L_\delta \preceq_n L_\gamma$  (as  $L_\gamma \preceq_n L_{\omega_1}$  and so this condition is the same as  $L_\delta \preceq_n L_{\omega_1}$ ) and  $L_\delta \models T$ . To say that  $L_\delta \preceq_n L_\gamma$  is to say that for every  $c \in L_\delta$  and every  $\Sigma_n$  formula  $\phi(cx)$ ,  $L_\gamma \models \phi(c) \rightarrow L_\delta \models \phi(c)$ . By the  $\Sigma_n$  admissibility of  $L_\gamma$  this is equivalent to a  $\Pi_n$  formula and so its negation to a  $\Sigma_n$  one. Thus the sequence  $\gamma_j$  is  $\Sigma_{n+1}$  as required.  $\square$

**Corollary 6.9.** *If  $T$  is a true  $\Pi_3^1$  formula then, for  $n \geq 2$ ,  $\Pi_n^1\text{-CA} + T + \Pi_\infty^1\text{-TI} \not\vdash \Delta_{n+1}^1\text{-CA}_0$ .*

Once again this Proposition and Corollary generalize to true  $\Sigma_4^1$  sentences by relativization. More generally, the results of this section can be suitably generalized to  $\Sigma_m^1$  formulas  $T$  which are theorems of ZFC or at least consistent with  $\text{ZFC} + V=L(X)$  for some subset  $X$  of  $\omega$  as they are then true in a model  $M$  of  $V = L(X)$  and so in  $L_{(\omega_1)^M}(X)$ . The proofs are essentially the same beginning with the fact that  $T$  is true in  $L_{\omega_1}(X)$  (of a model  $M$ ).

**Corollary 6.10.** *If, for  $m > 4$ ,  $T$  is a  $\Sigma_m^1$  theorem of ZFC (or is even consistent with  $\text{ZFC} + V = L(X)$  for some subset  $X$  of  $\omega$ ) and  $n \geq m - 2$ , then  $\Pi_{n-1}^1\text{-CA} + T + \Pi_\infty^1\text{-TI}_0 \not\vdash \Delta_n^1\text{-CA}_0$  and  $\Delta_n^1\text{-CA} + T + \Pi_\infty^1\text{-TI}_0 \not\vdash \Pi_n^1\text{-CA}_0$ . The proofs follow those of Theorem 6.1 (with Corollary 6.2) and Proposition 6.5 (with Corollary 6.6), respectively.*

We close by pointing out that the results of this section are optimal in terms of the quantifier complexity of theorems that cannot imply  $\Pi_n^1\text{-CA}_0$  or even  $\Delta_n^1\text{-CA}_0$ .

*Remark 6.11.* For  $n \geq 1$ , there is a  $\Pi_{n+2}^1$  theorem  $T$  of second order arithmetic that implies  $\Pi_n^1\text{-CA}_0$  (and so  $\Delta_n^1\text{-CA}_0$ ) over  $\text{RCA}_0$ . The sentence  $T$  says that for every  $\Pi_n^1$  formula  $\forall Z\psi(n, Z, A)$  and for every choice of parameter  $A$ , there is an  $X$  such that  $\forall n(n \in X^{[0]} \rightarrow \forall Z\psi(n, Z, A)) \ \& \ \forall n(n \notin X^{[0]} \rightarrow \neg\psi(n, X^{[n+1]}, A))$ .

## REFERENCES

- [1] Barwise, J. [1975], *Admissible Sets and Structures, Perspectives in Mathematical Logic*, Springer-Verlag, Berlin.

- [2] Devlin, K. J. [1984], *Constructibility, Perspectives in Mathematical Logic*, Springer-Verlag, Berlin.
- [3] Foreman, M. ed. [2000], *Descriptive Set Theory and Dynamical Systems*, Cambridge University Press, Cambridge.
- [4] Friedman, H. [1971], Higher set theory and mathematical practice, *Annals of Mathematical Logic* **2**, 325-357.
- [5] Friedman, H. [1975], Some systems of second order arithmetic and their use, Proceedings of the *International Congress of Math.*, Vancouver 1974, v. 1, Can. Math. Congress, 1975, 235-242.
- [6] Kechris, A. [1995], *Classical Descriptive Set Theory, Graduate Texts in Mathematics* **156**, Springer-Verlag, New York.
- [7] Kechris, A. and Becker, H. [1996], *The Descriptive Set Theory of Polish Group Actions*, Cambridge University Press, Cambridge.
- [8] Kechris, A. and Louveau, A. [1987], *Descriptive Set Theory and the Structure of Sets of Uniqueness, London Math. Soc. Lect. Notes* **128**, Cambridge University Press, Cambridge.
- [9] Kechris, A. and Miller, B. D. [2004], *Topics in orbit equivalence, Lecture Notes in Mathematics* **1853**, Springer, New York.
- [10] Kuratowski, K. [1958], *Topologie* v.1, Państwowe Wydawnidwo Naukowe, Warsaw, English translation 1966.
- [11] Martin, D. A. [1974], Analysis  $\vdash \Delta_4^0$ -determinacy, circulated handwritten notes dated April 24, 1974.
- [12] Martin, D. A. [1974a],  $\Sigma_4^0$ -determinacy, circulated handwritten notes dated March 20, 1974.
- [13] Martin, D. A. [1975], Borel Determinacy, *Ann. Math.* **102**, 363-371.
- [14] Martin, D. A. [n.d.] *Determinacy*, circulated drafts, about 578 pp.
- [15] Martin, D. A and Steel, J. R. [1989], A proof of projective determinacy. *Journal of the American Mathematical Society* **2**, 71-125.
- [16] MedSalem, M. O. and Tanaka, K. [2007],  $\Delta_3^0$ -determinacy, comprehension and induction, *Journal of Symbolic Logic* **72**, 452-462.
- [17] MedSalem, M. O. and Tanaka, K. [2008], Weak determinacy and iterations of inductive definitions, in *Computational Prospects of Infinity, Part II: Presented Talks*, Chong, C., Feng, Q., Slaman, T. A., Woodin, W. H. and Yang, Y. eds., *LNS, Institute for Mathematical Sciences, National University of Singapore*, World Scientific, Singapore, 333-353.
- [18] Moschovakis, Y. N. [2009], *Descriptive Set Theory: Second Edition, Mathematical Surveys and Monographs* **155**, AMS, Providence.
- [19] Mummert, C. and Simpson, S. G. [2005], Reverse Mathematics and  $\Pi_2^1$  comprehension, *Bulletin of Symbolic Logic* **11**, 526-533.
- [20] Nemoto, T., MedSalem, M. O. and Tanaka, K. [2007], Infinite games in the Cantor space and subsystems of second order arithmetic, *Mathematical Logic Quarterly* **53**, 226-236.
- [21] Simpson, S. G. [2009], *Subsystems of Second Order Arithmetic* 2<sup>nd</sup> ed., *Perspectives in Logic*, ASL and Cambridge University Press, New York.
- [22] Steel, J. R. [1976], *Determinateness and Subsystems of Analysis*, Ph.D. Thesis, University of California, Berkeley.
- [23] Tanaka, K. [1990], Weak axioms of determinacy and subsystems of analysis I:  $\Delta_2^0$  games), *Zeitschr. f. math. Logik und Grundlagen d. Math.* **36**, 481-491.
- [24] Tanaka, K. [1991], Weak axioms of determinacy and subsystems of analysis II ( $\Sigma_2^0$  games), *Annals of Pure and Applied Logic* **52**, 181-193.
- [25] Welch, P. [2009], Weak systems of determinacy and arithmetical quasi-inductive definitions, preprints.
- [26] Woodin, H. [1988], Supercompact cardinals, sets of reals, and weakly homogeneous trees, *Proceedings of the National Academy of Sciences of the U.S.A.* **85**, 6587-6591.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CHICAGO, 5734 S. UNIVERSITY AVE., CHICAGO, IL 60637, USA

*E-mail address:* `antonio@math.uchicago.edu`

DEPARTMENT OF MATHEMATICS, MALOTT HALL, CORNELL UNIVERSITY, ITHACA NY 14853 USA

*E-mail address:* `shore@math.cornell.edu`

*URL:* `www.math.uchicago.edu/~antonio`

*URL:* `http://www.math.cornell.edu/~shore/`